

Have library access? [Log in through your library.](#)

[Register](#)

[Log in](#)

 [Workspace](#)

[Search](#) ▾

[About](#) [Support](#)



[All Content](#)

[Images](#)

Search journals, books, image 

[Browse](#) ▾

[Tools](#) ▾

JOURNAL ARTICLE

Generalized Likelihood Ratio Tests and Uniformly Most Powerful Tests

David Birkes

The American Statistician

Vol. 44, No. 2 (May, 1990), pp. 163-166 (4 pages)

Published By: Taylor & Francis, Ltd.



ITHAKA websites, which ITHAKA manages from its location in the United States, use cookies for different purposes, such as to ensure web site function, display non-targeted ads, provide social media features, and track usage, engaging with third party service providers such as Google Analytics. You may manage non-essential cookies in “Cookie Settings”. For more information, please see our [Cookie Policy](#).

[Cookie Settings](#)

[OK, proceed](#)

Preview

ITHAKA websites, which ITHAKA manages from its location in the United States, use cookies for different purposes, such as to ensure web site function, display non-targeted ads, provide social media features, and track usage, engaging with third party service providers such as Google Analytics. You may manage non-essential cookies in “Cookie Settings”. For more information, please see our [Cookie Policy](#).

[Cookie Settings](#)

OK, proceed

TEACHER'S CORNER

In this department *The American Statistician* publishes articles, reviews, and notes of interest to teachers of the first mathematical statistics course and of applied statistics courses. The department includes the Accent on Teaching Materials section; suitable contents for the section are described

under the section heading. Articles and notes for the department, but not intended specifically for the section, should be useful to a substantial number of teachers of the indicated types of courses or should have the potential for fundamentally affecting the way in which a course is taught.

Generalized Likelihood Ratio Tests and Uniformly Most Powerful Tests

DAVID BIRKES*

For testing a one-sided hypothesis in a one-parameter family of distributions, it is shown that the generalized likelihood ratio (GLR) test coincides with the uniformly most powerful (UMP) test, assuming certain monotonicity properties for the likelihood function. In particular, the equivalence of GLR tests and UMP tests holds for one-parameter exponential families. In addition, the relationship between GLR and UMPU (UMP unbiased) tests is considered when testing two-sided hypotheses.

KEY WORDS: Exponential family; Monotone likelihood ratio property; Unbiased test.

1. INTRODUCTION

A first course in mathematical statistics taught at the level of Mood, Graybill, and Boes (1974) or Bain and Engelhardt (1987) would present the concepts of a uniformly most powerful (UMP) test and of a generalized likelihood ratio (GLR) test. In this article a relationship between these two concepts is demonstrated.

When a GLR test is calculated for a hypothesis for which a UMP test exists, it seems that the GLR test almost always turns out to coincide with the UMP test [but see exercise 24.25 in Kendall and Stuart (1979)]. We show this to be true for tests of one-sided hypotheses in a one-parameter family of distributions under certain conditions on the likelihood function (see Theorem 2 in Sec. 2). In particular, the equivalence of GLR tests and UMP tests is seen to hold for one-parameter exponential families (see Sec. 3). Section 5 considers the relationship between GLR and UMPU (UMP unbiased) tests for two-sided hypotheses.

Suppose that X_1, \dots, X_n is a random sample from a distribution with density function $f(x; \theta)$ for some real-valued parameter θ in a parameter set Ω . The density func-

We will assume that there is a real-valued sufficient statistic $T(X_1, \dots, X_n)$. Because of the factorization criterion, likelihood procedures depend on the data only through the sufficient statistic. Rather than the likelihood function of the entire sample, we can use the likelihood function of T , which we will denote by $L(\theta; t)$.

Let us test the one-sided hypothesis $H_0: \theta \leq b$ versus $H_1: \theta > b$. Take any $\theta_1 > b$. The Neyman-Pearson lemma says that the most powerful test for testing the simple hypothesis $H_0: \theta = b$ versus the simple alternative $H_1: \theta = \theta_1$ is the test that rejects for large values of the likelihood ratio

$$R_s(t) = L(\theta_1; t)/L(b; t).$$

For testing the composite hypothesis $H_0: \theta \leq b$ versus $H_1: \theta > b$, the likelihood ratio can be generalized to the ratio

$$\sup_{\theta_1 \in \Omega_1} L(\theta_1; t) / \sup_{\theta_0 \in \Omega_0} L(\theta_0; t),$$

where $\Omega_0 = \{\theta \in \Omega: \theta \leq b\}$ and $\Omega_1 = \{\theta \in \Omega: \theta > b\}$. As a test statistic this ratio is essentially equivalent to the ratio

$$R_g(t) = \sup_{\theta \in \Omega} L(\theta; t) / \sup_{\theta_0 \in \Omega_0} L(\theta_0; t).$$

Letting $\hat{\theta}(t)$ denote the maximum likelihood estimate (MLE) of θ in the full model (parameterized by Ω) and letting $\hat{\theta}_0(t)$ denote the MLE of θ in the hypothesized model (parameterized by Ω_0), we can write

$$R_g(t) = L(\hat{\theta}(t); t) / L(\hat{\theta}_0(t); t).$$

The reciprocal of R_g is called the GLR. A test with rejection region of the form $\{t: R_g(t) > c\}$ or $\{t: R_g(t) \geq c\}$ is called a GLR test. [If $R_g(T)$ is a continuous random variable under all values of θ , then the rejection regions $\{t: R_g(t) > c\}$ and $\{t: R_g(t) \geq c\}$ are equivalent, since they

Abstract

ITHAKA websites, which ITHAKA manages from its location in the United States, use cookies for different purposes, such as to ensure web site function, display non-targeted ads, provide social media features, and track usage, engaging with third party service providers such as Google Analytics. You may manage non-essential cookies in "Cookie Settings". For more information, please see our [Cookie Policy](#).

[Cookie Settings](#)

OK, proceed

Publisher Information

Building on two centuries' experience, Taylor & Francis has grown rapidly over the last two decades to become a leading international academic publisher. The Group publishes over 800 journals and over 1,800 new books each year, covering a wide variety of subject areas and incorporating the journal...

Rights & Usage

This item is part of a JSTOR Collection.
For terms and use, please refer to our [Terms and Conditions](#)
The American Statistician © 1990 [American Statistical Association](#)
[Request Permissions](#)

Explore JSTOR

- By Subject
- By Title
- Collections
- Publisher
- Advanced Search
- Images
- Data for Research
- Get Access
- Get Support
- LibGuides
- Research Basics

ITHAKA websites, which ITHAKA manages from its location in the United States, use cookies for different purposes, such as to ensure web site function, display non-targeted ads, provide social media features, and track usage, engaging with third party service providers such as Google Analytics. You may manage non-essential cookies in “Cookie Settings”. For more information, please see our [Cookie Policy](#).

[Cookie Settings](#)

OK, proceed

For Publishers

JSTOR is part of ITHAKA, a not-for-profit organization helping the academic community use digital technologies to preserve the scholarly record and to advance research and teaching in sustainable ways.

©2000-2022 ITHAKA. All Rights Reserved. JSTOR®, the JSTOR logo, JPASS®, Artstor®, Reveal Digital™ and ITHAKA® are registered trademarks of ITHAKA.

[Terms & Conditions of Use](#)

[Privacy Policy](#) [Accessibility](#)

[Cookie Policy](#) [Cookie Settings](#)

[Select Language](#)

ITHAKA websites, which ITHAKA manages from its location in the United States, use cookies for different purposes, such as to ensure web site function, display non-targeted ads, provide social media features, and track usage, engaging with third party service providers such as Google Analytics. You may manage non-essential cookies in “Cookie Settings”. For more information, please see our [Cookie Policy](#).

[Cookie Settings](#)

OK, proceed