

Houghton Mifflin Harcourt (http://www.hmhco.com)

Sign In (/account/Sign-In-Register?ReturnUrl=%2fstudy-guides%2falgebra%2flinear-algebra%2freal-euclidean-vector-spaces%2fthenullspace-of-a-matrix)

Sign Up (/account/Sign-In-Register?ReturnUrl=%2fstudy-guides%2falgebra%2flinear-algebra%2freal-euclidean-vectorspaces%2fthe-nullspace-of-a-matrix#signup)



<u>(/)</u>

Search CliffsNotes

Q

Literature Notes (/literature)Test Prep (/test-prep)Study Guides (/study-guides)Student Life (/student-life)





The Nullspace of a Matrix


The Nullspace of a Matrix

<u>Vector Algebra</u> The Space R² (/study-quides/algebra/linear-algebra/vectoralgebra/the-snace-r2) Matrix Algebra Matrices (/study-guides/algebra/linear-algebra/matrix-The SpaceR+ (/study-guides/algebra/linear-algebra/vector-algebra/matrices) Linear Systems Operstion & With Matricest 4054-uply desides designed by a designed by algebra/the-space-rn) Real Euclidean Meichant Senatoristation Roan Operation is tudy Deptedent / Dep aldelessate especial contractions and a second contraction of the cont **sper**ations-to-determine-a1) The Determinan ramer's Rule (/study-quides/algebra/linear-algebra/thethitteen in the produce of the continue of the electioned in very stearce/st/limetare indented interared vstems) Eigenvalues and Eigenvalues an **Soliele State of Sallitations (distribute state of Sallitation Sallitation State of Sallitat** The Nulls State of the 1015 project the the eigenvectors/determining-the-eigenvectors-of-a-matrix) The solution Explaction xpratros io Static Static Planting interest in the solution Explanting interest in the solution in the solution is a set of homographic interest in the solution in the solution is a set of homographic interest in the solution in the solution is a set of homographic interest in the solution in the solution is a set of homographic interest in the solution in the solution is a set of homographic interest in the solution in the solution is a set of homographic interest in the solution in the solution is a set of homographic interest in the solution in the solution is a set of homographic interest in the solution in the solution is a set of homographic interest in the solution in the solution is a set of homographic interest in the solution in the solution is a set of homographic interest in the solution in the solution is a set of homographic interest in the solution in the solution is a set of homographic interest in the solution in the solution is a set of homographic interest in the solution in the s burce of vector spaces. Let A be an m by n matrix, and consider the months in the consideration of the con Row Space and Column Space of a Matrix (/study-Diagonalization (/study-guides/algebra/linear-algebra/eigenvalues-Ax = 0and-eigenvectors/diagonalization) Eigenvalue and Eigenvector Defined (/study-guides/algebra/linearalgebra/eigenvalues-and-eigenvectors/eigenvalue-and-eigenvectordefined)

Since A is m by n, the set of all vectors \mathbf{x} which satisfy this equation forms a subset of \mathbf{R}^n . (This subset is nonempty, since it clearly contains the zero vector: $\mathbf{x} = \mathbf{0}$ always satisfies $A \mathbf{x} = \mathbf{0}$.) This subset actually forms a subspace of \mathbf{R}^n , called the **nullspace** of the

matrix A and denoted N(A). To prove that N(A) is a subspace of \mathbf{R}^n , closure under both addition and scalar multiplication must be established. If \mathbf{x}_1 and \mathbf{x}_2 are in N(A), then, by definition, $A\mathbf{x}_1 = \mathbf{0}$ and $A\mathbf{x}_2 = \mathbf{0}$. Adding these equations yields

$$A\mathbf{x}_1 + A\mathbf{x}_2 = \mathbf{0} \implies A(\mathbf{x}_1 + \mathbf{x}_2) = \mathbf{0} \implies \mathbf{x}_1 + \mathbf{x}_2 \in N(A)$$



which verifies closure under addition. Next, if \mathbf{x} is in N(A), then $A \mathbf{x} = \mathbf{0}$, so if k is any scalar,

$$k(A\mathbf{x}) = \mathbf{0} \implies A(k\mathbf{x}) = \mathbf{0} \implies k\mathbf{x} \in N(A)$$

verifying closure under scalar multiplication. Thus, the solution set of a homogeneous linear system forms a vector space. Note carefully that if the

system is *not* homogeneous, then the set of solutions is *not* a vector space since the set will not contain the zero vector.

Example 1: The plane P in Example 7, given by 2x + y - 3z = 0, was shown to be a subspace of \mathbb{R}^3 . Another proof that this defines a subspace of \mathbb{R}^3 follows from the observation that 2x + y - 3z = 0 is equivalent to the homogeneous system

$$A\begin{bmatrix} x \\ y \\ z \end{bmatrix} = 0$$

where A is the 1 x 3 matrix [2 1 -3]. P is the nullspace of A.

Fxamnle 2. The set of solutions of the homogeneous system

Example 2. The out of oblations of the homogeneous system

$$\begin{bmatrix} -1 & 1 & 2 & 4 \\ 2 & 0 & 1 & -7 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

forms a subspace of \mathbb{R}^n for some n. State the value of n and explicitly determine this subspace.

Since the coefficient matrix is 2 by 4, \mathbf{x} must be a 4-vector. Thus, n = 4: The nullspace of this matrix is a subspace of \mathbf{R}^4 . To determine this subspace, the equation is solved by first row-reducing the given matrix:

$$\begin{bmatrix} -1 & 1 & 2 & 4 \\ 2 & 0 & 1 & -7 \end{bmatrix} \xrightarrow{\begin{array}{c} 2r_1 \text{ added to } r_2 \\ -(-1)r_1 \end{array}} \begin{bmatrix} 1 & -1 & -2 & -4 \\ 0 & 2 & 5 & 1 \end{bmatrix}$$

Therefore, the system is equivalent to

$$\begin{bmatrix} 1 & -1 & -2 & -4 \\ 0 & 2 & 5 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

that is,

$$x_1 - x_2 - 2x_3 - 4x_4 = 0$$
$$2x_2 + 5x_3 + x_4 = 0$$

If you let x_3 and x_4 be free variables, the second equation directly above implies

$$x_2 = -\frac{1}{2}(5x_3 + x_4)$$

Substituting this result into the other equation determines x_1 :

$$x_1 - \left[-\frac{1}{2}(5x_3 + x_4)\right] - 2x_3 - 4x_4 = 0$$
$$x_1 = -\frac{1}{2}(x_3 - 7x_4)$$

Therefore, the set of solutions of the given homogeneous system can be written as

$$\begin{bmatrix}
-\frac{1}{2}(x_3 - 7x_4) \\
-\frac{1}{2}(5x_3 + x_4) \\
x_3 \\
x_4
\end{bmatrix} : x_3, x_4 \in \mathbb{R}$$

which is a subspace of ${\bf R}^4$. This is the nullspace of the matrix

$$\begin{bmatrix} -1 & 1 & 2 & 4 \\ 2 & 0 & 1 & -7 \end{bmatrix}$$

Example 3: Find the nullspace of the matrix

$$A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$$

By definition, the nullspace of A consists of all vectors \mathbf{x} such that A \mathbf{x} = $\mathbf{0}$. Perform the following elementary row operations on A,

$$\begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \xrightarrow{\mathbf{r}_1 \leftrightarrow \mathbf{r}_2} \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \xrightarrow{-2\mathbf{r}_1 \text{ added to } \mathbf{r}_2} \begin{bmatrix} 1 & 2 \\ 0 & -3 \end{bmatrix}$$

to conclude that $A \mathbf{x} = \mathbf{0}$ is equivalent to the simpler system

$$\begin{bmatrix} 1 & 2 \\ 0 & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The second row implies that $x_2 = 0$, and back-substituting this into the first row implies that $x_1 = 0$ also. Since the only solution of $A \mathbf{x} = \mathbf{0}$ is $\mathbf{x} = \mathbf{0}$, the nullspace of A consists of the zero vector alone. This subspace, $\{\mathbf{0}\}$, is called the **trivial subspace** (of \mathbf{R}^2).

Example 4: Find the nullspace of the matrix

$$B = \begin{bmatrix} 2 & 1 \\ -4 & -2 \end{bmatrix}$$

To solve $B \mathbf{x} = \mathbf{0}$, begin by row-reducing B:

$$\begin{bmatrix} 2 & 1 \\ -4 & -2 \end{bmatrix} \xrightarrow{2r_1 \text{ added to } r_2} \begin{bmatrix} 2 & 1 \\ 0 & 0 \end{bmatrix}$$

The system $B \mathbf{x} = \mathbf{0}$ is therefore equivalent to the simpler system

$$\begin{bmatrix} 2 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Since the bottom row of this coefficient matrix contains only zeros, x_2 can be taken as a free variable. The first row then gives $2x_1+x_2=0 \Rightarrow x_1=-\frac{1}{2}x_2$ so any vector of the form

$$\begin{bmatrix} -\frac{1}{2}x_2 \\ x_2 \end{bmatrix}$$

satisfies $B \mathbf{x} = \mathbf{0}$. The collection of all such vectors is the nullspace of B, a subspace of \mathbf{R}^2 :

$$N(B) = \left\{ \begin{bmatrix} -\frac{1}{2}x \\ x \end{bmatrix} : x \in \mathbb{R} \right\}$$

Notes

CliffsNotes study guides are written by real teachers and professors, so no matter what you're studying, CliffsNotes can ease your homework headaches and help you score high on exams.

About CliffsNotes (/discover-about)

Advertise with Us (/discover-advertise)

(https://instagram.com/cliffsnotesofficial/)

Contact Us (/discover-contact)

Follow us:





(https://www.youtube.com/user/cliffsnotes/videos)

© 2016 Houghton Mifflin Harcourt. All rights reserved.

Terms & Conditions (http://www.hmhco.com/common/terms-conditions)

Privacy Policy (http://www.hmhco.com/common/privacy-policy)

Disclaimer (/disclaimer)

Sitemap (/sitemap)

A18ACD436D5A3997E3DA2573E3FD792A