

$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$

$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$

We wish to estimate:

$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$

We wish to estimate:

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\}$$

$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$


We wish to estimate:

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} = \mathbf{P}\{|S_n - np| > n\epsilon\}$$

$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$

We wish to estimate:


$$\mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} = \mathbf{P}\{|S_n - np| > n\epsilon\}$$



$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$

We wish to estimate:

$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} &= \mathbf{P}\{|S_n - np| > n\epsilon\} \\ &= \sum_{k: |k - np| > n\epsilon} b_n(k; p) \end{aligned}$$




$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$

We wish to estimate:

$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} &= \mathbf{P}\{|S_n - np| > n\epsilon\} \\ &= \sum_{k: |k - np| > n\epsilon} b_n(k; p) \end{aligned}$$

This means: sum over all integers k such that $|k - np| > n\epsilon$.
(That is to say, either $0 \leq k < np - n\epsilon$ or $np + n\epsilon < k \leq n$.)



$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$


We wish to estimate:

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} = \mathbf{P}\{|S_n - np| > n\epsilon\}$$

$$= \sum_{k: |k - np| > n\epsilon} b_n(k; p)$$

$$= \sum_{k: |k - np| > n\epsilon} 1 \cdot b_n(k; p)$$

This means: sum over all integers k such that $|k - np| > n\epsilon$.
(That is to say, either $0 \leq k < np - n\epsilon$ or $np + n\epsilon < k \leq n$.)




$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$

We wish to estimate:

$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} &= \mathbf{P}\{|S_n - np| > n\epsilon\} \\ &= \sum_{k: |k - np| > n\epsilon} b_n(k; p) \\ &= \sum_{k: |k - np| > n\epsilon} 1 \cdot b_n(k; p) \end{aligned}$$

This means: sum over all integers k such that $|k - np| > n\epsilon$.
(That is to say, either $0 \leq k < np - n\epsilon$ or $np + n\epsilon < k \leq n$.)

Which means: $\frac{|k - np|}{n\epsilon} > 1$.



$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$


We wish to estimate:

$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} &= \mathbf{P}\{|S_n - np| > n\epsilon\} \\ &= \sum_{k: |k - np| > n\epsilon} b_n(k; p) \\ &= \sum_{k: |k - np| > n\epsilon} 1 \cdot b_n(k; p) \end{aligned}$$

This means: sum over all integers k such that $|k - np| > n\epsilon$.
(That is to say, either $0 \leq k < np - n\epsilon$ or $np + n\epsilon < k \leq n$.)

Which means: $\frac{|k - np|}{n\epsilon} > 1$.

Or, equivalently: $\frac{(k - np)^2}{n^2 \epsilon^2} > 1$.



$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$


We wish to estimate:

$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} &= \mathbf{P}\{|S_n - np| > n\epsilon\} \\ &= \sum_{k: |k - np| > n\epsilon} b_n(k; p) \\ &= \sum_{k: |k - np| > n\epsilon} 1 \cdot b_n(k; p) \\ &\leq \sum_{k: |k - np| > n\epsilon} \frac{(k - np)^2}{n^2 \epsilon^2} \cdot b_n(k; p) \end{aligned}$$

This means: sum over all integers k such that $|k - np| > n\epsilon$.
(That is to say, either $0 \leq k < np - n\epsilon$ or $np + n\epsilon < k \leq n$.)

Which means: $\frac{|k - np|}{n\epsilon} > 1$.

Or, equivalently: $\frac{(k - np)^2}{n^2 \epsilon^2} > 1$.



$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$


We wish to estimate:

$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} &= \mathbf{P}\{|S_n - np| > n\epsilon\} \\ &= \sum_{k: |k - np| > n\epsilon} b_n(k; p) \\ &= \sum_{k: |k - np| > n\epsilon} 1 \cdot b_n(k; p) \\ &\leq \sum_{k: |k - np| > n\epsilon} \frac{(k - np)^2}{n^2 \epsilon^2} \cdot b_n(k; p) \\ &\leq \sum_{k=0}^n \frac{(k - np)^2}{n^2 \epsilon^2} \cdot b_n(k; p) \end{aligned}$$

This means: sum over all integers k such that $|k - np| > n\epsilon$.
(That is to say, either $0 \leq k < np - n\epsilon$ or $np + n\epsilon < k \leq n$.)

Which means: $\frac{|k - np|}{n\epsilon} > 1$.


Or, equivalently: $\frac{(k - np)^2}{n^2 \epsilon^2} > 1$.



$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$

We wish to estimate:

$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} &= \mathbf{P}\{|S_n - np| > n\epsilon\} \\ &= \sum_{k: |k - np| > n\epsilon} b_n(k; p) \\ &= \sum_{k: |k - np| > n\epsilon} 1 \cdot b_n(k; p) \\ &\leq \sum_{k: |k - np| > n\epsilon} \frac{(k - np)^2}{n^2 \epsilon^2} \cdot b_n(k; p) \\ &\leq \sum_{k=0}^n \frac{(k - np)^2}{n^2 \epsilon^2} \cdot b_n(k; p) \\ &= \frac{1}{n^2 \epsilon^2} \sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) \end{aligned}$$



This means: sum over all integers k such that $|k - np| > n\epsilon$.
(That is to say, either $0 \leq k < np - n\epsilon$ or $np + n\epsilon < k \leq n$.)


Which means: $\frac{|k - np|}{n\epsilon} > 1$.

Or, equivalently: $\frac{(k - np)^2}{n^2 \epsilon^2} > 1$.

$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$

We wish to estimate:

$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} &= \mathbf{P}\{|S_n - np| > n\epsilon\} \\ &= \sum_{k: |k - np| > n\epsilon} b_n(k; p) \\ &= \sum_{k: |k - np| > n\epsilon} 1 \cdot b_n(k; p) \\ &\leq \sum_{k: |k - np| > n\epsilon} \frac{(k - np)^2}{n^2 \epsilon^2} \cdot b_n(k; p) \\ &\leq \sum_{k=0}^n \frac{(k - np)^2}{n^2 \epsilon^2} \cdot b_n(k; p) \\ &= \frac{1}{n^2 \epsilon^2} \sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) \\ &= \frac{1}{n^2 \epsilon^2} \cdot npq = \frac{p(1 - p)}{n\epsilon^2} \end{aligned}$$



This means: sum over all integers k such that $|k - np| > n\epsilon$.
(That is to say, either $0 \leq k < np - n\epsilon$ or $np + n\epsilon < k \leq n$.)


Which means: $\frac{|k - np|}{n\epsilon} > 1$.

Or, equivalently: $\frac{(k - np)^2}{n^2 \epsilon^2} > 1$.

$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$

We wish to estimate:


$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} &= \mathbf{P}\{|S_n - np| > n\epsilon\} \\ &= \sum_{k: |k - np| > n\epsilon} b_n(k; p) \\ &= \sum_{k: |k - np| > n\epsilon} 1 \cdot b_n(k; p) \\ &\leq \sum_{k: |k - np| > n\epsilon} \frac{(k - np)^2}{n^2 \epsilon^2} \cdot b_n(k; p) \\ &\leq \sum_{k=0}^n \frac{(k - np)^2}{n^2 \epsilon^2} \cdot b_n(k; p) \\ &= \frac{1}{n^2 \epsilon^2} \sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) \\ &= \frac{1}{n^2 \epsilon^2} \cdot npq = \frac{p(1 - p)}{n\epsilon^2} \end{aligned}$$



This means: sum over all integers k such that $|k - np| > n\epsilon$.
(That is to say, either $0 \leq k < np - n\epsilon$ or $np + n\epsilon < k \leq n$.)

Which means: $\frac{|k - np|}{n\epsilon} > 1$.


Or, equivalently: $\frac{(k - np)^2}{n^2 \epsilon^2} > 1$.



$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$

We wish to estimate:


$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} &= \mathbf{P}\{|S_n - np| > n\epsilon\} \\ &= \sum_{k: |k - np| > n\epsilon} b_n(k; p) \\ &= \sum_{k: |k - np| > n\epsilon} 1 \cdot b_n(k; p) \\ &\leq \sum_{k: |k - np| > n\epsilon} \frac{(k - np)^2}{n^2 \epsilon^2} \cdot b_n(k; p) \\ &\leq \sum_{k=0}^n \frac{(k - np)^2}{n^2 \epsilon^2} \cdot b_n(k; p) \\ &= \frac{1}{n^2 \epsilon^2} \sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) \\ &= \frac{1}{n^2 \epsilon^2} \cdot npq = \frac{p(1 - p)}{n\epsilon^2} \leq \frac{1}{4n\epsilon^2} \end{aligned}$$



This means: sum over all integers k such that $|k - np| > n\epsilon$.
(That is to say, either $0 \leq k < np - n\epsilon$ or $np + n\epsilon < k \leq n$.)

Which means: $\frac{|k - np|}{n\epsilon} > 1$.

Or, equivalently: $\frac{(k - np)^2}{n^2 \epsilon^2} > 1$.




$$\sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) = npq$$

We wish to estimate:

$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - p\right| > \epsilon\right\} &= \mathbf{P}\{|S_n - np| > n\epsilon\} \\ &= \sum_{k: |k - np| > n\epsilon} b_n(k; p) \\ &= \sum_{k: |k - np| > n\epsilon} 1 \cdot b_n(k; p) \\ &\leq \sum_{k: |k - np| > n\epsilon} \frac{(k - np)^2}{n^2 \epsilon^2} \cdot b_n(k; p) \\ &\leq \sum_{k=0}^n \frac{(k - np)^2}{n^2 \epsilon^2} \cdot b_n(k; p) \\ &= \frac{1}{n^2 \epsilon^2} \sum_{k=0}^n (k - np)^2 \cdot b_n(k; p) \\ &= \frac{1}{n^2 \epsilon^2} \cdot npq = \frac{p(1-p)}{n\epsilon^2} \leq \frac{1}{4n\epsilon^2} \end{aligned}$$

p has vanished!



This means: sum over all integers k such that $|k - np| > n\epsilon$.
(That is to say, either $0 \leq k < np - n\epsilon$ or $np + n\epsilon < k \leq n$.)

Which means: $\frac{|k - np|}{n\epsilon} > 1$.

Or, equivalently: $\frac{(k - np)^2}{n^2 \epsilon^2} > 1$.

