



The distribution of accumulated successes



Is $\frac{S_n}{n}$ a good approximation to p ?

The distribution of accumulated successes




Is $\frac{S_n}{n}$ a good approximation to p ?

Three Bernoulli trials, success probability p , failure probability $q = 1 - p$:

$$X_1, X_2, X_3$$

The distribution of accumulated successes




Is $\frac{S_n}{n}$ a good approximation to p ?

Three Bernoulli trials, success probability p , failure probability $q = 1 - p$:
 X_1, X_2, X_3

Atoms	0,0,0	0,0,1	0,1,0	0,1,1	1,0,0	1,0,1	1,1,0	1,1,1
Probabilities	q^3	q^2p	q^2p	qp^2	q^2p	qp^2	qp^2	p^3

Chances multiply under independent selection



The distribution of accumulated successes

Is $\frac{S_n}{n}$ a good approximation to p ?

Three Bernoulli trials, success probability p , failure probability $q = 1 - p$:


$$X_1, X_2, X_3$$

Atoms	0,0,0	0,0,1	0,1,0	0,1,1	1,0,0	1,0,1	1,1,0	1,1,1
Probabilities	q^3	q^2p	q^2p	qp^2	q^2p	qp^2	qp^2	p^3

Chances multiply under independent selection

Accumulated successes: $S_3 = X_1 + X_2 + X_3$

Inherited atomic probabilities: $b(k) := \mathbf{P}\{S_3 = k\}$



The distribution of accumulated successes

Is $\frac{S_n}{n}$ a good approximation to p ?

Three Bernoulli trials, success probability p , failure probability $q = 1 - p$:

$$X_1, X_2, X_3$$

Atoms	0,0,0	0,0,1	0,1,0	0,1,1	1,0,0	1,0,1	1,1,0	1,1,1
Probabilities	q^3	q^2p	q^2p	qp^2	q^2p	qp^2	qp^2	p^3


Chances multiply under independent selection

Accumulated successes: $S_3 = X_1 + X_2 + X_3$

Inherited atomic probabilities: $b(k) := \mathbf{P}\{S_3 = k\}$

k	0	1	2	3
---	---	---	---	---

The distribution of accumulated successes



Is $\frac{S_n}{n}$ a good approximation to p ?

Three Bernoulli trials, success probability p , failure probability $q = 1 - p$:

$$X_1, X_2, X_3$$

Atoms	0,0,0	0,0,1	0,1,0	0,1,1	1,0,0	1,0,1	1,1,0	1,1,1
Probabilities	q^3	q^2p	q^2p	qp^2	q^2p	qp^2	qp^2	p^3


Chances multiply under independent selection

Accumulated successes: $S_3 = X_1 + X_2 + X_3$

Inherited atomic probabilities: $b(k) := \mathbf{P}\{S_3 = k\}$

k	0	1	2	3
---	---	---	---	---

The distribution of accumulated successes



Is $\frac{S_n}{n}$ a good approximation to p ?

Three Bernoulli trials, success probability p , failure probability $q = 1 - p$:

$$X_1, X_2, X_3$$

Atoms	0,0,0	0,0,1	0,1,0	0,1,1	1,0,0	1,0,1	1,1,0	1,1,1
Probabilities	q^3	q^2p	q^2p	qp^2	q^2p	qp^2	qp^2	p^3


Chances multiply under independent selection

Accumulated successes: $S_3 = X_1 + X_2 + X_3$

Inherited atomic probabilities: $b(k) := \mathbf{P}\{S_3 = k\}$

k	0	1	2	3
---	---	---	---	---

The distribution of accumulated successes



Is $\frac{S_n}{n}$ a good approximation to p ?

Three Bernoulli trials, success probability p , failure probability $q = 1 - p$:

$$X_1, X_2, X_3$$


Atoms	0,0,0	0,0,1	0,1,0	0,1,1	1,0,0	1,0,1	1,1,0	1,1,1
Probabilities	q^3	q^2p	q^2p	qp^2	q^2p	qp^2	qp^2	p^3

Chances multiply under independent selection

Accumulated successes: $S_3 = X_1 + X_2 + X_3$

Inherited atomic probabilities: $b(k) := \mathbf{P}\{S_3 = k\}$

k	0	1	2	3
---	---	---	---	---



The distribution of accumulated successes

Is $\frac{S_n}{n}$ a good approximation to p ?

Three Bernoulli trials, success probability p , failure probability $q = 1 - p$:
 X_1, X_2, X_3

Atoms	0,0,0	0,0,1	0,1,0	0,1,1	1,0,0	1,0,1	1,1,0	1,1,1
Probabilities	q^3	q^2p	q^2p	qp^2	q^2p	qp^2	qp^2	p^3

Chances multiply under independent selection

Accumulated successes: $S_3 = X_1 + X_2 + X_3$

Inherited atomic probabilities: $b(k) := \mathbf{P}\{S_3 = k\}$

k	0	1	2	3
$b(k)$	q^3	$3q^2p$	$3qp^2$	p^3

k	0	1	2	3
b(k)	q^3	$3q^2p$	$3qp^2$	p^3

The sample: a sequence of Bernoulli trials X_1, X_2, \dots, X_n

Accumulated successes: $S_n = X_1 + X_2 + \dots + X_n$

k	0	1	2	3
b(k)	q^3	$3q^2p$	$3qp^2$	p^3

Independent; success probability p , failure probability $q = 1 - p$

The sample: a sequence of Bernoulli trials X_1, X_2, \dots, X_n


Accumulated successes: $S_n = X_1 + X_2 + \dots + X_n$

k	0	1	2	3
b(k)	q^3	$3q^2p$	$3qp^2$	p^3

Independent; success probability p , failure probability $q = 1 - p$

The sample: a sequence of Bernoulli trials X_1, X_2, \dots, X_n

Accumulated successes: $S_n = X_1 + X_2 + \dots + X_n$



k	0	1	2	3
b(k)	q^3	$3q^2p$	$3qp^2$	p^3

Independent; success probability p , failure probability $q = 1 - p$

The sample: a sequence of Bernoulli trials X_1, X_2, \dots, X_n

Accumulated successes: $S_n = X_1 + X_2 + \dots + X_n$



Probability of observing a particular sequence of 4 successes in 12 trials: p^4q^{12-4}

k	0	1	2	3
b(k)	q^3	$3q^2p$	$3qp^2$	p^3

Independent; success probability p , failure probability $q = 1 - p$

The sample: a sequence of Bernoulli trials X_1, X_2, \dots, X_n

Accumulated successes: $S_n = X_1 + X_2 + \dots + X_n$



Probability of observing a particular sequence of 4 successes in 12 trials: $p^4 q^{12-4}$

Probability of observing exactly 4 successes in 12 trials: $\binom{12}{4} p^4 q^{12-4}$

k	0	1	2	3
b(k)	q^3	$3q^2p$	$3qp^2$	p^3

Independent; success probability p , failure probability $q = 1 - p$

The sample: a sequence of Bernoulli trials X_1, X_2, \dots, X_n

Accumulated successes: $S_n = X_1 + X_2 + \dots + X_n$



Probability of observing a particular sequence of k successes in n trials: $p^k q^{n-k}$

Probability of observing exactly 4 successes in 12 trials: $\binom{12}{4} p^4 q^{12-4}$

k	0	1	2	3
b(k)	q^3	$3q^2p$	$3qp^2$	p^3

Independent; success probability p , failure probability $q = 1 - p$

The sample: a sequence of Bernoulli trials X_1, X_2, \dots, X_n

Accumulated successes: $S_n = X_1 + X_2 + \dots + X_n$



Probability of observing a particular sequence of k successes in n trials: $p^k q^{n-k}$

Probability of observing exactly k successes in n trials: $\binom{n}{k} p^k q^{n-k}$

k	0	1	2	3
b(k)	q^3	$3q^2p$	$3qp^2$	p^3

Independent; success probability p , failure probability $q = 1 - p$

The sample: a sequence of Bernoulli trials X_1, X_2, \dots, X_n

Accumulated successes: $S_n = X_1 + X_2 + \dots + X_n$



Probability of observing a particular sequence of k successes in n trials: $p^k q^{n-k}$

Probability of observing exactly k successes in n trials: $\binom{n}{k} p^k q^{n-k} =: b(k) = b_n(k; p)$

The binomial distribution

- ❖ Accumulated successes: New chance variables from old.
 - ❖ The (new) sample space: $\Omega = \{0, 1, \dots, n\}$.
 - ❖ The probability measure is induced from the atomic probabilities: $b(k) = b_n(k; p) := \mathbf{P}\{S_n = k\}$.

The binomial distribution

- ❖ Accumulated successes: New chance variables from old.

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p)$$

$$S_n = X_1 + X_2 + \dots + X_n \sim \text{Binomial}(n, p)$$

- ❖ The (new) sample space: $\Omega = \{0, 1, \dots, n\}$.
- ❖ The probability measure is induced from the atomic probabilities: $b(k) = b_n(k; p) := \mathbf{P}\{S_n = k\}$.

The binomial distribution

- ❖ Accumulated successes: New chance variables from old.

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p)$$

$$S_n = X_1 + X_2 + \dots + X_n \sim \text{Binomial}(n, p)$$

- ❖ The (new) sample space: $\Omega = \{0, 1, \dots, n\}$.
- ❖ The probability measure is induced from the atomic probabilities: $b(k) = b_n(k; p) := \mathbf{P}\{S_n = k\}$.

$$b(k) = \binom{n}{k} p^k q^{n-k} \quad (k = 0, 1, \dots, n)$$

The binomial distribution

- ❖ Accumulated successes: New chance variables from old.

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p)$$

$$S_n = X_1 + X_2 + \dots + X_n \sim \text{Binomial}(n, p)$$

- ❖ The (new) sample space: $\Omega = \{0, 1, \dots, n\}$.
- ❖ The probability measure is induced from the atomic probabilities: $b(k) = b_n(k; p) := \mathbf{P}\{S_n = k\}$.

$$b(k) = \binom{n}{k} p^k q^{n-k} \quad (k = 0, 1, \dots, n)$$

Positivity ?

The binomial distribution

- ❖ Accumulated successes: New chance variables from old.

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p)$$

$$S_n = X_1 + X_2 + \dots + X_n \sim \text{Binomial}(n, p)$$

- ❖ The (new) sample space: $\Omega = \{0, 1, \dots, n\}$.
- ❖ The probability measure is induced from the atomic probabilities: $b(k) = b_n(k; p) := \mathbf{P}\{S_n = k\}$.

$$b(k) = \binom{n}{k} p^k q^{n-k} \quad (k = 0, 1, \dots, n)$$

Positivity ✓ $b_n(k; p) \geq 0$ for $k = 0, 1, \dots, n$

The binomial distribution

- ❖ Accumulated successes: New chance variables from old.

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p)$$

$$S_n = X_1 + X_2 + \dots + X_n \sim \text{Binomial}(n, p)$$

- ❖ The (new) sample space: $\Omega = \{0, 1, \dots, n\}$.
- ❖ The probability measure is induced from the atomic probabilities: $b(k) = b_n(k; p) := \mathbf{P}\{S_n = k\}$.

$$b(k) = \binom{n}{k} p^k q^{n-k} \quad (k = 0, 1, \dots, n)$$

Positivity ✓ $b_n(k; p) \geq 0$ for $k = 0, 1, \dots, n$

Normalisation ?

The binomial distribution

- ❖ Accumulated successes: New chance variables from old.

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p)$$

$$S_n = X_1 + X_2 + \dots + X_n \sim \text{Binomial}(n, p)$$

- ❖ The (new) sample space: $\Omega = \{0, 1, \dots, n\}$.
- ❖ The probability measure is induced from the atomic probabilities: $b(k) = b_n(k; p) := \mathbf{P}\{S_n = k\}$.

$$b(k) = \binom{n}{k} p^k q^{n-k} \quad (k = 0, 1, \dots, n)$$

Positivity ✓ $b_n(k; p) \geq 0$ for $k = 0, 1, \dots, n$

Normalisation ✓ $\sum_k b_n(k; p) = \sum_{k=0}^n \binom{n}{k} p^k q^{n-k} = (p + q)^n = 1$

The binomial theorem!

$$b(k) = \binom{n}{k} p^k q^{n-k} \quad (k = 0, 1, \dots, n)$$

$$b(k) = \binom{n}{k} p^k q^{n-k} \quad (k = 0, 1, \dots, n)$$

$n = 100$

