

Lesson 3.2-3.3

[Back to Week 1](#)

11/11 points earned
(100%)

Quiz passed!



Be Recognized for Your Achievements. "Course Certificates give you the recognition you need to get the job, the material gives you the skills to do the job. It makes you look more valuable because you are more valuable." - Peter B., USA, Software Developer

Showcase Your Accomplishment! Earn Your Course Certificate! ₹1,953 ➔



1 / 1
points

1.

If continuous random variable X has probability density function (PDF) $f(x)$, what is the interpretation of the following integral:

$$\int_{-2}^5 f(x) dx ?$$

☐ $P(X \geq -2 \cup X \leq 5)$

☐ $P(X \leq -2 \cap X \geq 5)$



☐ $P(X \leq -2 \cap X \leq 5)$

☒ $P(X \geq -2 \cap X \leq 5)$

Correct Response

This could also be written $P(-2 \leq X \leq 5)$.



1 / 1
points

2. If $X \sim \text{Uniform}(0, 1)$, then what is the value of $P(-3 < X < 0.2)$?

0.2

Correct Response

$$\int_{-3}^{0.2} f(x)dx = \int_{-3}^{0.2} I_{\{0 < x < 1\}}(x)dx = \int_0^{0.2} 1dx = 0.2.$$



1 / 1
points

3. If $X \sim \text{Exponential}(5)$, find the expected value $E(X)$.

0.2

Correct Response

With $X \sim \text{Exponential}(\lambda)$, we have $E(X) = 1/\lambda$.



1 / 1
points

4.

Which of the following scenarios could we most appropriately model using an exponentially distributed random variable?

- ☐ The number of failed lightbulbs in a batch of 5000 after 100 hours in service
- ☒ The lifetime in hours of a particular lightbulb

Correct Response

This is a positive, continuous quantity.

- ☐ The hours of service until all light bulbs in a batch of 5000 fail
- ☐ The probability of a light bulb failure before 100 hours in service







1 / 1
points

5.

If $X \sim \text{Uniform}(2, 6)$, which of the following is the PDF of X ?




☐☐

**Correct Response**

This PDF has uniform value (1/4) over the interval $[2, 6]$ and is 0 everywhere else.





1 / 1
points

6. If $X \sim \text{Uniform}(2, 6)$, what is $P(2 < X \leq 3)$? Round your answer to two decimal places.

0.25

Correct Response



This is $\int_2^3 1/4 dx$.




1 / 1
points


7.
If $X \sim N(0, 1)$, which of the following is the PDF of X ?






**Correct Response**

This is the standard normal distribution.





1 / 1
points

8. If $X \sim N(2, 1)$, what is the expected value of $-5X$? This is denoted as $E(-5X)$.

-10

Correct Response

For any number c and any random variable with expectation $E(X)$, we have $E(cX) = cE(X)$.



1 / 1
points

9. Let $X \sim N(1, 1)$ and $Y \sim N(4, 3^2)$. What is the value of $E(X + Y)$?

5

Correct Response

For random variables X and Y with expectations $E(X)$ and $E(Y)$, we always have $E(X + Y) = E(X) + E(Y)$.



1 / 1
points

10.

The normal distribution is also linear in the sense that if $X \sim N(\mu, \sigma^2)$, then for any real constants $a \neq 0$ and b , the distribution of $Y = aX + b$ is distributed $N(a\mu + b, a^2\sigma^2)$.

Using this fact, what is the distribution of $Z = \frac{X - \mu}{\sigma}$?

- ☐ $N(\mu, \sigma^2)$
- ☐ $N(\mu, \sigma)$
- ☐ $N(\mu/\sigma, 1)$
- ☐ $N(1, \sigma^2)$
- ☒ $N(0, 1)$

Correct Response

Here $a = 1/\sigma$ and $b = -\mu/\sigma$. Subtracting the mean and dividing by the standard deviation is referred to as standardizing a random variable.



1 / 1
points

11.

Which of the following random variables would yield the highest value of $P(-1 < X < 1)$?

Hint: Random variables with larger variance are more dispersed.

☒ $X \sim N(0, 0.1)$

Correct Response

Of the four options, this is the least dispersed, meaning that most of the probability is associated with small values of X .

☐ $X \sim N(0, 1)$

☐ $X \sim N(0, 10)$

☐ $X \sim N(0, 100)$

