

Why drug testing works

Drug	Y_1, Y_2, \dots, Y_n
Placebo	Z_1, Z_2, \dots, Z_n

Drug	Y_1, Y_2, \dots, Y_n
Placebo	Z_1, Z_2, \dots, Z_n

$$\Delta := \left| \frac{Y_1 + Y_2 + \dots + Y_n}{n} - \frac{Z_1 + Z_2 + \dots + Z_n}{n} \right|$$

Drug	Y_1, Y_2, \dots, Y_n
Placebo	Z_1, Z_2, \dots, Z_n

$$\Delta := \left| \frac{Y_1 + Y_2 + \dots + Y_n}{n} - \frac{Z_1 + Z_2 + \dots + Z_n}{n} \right|$$

Is the **observed discrepancy** Δ larger than we would expect from random variation if the drug were truly safe?

Drug	Y_1, Y_2, \dots, Y_n
Placebo	Z_1, Z_2, \dots, Z_n

$$\Delta := \left| \frac{Y_1 + Y_2 + \dots + Y_n}{n} - \frac{Z_1 + Z_2 + \dots + Z_n}{n} \right|$$

Is the **observed discrepancy** Δ larger than we would expect from random variation if the drug were truly safe?

The basis of a statistical test:

Assess the likelihood of the **observed discrepancy** *if the drug were indeed truly harmless* by comparison with a *gedanken* experiment consisting of a double sample of placebo patients.

A *gedkanken* or thought experiment: a double placebo sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p)$$

$$X'_1, X'_2, \dots, X'_n$$

A *gedkanken* or thought experiment: a double placebo sample

$$\begin{array}{l} X_1, X_2, \dots, X_n \\ X'_1, X'_2, \dots, X'_n \end{array} \sim \text{Bernoulli}(p) \quad \rightarrow \quad \begin{array}{l} S_n = X_1 + X_2 + \dots + X_n \\ S'_n = X'_1 + X'_2 + \dots + X'_n \end{array} \sim \text{Binomial}(n, p)$$

A *gedkanken* or thought experiment: a double placebo sample

$$\begin{array}{l} X_1, X_2, \dots, X_n \\ X'_1, X'_2, \dots, X'_n \end{array} \sim \text{Bernoulli}(p) \quad \rightarrow \quad \begin{array}{l} S_n = X_1 + X_2 + \dots + X_n \\ S'_n = X'_1 + X'_2 + \dots + X'_n \end{array} \sim \text{Binomial}(n, p)$$

We know n but p is the *unknown* rate at which (cardiopulmonary or other) complications occur in the general population

A *gedanken* or thought experiment: a double placebo sample

$$\begin{array}{l} X_1, X_2, \dots, X_n \\ X'_1, X'_2, \dots, X'_n \end{array} \sim \text{Bernoulli}(p) \quad \rightarrow \quad \begin{array}{l} S_n = X_1 + X_2 + \dots + X_n \\ S'_n = X'_1 + X'_2 + \dots + X'_n \end{array} \sim \text{Binomial}(n, p)$$

We know n but p is the *unknown* rate at which (cardiopulmonary or other) complications occur in the general population

Key idea: how likely is it that the difference in the relative frequencies of “success” in the *gedanken* double placebo sample exceeds the **observed discrepancy** Δ ?

A *gedanken* or thought experiment: a double placebo sample

$$\begin{array}{l} X_1, X_2, \dots, X_n \\ X'_1, X'_2, \dots, X'_n \end{array} \sim \text{Bernoulli}(p) \quad \longrightarrow \quad \begin{array}{l} S_n = X_1 + X_2 + \dots + X_n \\ S'_n = X'_1 + X'_2 + \dots + X'_n \end{array} \sim \text{Binomial}(n, p)$$

We know n but p is the *unknown* rate at which (cardiopulmonary or other) complications occur in the general population

Key idea: how likely is it that the difference in the relative frequencies of “success” in the *gedanken* double placebo sample exceeds the **observed discrepancy** Δ ?

We want to evaluate

$$\mathbf{P} \left\{ \left| \frac{S_n}{n} - \frac{S'_n}{n} \right| > \Delta \right\}$$

A *gedanken* or thought experiment: a double placebo sample

$$\begin{array}{l} X_1, X_2, \dots, X_n \\ X'_1, X'_2, \dots, X'_n \end{array} \sim \text{Bernoulli}(p) \quad \rightarrow \quad \begin{array}{l} S_n = X_1 + X_2 + \dots + X_n \\ S'_n = X'_1 + X'_2 + \dots + X'_n \end{array} \sim \text{Binomial}(n, p)$$

We know n but p is the *unknown* rate at which (cardiopulmonary or other) complications occur in the general population

Key idea: how likely is it that the difference in the relative frequencies of “success” in the *gedanken* double placebo sample exceeds the **observed discrepancy** Δ ?

We want to evaluate

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \frac{S'_n}{n}\right| > \Delta\right\} = \mathbf{P}\{|S_n - S'_n| > n\Delta\}$$

$$S_n = X_1 + X_2 + \cdots + X_n \sim \text{Binomial}(n, p)$$

$$S'_n = X'_1 + X'_2 + \cdots + X'_n$$

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \frac{S'_n}{n}\right| > \Delta\right\}$$

$$S_n = X_1 + X_2 + \cdots + X_n \sim \text{Binomial}(n, p)$$

$$S'_n = X'_1 + X'_2 + \cdots + X'_n$$

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \frac{S'_n}{n}\right| > \Delta\right\} = \mathbf{P}\{|S_n - S'_n| > n\Delta\}$$

$$S_n = X_1 + X_2 + \cdots + X_n \sim \text{Binomial}(n, p)$$

$$S'_n = X'_1 + X'_2 + \cdots + X'_n$$

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \frac{S'_n}{n}\right| > \Delta\right\} = \mathbf{P}\{|S_n - S'_n| > n\Delta\}$$

$$|S_n - S'_n| = |(S_n - np) - (S'_n - np)|$$

$$S_n = X_1 + X_2 + \cdots + X_n \sim \text{Binomial}(n, p)$$

$$S'_n = X'_1 + X'_2 + \cdots + X'_n$$

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \frac{S'_n}{n}\right| > \Delta\right\} = \mathbf{P}\{|S_n - S'_n| > n\Delta\}$$

$$|S_n - S'_n| = |(S_n - np) - (S'_n - np)| \quad \text{Triangle inequality: } |a - b| \leq |a| + |b|$$

$$S_n = X_1 + X_2 + \cdots + X_n \sim \text{Binomial}(n, p)$$

$$S'_n = X'_1 + X'_2 + \cdots + X'_n$$

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \frac{S'_n}{n}\right| > \Delta\right\} = \mathbf{P}\{|S_n - S'_n| > n\Delta\}$$

$$|S_n - S'_n| = |(S_n - np) - (S'_n - np)|$$

Triangle inequality: $|a - b| \leq |a| + |b|$

$$\leq |S_n - np| + |S'_n - np|$$

$$S_n = X_1 + X_2 + \cdots + X_n \sim \text{Binomial}(n, p)$$

$$S'_n = X'_1 + X'_2 + \cdots + X'_n$$

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \frac{S'_n}{n}\right| > \Delta\right\} = \mathbf{P}\{|S_n - S'_n| > n\Delta\}$$

$$|S_n - S'_n| = |(S_n - np) - (S'_n - np)|$$

Triangle inequality: $|a - b| \leq |a| + |b|$

$$\leq |S_n - np| + |S'_n - np|$$

If $|a - b| > c$ then $|a| > c/2$ or $|b| > c/2$

$$S_n = X_1 + X_2 + \cdots + X_n \sim \text{Binomial}(n, p)$$

$$S'_n = X'_1 + X'_2 + \cdots + X'_n$$

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \frac{S'_n}{n}\right| > \Delta\right\} = \mathbf{P}\{|S_n - S'_n| > n\Delta\}$$

$$|S_n - S'_n| = |(S_n - np) - (S'_n - np)|$$

Triangle inequality: $|a - b| \leq |a| + |b|$

$$\leq |S_n - np| + |S'_n - np|$$

If $|a - b| > c$ then $|a| > c/2$ or $|b| > c/2$

If the event $|S_n - S'_n| > n\Delta$ occurs then either $|S_n - np| > n\Delta/2$ or $|S'_n - np| > n\Delta/2$ or both occur

$$S_n = X_1 + X_2 + \cdots + X_n \sim \text{Binomial}(n, p)$$

$$S'_n = X'_1 + X'_2 + \cdots + X'_n$$

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \frac{S'_n}{n}\right| > \Delta\right\} = \mathbf{P}\{|S_n - S'_n| > n\Delta\}$$

$$|S_n - S'_n| = |(S_n - np) - (S'_n - np)|$$

Triangle inequality: $|a - b| \leq |a| + |b|$

$$\leq |S_n - np| + |S'_n - np|$$

If $|a - b| > c$ then $|a| > c/2$ or $|b| > c/2$

If the event $|S_n - S'_n| > n\Delta$ occurs then either $|S_n - np| > n\Delta/2$ or $|S'_n - np| > n\Delta/2$ or both occur

$$\{|S_n - S'_n| > n\Delta\} \subseteq \{|S_n - np| > n\Delta/2\} \cup \{|S'_n - np| > n\Delta/2\}$$