



The coupon collector's problem

The coupon collector's problem

A randomly selected coupon (playbill, collectible) from a collection of n distinct coupons is included with each piece of merchandise purchased.

The coupon collector's problem

A randomly selected coupon (playbill, collectible) from a collection of n distinct coupons is included with each piece of merchandise purchased.

- * How many distinct coupons are obtained if t purchases are made?

The coupon collector's problem

A randomly selected coupon (playbill, collectible) from a collection of n distinct coupons is included with each piece of merchandise purchased.

- ✧ How many distinct coupons are obtained if t purchases are made?
- ✧ How many purchases must be made in order to obtain a complete set of coupons?

- ❖ What are the sample points of the experiment?

❖ What are the sample points of the experiment?

Sampling with replacement: $(k_1, k_2, \dots, k_t) \in \{1, 2, \dots, n\}^t$

- ❖ What are the sample points of the experiment?

Sampling with replacement: $(k_1, k_2, \dots, k_t) \in \{1, 2, \dots, n\}^t$

- ❖ What is the probability measure?

- ❖ What are the sample points of the experiment?

Sampling with replacement: $(k_1, k_2, \dots, k_t) \in \{1, 2, \dots, n\}^t$

- ❖ What is the probability measure?

Combinatorial measure placing equal mass on each atom: $\mathbf{P}\{(k_1, k_2, \dots, k_n)\} = \frac{1}{n^t}$

- ❖ What are the sample points of the experiment?

Sampling with replacement: $(k_1, k_2, \dots, k_t) \in \{1, 2, \dots, n\}^t$

- ❖ What is the probability measure?

Combinatorial measure placing equal mass on each atom: $\mathbf{P}\{(k_1, k_2, \dots, k_n)\} = \frac{1}{n^t}$

- ❖ What are the events of interest?

- ❖ What are the sample points of the experiment?

Sampling with replacement: $(k_1, k_2, \dots, k_t) \in \{1, 2, \dots, n\}^t$

- ❖ What is the probability measure?

Combinatorial measure placing equal mass on each atom: $\mathbf{P}\{(k_1, k_2, \dots, k_n)\} = \frac{1}{n^t}$

- ❖ What are the events of interest?

$A_j :=$ the j th coupon is not obtained in t purchases

- ❖ What are the sample points of the experiment?

Sampling with replacement: $(k_1, k_2, \dots, k_t) \in \{1, 2, \dots, n\}^t$

- ❖ What is the probability measure?

Combinatorial measure placing equal mass on each atom: $\mathbf{P}\{(k_1, k_2, \dots, k_n)\} = \frac{1}{n^t}$

- ❖ What are the events of interest?

$A_j :=$ the j th coupon is not obtained in t purchases

$$X_j = \begin{cases} 1 & \text{if } A_j \text{ occurs} \\ 0 & \text{otherwise} \end{cases}$$

- ❖ What are the sample points of the experiment?

Sampling with replacement: $(k_1, k_2, \dots, k_t) \in \{1, 2, \dots, n\}^t$

- ❖ What is the probability measure?

Combinatorial measure placing equal mass on each atom: $\mathbf{P}\{(k_1, k_2, \dots, k_n)\} = \frac{1}{n^t}$

- ❖ What are the events of interest?

$A_j :=$ the j th coupon is not obtained in t purchases

$$X_j = \begin{cases} 1 & \text{if } A_j \text{ occurs} \\ 0 & \text{otherwise} \end{cases}$$

$A_1^c \cap \dots \cap A_n^c = \{X_1 + \dots + X_n = 0\}$ all the coupons are acquired

- ❖ What are the sample points of the experiment?

Sampling with replacement: $(k_1, k_2, \dots, k_t) \in \{1, 2, \dots, n\}^t$

- ❖ What is the probability measure?

Combinatorial measure placing equal mass on each atom: $\mathbf{P}\{(k_1, k_2, \dots, k_n)\} = \frac{1}{n^t}$

- ❖ What are the events of interest?

$A_j :=$ the j th coupon is not obtained in t purchases

$$X_j = \begin{cases} 1 & \text{if } A_j \text{ occurs} \\ 0 & \text{otherwise} \end{cases}$$

$A_1^c \cap \dots \cap A_n^c = \{X_1 + \dots + X_n = 0\}$ all the coupons are acquired

$\{X_1 + \dots + X_n = k\}$ k of the coupons are still to be acquired

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \quad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

The 1st term $\mathbf{P}(A_1) = \mathbf{P}\{(k_1, k_2, \dots, k_n) : k_i \neq 1 \text{ (each } i)\}$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

The 1st term $\mathbf{P}(A_1) = \mathbf{P}\{(k_1, k_2, \dots, k_n) : k_i \neq 1 \text{ (each } i)\} = \left(1 - \frac{1}{n}\right)^t$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

The 1st term

$$\mathbf{P}(A_1) = \mathbf{P}\{(k_1, k_2, \dots, k_n) : k_i \neq 1 \text{ (each } i)\} = \left(1 - \frac{1}{n}\right)^t$$

$$S_1 = \binom{n}{1} \mathbf{P}(A_1) = n \left(1 - \frac{1}{n}\right)^t$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

The 1st term

$$\mathbf{P}(A_1) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \neq 1 \text{ (each } i) \} = \left(1 - \frac{1}{n}\right)^t$$

$$S_1 = \binom{n}{1} \mathbf{P}(A_1) = n \left(1 - \frac{1}{n}\right)^t$$

The 2nd term

$$\mathbf{P}(A_1 \cap A_2) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \notin \{1, 2\} \text{ (each } i) \}$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

The 1st term

$$\mathbf{P}(A_1) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \neq 1 \text{ (each } i) \} = \left(1 - \frac{1}{n}\right)^t$$

$$S_1 = \binom{n}{1} \mathbf{P}(A_1) = n \left(1 - \frac{1}{n}\right)^t$$

The 2nd term

$$\mathbf{P}(A_1 \cap A_2) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \notin \{1, 2\} \text{ (each } i) \} = \left(1 - \frac{2}{n}\right)^t$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

The 1st term

$$\mathbf{P}(A_1) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \neq 1 \text{ (each } i) \} = \left(1 - \frac{1}{n}\right)^t$$

$$S_1 = \binom{n}{1} \mathbf{P}(A_1) = n \left(1 - \frac{1}{n}\right)^t$$

The 2nd term

$$\mathbf{P}(A_1 \cap A_2) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \notin \{1, 2\} \text{ (each } i) \} = \left(1 - \frac{2}{n}\right)^t$$

$$S_2 = \binom{n}{2} \mathbf{P}(A_1 \cap A_2) = \frac{n^2}{2!} \left(1 - \frac{2}{n}\right)^t$$

The jth term

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

The 1st term

$$\mathbf{P}(A_1) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \neq 1 \text{ (each } i) \} = \left(1 - \frac{1}{n}\right)^t$$

$$S_1 = \binom{n}{1} \mathbf{P}(A_1) = n \left(1 - \frac{1}{n}\right)^t$$

The 2nd term

$$\mathbf{P}(A_1 \cap A_2) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \notin \{1, 2\} \text{ (each } i) \} = \left(1 - \frac{2}{n}\right)^t$$

$$S_2 = \binom{n}{2} \mathbf{P}(A_1 \cap A_2) = \frac{n^2}{2!} \left(1 - \frac{2}{n}\right)^t$$

The jth term

$$\mathbf{P}(A_1 \cap A_2 \cdots \cap A_j) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \notin \{1, 2, \dots, j\} \text{ (each } i) \}$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

The 1st term

$$\mathbf{P}(A_1) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \neq 1 \text{ (each } i) \} = \left(1 - \frac{1}{n}\right)^t$$

$$S_1 = \binom{n}{1} \mathbf{P}(A_1) = n \left(1 - \frac{1}{n}\right)^t$$

The 2nd term

$$\mathbf{P}(A_1 \cap A_2) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \notin \{1, 2\} \text{ (each } i) \} = \left(1 - \frac{2}{n}\right)^t$$

$$S_2 = \binom{n}{2} \mathbf{P}(A_1 \cap A_2) = \frac{n^2}{2!} \left(1 - \frac{2}{n}\right)^t$$

The jth term

$$\mathbf{P}(A_1 \cap A_2 \cdots \cap A_j) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \notin \{1, 2, \dots, j\} \text{ (each } i) \} = \left(1 - \frac{j}{n}\right)^t$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

The 1st term

$$\mathbf{P}(A_1) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \neq 1 \text{ (each } i) \} = \left(1 - \frac{1}{n}\right)^t$$

$$S_1 = \binom{n}{1} \mathbf{P}(A_1) = n \left(1 - \frac{1}{n}\right)^t$$

The 2nd term

$$\mathbf{P}(A_1 \cap A_2) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \notin \{1, 2\} \text{ (each } i) \} = \left(1 - \frac{2}{n}\right)^t$$

$$S_2 = \binom{n}{2} \mathbf{P}(A_1 \cap A_2) = \frac{n^2}{2!} \left(1 - \frac{2}{n}\right)^t$$

The jth term

$$\mathbf{P}(A_1 \cap A_2 \cdots \cap A_j) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \notin \{1, 2, \dots, j\} \text{ (each } i) \} = \left(1 - \frac{j}{n}\right)^t$$

$$S_j = \binom{n}{j} \mathbf{P}(A_1 \cap A_2 \cap \cdots \cap A_j)$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

The 1st term

$$\mathbf{P}(A_1) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \neq 1 \text{ (each } i) \} = \left(1 - \frac{1}{n}\right)^t$$

$$S_1 = \binom{n}{1} \mathbf{P}(A_1) = n \left(1 - \frac{1}{n}\right)^t$$

The 2nd term

$$\mathbf{P}(A_1 \cap A_2) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \notin \{1, 2\} \text{ (each } i) \} = \left(1 - \frac{2}{n}\right)^t$$

$$S_2 = \binom{n}{2} \mathbf{P}(A_1 \cap A_2) = \frac{n^2}{2!} \left(1 - \frac{2}{n}\right)^t$$

The jth term

$$\mathbf{P}(A_1 \cap A_2 \cdots \cap A_j) = \mathbf{P}\{ (k_1, k_2, \dots, k_n) : k_i \notin \{1, 2, \dots, j\} \text{ (each } i) \} = \left(1 - \frac{j}{n}\right)^t$$

$$S_j = \binom{n}{j} \mathbf{P}(A_1 \cap A_2 \cap \cdots \cap A_j) = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$s_j = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$S_j = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t$$

$$\lim_{n \rightarrow \infty} \left(1 - \frac{x}{n}\right)^n = e^{-x}$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$S_j = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^{n \cdot \frac{t}{n}}$$

$$\lim_{n \rightarrow \infty} \left(1 - \frac{x}{n}\right)^n = e^{-x}$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\}$$

$$\{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$S_j = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^{n \cdot \frac{t}{n}} \approx \frac{n^j \cdot e^{-j \cdot t/n}}{j!}$$

$$\lim_{n \rightarrow \infty} \left(1 - \frac{x}{n}\right)^n = e^{-x}$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\}$$

$$\{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$S_j = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^{n \cdot \frac{t}{n}} \approx \frac{n^j \cdot e^{-j \cdot t/n}}{j!} = \frac{(ne^{-t/n})^j}{j!}$$

$$\lim_{n \rightarrow \infty} \left(1 - \frac{x}{n}\right)^n = e^{-x}$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$S_j = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^{n \cdot \frac{t}{n}} \approx \frac{n^j \cdot e^{-j \cdot t/n}}{j!} = \frac{(ne^{-t/n})^j}{j!}$$

$$\lim_{n \rightarrow \infty} \left(1 - \frac{x}{n}\right)^n = e^{-x}$$

If $ne^{-t/n} = \lambda$ or, equivalently, $t = n \log \frac{n}{\lambda}$, then $S_j \rightarrow \frac{\lambda^j}{j!}$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$S_j = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^{n \cdot \frac{t}{n}} \approx \frac{n^j \cdot e^{-j \cdot t/n}}{j!} = \frac{(ne^{-t/n})^j}{j!}$$

$$\lim_{n \rightarrow \infty} \left(1 - \frac{x}{n}\right)^n = e^{-x}$$

If $ne^{-t/n} = \lambda$ or, equivalently, $t = n \log \frac{n}{\lambda}$, then $S_j \rightarrow \frac{\lambda^j}{j!}$

$$\mathbf{P}(A_1^c \cap A_2^c \cap \cdots \cap A_n^c) = S_0 - S_1 + S_2 - \cdots + (-1)^n S_n$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$S_j = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^{n \cdot \frac{t}{n}} \approx \frac{n^j \cdot e^{-j \cdot t/n}}{j!} = \frac{(ne^{-t/n})^j}{j!}$$

$$\lim_{n \rightarrow \infty} \left(1 - \frac{x}{n}\right)^n = e^{-x}$$

If $ne^{-t/n} = \lambda$ or, equivalently, $t = n \log \frac{n}{\lambda}$, then $S_j \rightarrow \frac{\lambda^j}{j!}$

$$\mathbf{P}(A_1^c \cap A_2^c \cap \cdots \cap A_n^c) = S_0 - S_1 + S_2 - \cdots + (-1)^n S_n \rightarrow e^{-\lambda} \quad \text{desired confidence}$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$S_j = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^{n \cdot \frac{t}{n}} \approx \frac{n^j \cdot e^{-j \cdot t/n}}{j!} = \frac{(ne^{-t/n})^j}{j!}$$

$$\lim_{n \rightarrow \infty} \left(1 - \frac{x}{n}\right)^n = e^{-x}$$

If $ne^{-t/n} = \lambda$ or, equivalently, $t = n \log \frac{n}{\lambda}$, then $S_j \rightarrow \frac{\lambda^j}{j!}$

$$\mathbf{P}(A_1^c \cap A_2^c \cap \cdots \cap A_n^c) = S_0 - S_1 + S_2 - \cdots + (-1)^n S_n \rightarrow e^{-\lambda} \quad \text{desired confidence}$$

$$\mathbf{P}\{X_1 + X_2 + \cdots + X_n = k\} = S_k - \binom{k+1}{k} S_{k+1} + \binom{k+2}{k} S_{k+2} - \cdots + (-1)^{n-k} \binom{n}{k} S_n$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$S_j = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^{n \cdot \frac{t}{n}} \approx \frac{n^j \cdot e^{-j \cdot t/n}}{j!} = \frac{(ne^{-t/n})^j}{j!}$$

$$\lim_{n \rightarrow \infty} \left(1 - \frac{x}{n}\right)^n = e^{-x}$$

If $ne^{-t/n} = \lambda$ or, equivalently, $t = n \log \frac{n}{\lambda}$, then $S_j \rightarrow \frac{\lambda^j}{j!}$

$$\mathbf{P}(A_1^c \cap A_2^c \cap \cdots \cap A_n^c) = S_0 - S_1 + S_2 - \cdots + (-1)^n S_n \rightarrow e^{-\lambda} \quad \text{desired confidence}$$

$$\mathbf{P}\{X_1 + X_2 + \cdots + X_n = k\} = S_k - \binom{k+1}{k} S_{k+1} + \binom{k+2}{k} S_{k+2} - \cdots + (-1)^{n-k} \binom{n}{k} S_n \rightarrow e^{-\lambda} \frac{\lambda^k}{k!}$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$S_j = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^{n \cdot \frac{t}{n}} \approx \frac{n^j \cdot e^{-j \cdot t/n}}{j!} = \frac{(ne^{-t/n})^j}{j!}$$

$$\lim_{n \rightarrow \infty} \left(1 - \frac{x}{n}\right)^n = e^{-x}$$

If $ne^{-t/n} = \lambda$ or, equivalently, $t = n \log \frac{n}{\lambda}$, then $S_j \rightarrow \frac{\lambda^j}{j!}$

$$\mathbf{P}(A_1^c \cap A_2^c \cap \cdots \cap A_n^c) = S_0 - S_1 + S_2 - \cdots + (-1)^n S_n \rightarrow e^{-\lambda} \quad \text{desired confidence}$$

$$\mathbf{P}\{X_1 + X_2 + \cdots + X_n = k\} = S_k - \binom{k+1}{k} S_{k+1} + \binom{k+2}{k} S_{k+2} - \cdots + (-1)^{n-k} \binom{n}{k} S_n \rightarrow e^{-\lambda} \frac{\lambda^k}{k!}$$

The number of purchases needed to acquire all 14 Gilbert and Sullivan playbills:

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$S_j = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^{n \cdot \frac{t}{n}} \approx \frac{n^j \cdot e^{-j \cdot t/n}}{j!} = \frac{(ne^{-t/n})^j}{j!}$$

$$\lim_{n \rightarrow \infty} \left(1 - \frac{x}{n}\right)^n = e^{-x}$$

If $ne^{-t/n} = \lambda$ or, equivalently, $t = n \log \frac{n}{\lambda}$, then $S_j \rightarrow \frac{\lambda^j}{j!}$

$$\mathbf{P}(A_1^c \cap A_2^c \cap \cdots \cap A_n^c) = S_0 - S_1 + S_2 - \cdots + (-1)^n S_n \rightarrow e^{-\lambda} \quad \text{desired confidence}$$

$$\mathbf{P}\{X_1 + X_2 + \cdots + X_n = k\} = S_k - \binom{k+1}{k} S_{k+1} + \binom{k+2}{k} S_{k+2} - \cdots + (-1)^{n-k} \binom{n}{k} S_n \rightarrow e^{-\lambda} \frac{\lambda^k}{k!}$$

The number of purchases needed to acquire all 14 Gilbert and Sullivan playbills:

$$\text{For 90\% confidence } \lambda = -\log(0.9) = 0.10536 \text{ and } t = \lceil 14 \log \frac{14}{0.10536} \rceil = 69.$$

$$A_1^c \cap \cdots \cap A_n^c = \{X_1 + \cdots + X_n = 0\} \qquad \{X_1 + \cdots + X_n = k\}$$

The inclusion–exclusion sums

$$S_j = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^t = \frac{n^j}{j!} \left(1 - \frac{j}{n}\right)^{n \cdot \frac{t}{n}} \approx \frac{n^j \cdot e^{-j \cdot t/n}}{j!} = \frac{(ne^{-t/n})^j}{j!}$$

$$\lim_{n \rightarrow \infty} \left(1 - \frac{x}{n}\right)^n = e^{-x}$$

If $ne^{-t/n} = \lambda$ or, equivalently, $t = n \log \frac{n}{\lambda}$, then $S_j \rightarrow \frac{\lambda^j}{j!}$

$$\mathbf{P}(A_1^c \cap A_2^c \cap \cdots \cap A_n^c) = S_0 - S_1 + S_2 - \cdots + (-1)^n S_n \rightarrow e^{-\lambda} \quad \text{desired confidence}$$

$$\mathbf{P}\{X_1 + X_2 + \cdots + X_n = k\} = S_k - \binom{k+1}{k} S_{k+1} + \binom{k+2}{k} S_{k+2} - \cdots + (-1)^{n-k} \binom{n}{k} S_n \rightarrow e^{-\lambda} \frac{\lambda^k}{k!}$$

The number of purchases needed to acquire all 14 Gilbert and Sullivan playbills:

For 90% confidence $\lambda = -\log(0.9) = 0.10536$ and $t = \lceil 14 \log \frac{14}{0.10536} \rceil = 69$.

For 99% confidence $\lambda = -\log(0.99) = 0.01005$ and $t = \lceil 14 \log \frac{14}{0.01005} \rceil = 102$.