Tableau 6, Part 2

Probabilities in simple settings
Unexpected ramifications from the repeated toss of a coin, continuous spaces

Tableau 6, Part 2

Probabilities in simple settings
Unexpected ramifications from the repeated toss of a coin, continuous spaces



The probabilist's trinity (Ω , \mathcal{F} , P)

- * What is the collection of conceptual outcomes of the chance experiment?
- * What are the events of interest?
- * What are the chances associated with the events?

Sample space: Ω

Algebra of events: F

Probability measure: P



Toss a coin repeatedly without end

Sample space Ω : each sample point ω is an infinite sequence of heads and tails.

Toss a coin repeatedly without end

Is there a simple characterisation of the events?

Sample space Ω: each sample point ω is an infinite sequence of heads and tails.

Represent heads by 1 and tails by 0.

Toss a coin repeatedly without end

Is there a simple characterisation of the events?

Sample space Ω: each sample point ω is an infinite sequence of heads and tails.

Represent heads by 1 and tails by 0.

 $\Omega = \{ x_1 x_2 x_3 \dots : x_n \in \{0, 1\} \text{ for each } n = 1, 2, 3, \dots \}$

Toss a coin repeatedly without end

Is there a simple characterisation of the events?

- Sample space Ω: each sample point ω is an infinite sequence of heads and tails.
 - Represent heads by 1 and tails by 0.
 - $\Omega = \{ x_1 x_2 x_3 \dots : x_n \in \{0, 1\} \text{ for each } n = 1, 2, 3, \dots \}$

Toss a coin repeatedly without end

- Is there a simple characterisation of the events?
 - Event A that the first head occurs on the fourth toss.

- Sample space Ω : each sample point ω is an infinite sequence of heads and tails.
 - Represent heads by 1 and tails by 0.
 - $\Omega = \{ x_1 x_2 x_3 \dots : x_n \in \{0, 1\} \text{ for each } n = 1, 2, 3, \dots \}$

Toss a coin repeatedly without end

- Is there a simple characterisation of the events?
 - Event A that the first head occurs on the fourth toss.
 - $A = \{x_1x_2x_3 \dots : x_1 = x_2 = x_3 = 0, x_4 = 1, x_n \in \{0, 1\} \text{ for each } n = 5, 6, 7, \dots \}$
- What is an appropriate choice of probability measure P?