



Chevalier de Méré's paradox

Antoine Gombaud

Which of the following is it more advantageous to wager on?

A) At least one ace (ace = "6") in 4 throws of a die.

B) At least one double ace in 24 throws of a pair of dice?

Chevalier de Méré's paradox

Antoine Gombaud

Which of the following is it more advantageous to wager on?

A) At least one ace (ace = "6") in 4 throws of a die.

B) At least one double ace in 24 throws of a pair of dice?

✦ A specious (and incorrect) argument:

A) There is a one-in-six chance of an ace in a die throw. So in four throws the odds of an ace should be 4 : 6.

B) There is a one-in-36 chance of a double ace when a pair of dice are thrown. So in twenty four throws of the pair, the odds of a double ace are in the proportionate ratio 24 : 36.

Chevalier de Méré's paradox

Antoine Gombaud

Which of the following is it more advantageous to wager on?

A) At least one ace (ace = "6") in 4 throws of a die.

B) At least one double ace in 24 throws of a pair of dice?

❖ A specious (and incorrect) argument:

A) There is a one-in-six chance of an ace in a die throw. So in four throws the odds of an ace should be 4 : 6.

B) There is a one-in-36 chance of a double ace when a pair of dice are thrown. So in twenty four throws of the pair, the odds of a double ace are in the proportionate ratio 24 : 36.

❖ What is wrong with this line of reasoning?

Chevalier de Méré's paradox

Antoine Gombaud

Which of the following is it more advantageous to wager on?

A) At least one ace (ace = "6") in 4 throws of a die.

B) At least one double ace in 24 throws of a pair of dice?

❖ A specious (and incorrect) argument:

A) There is a one-in-six chance of an ace in a die throw. So in four throws the odds of an ace should be 4 : 6.


B) There is a one-in-36 chance of a double ace when a pair of dice are thrown. So in twenty four throws of the pair, the odds of a double ace are in the proportionate ratio 24 : 36.

❖ What is wrong with this line of reasoning?

The argument implicitly adopts the working principle that *the chances increase proportionately with the number of trials*. Thus, if the reasoning in A) is taken at face value then the odds of (at least) one ace in six throws of a die should be $6 : 6 = 1$, which is absurd. So something is wrong with the principle.


A balls and urns metaphor: one die

A) Four throws of a die: each throw represents a distinct ball, the face values that are possible represent six distinct urns.



A balls and urns metaphor: one die


A) Four throws of a die: each throw represents a distinct ball, the face values that are possible represent six distinct urns.



A balls and urns metaphor: one die

A) Four throws of a die: each throw represents a distinct ball, the face values that are possible represent six distinct urns.


(5, 1, 5, 2)



A balls and urns metaphor: one die

A) Four throws of a die: each throw represents a distinct ball, the face values that are possible represent six distinct urns.


(5, 1, 5, 2)



A balls and urns metaphor: one die

A) Four throws of a die: each throw represents a distinct ball, the face values that are possible represent six distinct urns.

(5, 1, 5, 2)



- Possible outcomes: (k_1, k_2, k_3, k_4) . Ordered sample (of urns) with replacement. The number of such sequences is 6^4 .
- The **complement** of the event that an ace is thrown occurs if **none** of k_1, k_2, k_3, k_4 is equal to six. There are 5^4 sequences favourable to this event. Under random selection this event hence has probability $5^4/6^4 \approx 0.482$.
- The event that there is an ace hence has probability $P(A) = 1 - \frac{5^4}{6^4} \approx 0.518$.

A balls and urns metaphor: two dice

$$P(A) = 1 - \frac{5^4}{6^4} \approx 0.518$$

- B) Twenty four throws of a pair of dice: each throw represents a distinct ball, the face values that are possible represent six distinct urns.

A balls and urns metaphor: two dice

$$P(A) = 1 - \frac{5^4}{6^4} \approx 0.518$$

B) Twenty four throws of a pair of dice: each throw represents a distinct ball, the face values that are possible represent six distinct urns.

6	6	21		22	15		16
5	2	11		1			10
				24			19
4				5			8
	18		7				
3					14		
						3	17
2	4						13
			20	9			
1				12	23		
	1	2	3	4	5	6	

A balls and urns metaphor: two dice

$$\mathbf{P(A)} = 1 - \frac{5^4}{6^4} \approx 0.518$$

B) Twenty four throws of a pair of dice: each throw represents a distinct ball, the face values that are possible represent six distinct urns.

6	6	21	22	15	16
5	2	11	1		10
4			24		19
3	18	7	5	14	8
2				3	17
1	4	20	9	12	13
	1	2	3	4	5

A balls and urns metaphor: two dice

$$P(A) = 1 - \frac{5^4}{6^4} \approx 0.518$$

B) Twenty four throws of a pair of dice: each throw represents a distinct ball, the face values that are possible represent six distinct urns.

- Possible outcomes: $(j_1, k_1), \dots, (j_{24}, k_{24})$. Ordered sample (of pairs of urns) with replacement. The number of such sequences is 36^{24} .
- The **complement** of the event that an ace pair is obtained occurs if **none** of the pairs $(j_1, k_1), \dots, (j_{24}, k_{24})$ is equal to $(6, 6)$. There are 35^{24} sequences favourable to this event. Under random selection this event hence has probability $35^{24} / 36^{24} \approx 0.509$.
- The event that there is an ace hence has probability $P(B) = 1 - \frac{35^{24}}{36^{24}} \approx 0.491$.

6	6 21		22	15		16
5	2 11		1 24			10 19 8
4			5 7			
3				14		
2	4				3	17 13
1			20 9	12	23	
	1	2	3	4	5	6

A balls and urns metaphor: two dice

$$P(A) = 1 - \frac{5^4}{6^4} \approx 0.518$$

B) Twenty four throws of a pair of dice: each throw represents a distinct ball, the face values that are possible represent six distinct urns.

- Possible outcomes: $(j_1, k_1), \dots, (j_{24}, k_{24})$. Ordered sample (of pairs of urns) with replacement. The number of such sequences is 36^{24} .
- The **complement** of the event that an ace pair is obtained occurs if **none** of the pairs $(j_1, k_1), \dots, (j_{24}, k_{24})$ is equal to $(6, 6)$. There are 35^{24} sequences favourable to this event. Under random selection this event hence has probability $35^{24} / 36^{24} \approx 0.509$.
- The event that there is an ace hence has probability $P(B) = 1 - \frac{35^{24}}{36^{24}} \approx 0.491$.

6	6 21		22	15		16
5	2 11		1 24			10 19 8
4			5 7			
3				14		
2	4				3	17 13
1			20 9	12	23	
	1	2	3	4	5	6

It is (slightly) more likely to obtain an ace in 4 throws of a die than it is to obtain a double ace in 24 throws of a pair of dice.