

A familial paradox

A family with two children is known to have (at least) one boy. What are the chances the other child is a boy?

A familial paradox

A family with two children is known to have (at least) one boy. What are the chances the other child is a boy?

1/2?

● *Sample space:* it is natural to list the gender of the child by age
 $\Omega = \{\mathfrak{B}\mathfrak{B}, \mathfrak{B}\mathfrak{G}, \mathfrak{G}\mathfrak{B}, \mathfrak{G}\mathfrak{G}\}$

● *Sample space*: it is natural to list the gender of the child by age
 $\Omega = \{\mathfrak{B}\mathfrak{B}, \mathfrak{B}\mathfrak{G}, \mathfrak{G}\mathfrak{B}, \mathfrak{G}\mathfrak{G}\}$

● The *events of interest*:

● *Sample space*: it is natural to list the gender of the child by age
 $\Omega = \{\mathfrak{B}\mathfrak{B}, \mathfrak{B}\mathfrak{G}, \mathfrak{G}\mathfrak{B}, \mathfrak{G}\mathfrak{G}\}$

● The *events of interest*:
 $A := \text{both children are boys} = \{\mathfrak{B}\mathfrak{B}\}$

● *Sample space:* it is natural to list the gender of the child by age
 $\Omega = \{\mathfrak{B}\mathfrak{B}, \mathfrak{B}\mathfrak{G}, \mathfrak{G}\mathfrak{B}, \mathfrak{G}\mathfrak{G}\}$

● *The events of interest:*

$A :=$ both children are boys $= \{\mathfrak{B}\mathfrak{B}\}$

$B :=$ (at least) one child is a boy $= \{\mathfrak{B}\mathfrak{B}, \mathfrak{B}\mathfrak{G}, \mathfrak{G}\mathfrak{B}\}$

● *Sample space*: it is natural to list the gender of the child by age
 $\Omega = \{\mathfrak{B}\mathfrak{B}, \mathfrak{B}\mathfrak{G}, \mathfrak{G}\mathfrak{B}, \mathfrak{G}\mathfrak{G}\}$

● *The events of interest*:

$A :=$ both children are boys $= \{\mathfrak{B}\mathfrak{B}\}$

$B :=$ (at least) one child is a boy $= \{\mathfrak{B}\mathfrak{B}, \mathfrak{B}\mathfrak{G}, \mathfrak{G}\mathfrak{B}\}$

● *Probability measure \mathbf{P}* :

- Combinatorial setting: to each *atom* (singleton set) assign equal *probability mass*

$$\mathbf{P}\{\mathfrak{B}\mathfrak{B}\} = \mathbf{P}\{\mathfrak{B}\mathfrak{G}\} = \mathbf{P}\{\mathfrak{G}\mathfrak{B}\} = \mathbf{P}\{\mathfrak{G}\mathfrak{G}\} = 1/4$$

- Event probabilities via additivity:

$$\mathbf{P}(A) = \mathbf{P}\{\mathfrak{B}\mathfrak{B}\} = 1/4$$

$$\mathbf{P}(B) = \mathbf{P}\{\mathfrak{B}\mathfrak{B}\} + \mathbf{P}\{\mathfrak{B}\mathfrak{G}\} + \mathbf{P}\{\mathfrak{G}\mathfrak{B}\} = 3/4$$

BB	BG
GB	GG

Ω

Sample space: it is natural to list the gender of the child by age
 $\Omega = \{BB, BG, GB, GG\}$

The *events of interest*:

$A := \text{both children are boys} = \{BB\}$

$B := \text{(at least) one child is a boy} = \{BB, BG, GB\}$

Probability measure P :

- Combinatorial setting: to each *atom* (singleton set) assign equal *probability mass*

$$P\{BB\} = P\{BG\} = P\{GB\} = P\{GG\} = 1/4$$

- Event probabilities via additivity:

$$P(A) = P\{BB\} = 1/4$$

$$P(B) = P\{BB\} + P\{BG\} + P\{GB\} = 3/4$$

The chance of **A**
(without side information)

$$P(A) = 1/4$$

BB	BG
GB	GG

Ω

Sample space: it is natural to list the gender of the child by age
 $\Omega = \{BB, BG, GB, GG\}$

The *events of interest*:

$A :=$ both children are boys $= \{BB\}$

$B :=$ (at least) one child is a boy $= \{BB, BG, GB\}$

Probability measure **P**:

- Combinatorial setting: to each *atom* (singleton set) assign equal *probability mass*

$$P\{BB\} = P\{BG\} = P\{GB\} = P\{GG\} = 1/4$$

- Event probabilities via additivity:

$$P(A) = P\{BB\} = 1/4$$

$$P(B) = P\{BB\} + P\{BG\} + P\{GB\} = 3/4$$

Sample space: it is natural to list the gender of the child by age
 $\Omega = \{\mathfrak{B}\mathfrak{B}, \mathfrak{B}\mathfrak{G}, \mathfrak{G}\mathfrak{B}, \mathfrak{G}\mathfrak{G}\}$

The *events of interest*:
 $A := \text{both children are boys} = \{\mathfrak{B}\mathfrak{B}\}$
 $B := \text{(at least) one child is a boy} = \{\mathfrak{B}\mathfrak{B}, \mathfrak{B}\mathfrak{G}, \mathfrak{G}\mathfrak{B}\}$

Probability measure \mathbf{P} :

- Combinatorial setting: to each *atom* (singleton set) assign equal *probability mass*

$$\mathbf{P}\{\mathfrak{B}\mathfrak{B}\} = \mathbf{P}\{\mathfrak{B}\mathfrak{G}\} = \mathbf{P}\{\mathfrak{G}\mathfrak{B}\} = \mathbf{P}\{\mathfrak{G}\mathfrak{G}\} = 1/4$$

- Event probabilities via additivity:

$$\mathbf{P}(A) = \mathbf{P}\{\mathfrak{B}\mathfrak{B}\} = 1/4$$

$$\mathbf{P}(B) = \mathbf{P}\{\mathfrak{B}\mathfrak{B}\} + \mathbf{P}\{\mathfrak{B}\mathfrak{G}\} + \mathbf{P}\{\mathfrak{G}\mathfrak{B}\} = 3/4$$

The chance of A
 (without side information)

$$\mathbf{P}(A) = 1/4$$

Conditioned on (at least)
 one boy



Sample space: it is natural to list the gender of the child by age
 $\Omega = \{\mathfrak{B}\mathfrak{B}, \mathfrak{B}\mathfrak{G}, \mathfrak{G}\mathfrak{B}, \mathfrak{G}\mathfrak{G}\}$

The *events of interest*:
 $A := \text{both children are boys} = \{\mathfrak{B}\mathfrak{B}\}$
 $B := \text{(at least) one child is a boy} = \{\mathfrak{B}\mathfrak{B}, \mathfrak{B}\mathfrak{G}, \mathfrak{G}\mathfrak{B}\}$

Probability measure \mathbf{P} :

- Combinatorial setting: to each *atom* (singleton set) assign equal *probability mass*

$$\mathbf{P}\{\mathfrak{B}\mathfrak{B}\} = \mathbf{P}\{\mathfrak{B}\mathfrak{G}\} = \mathbf{P}\{\mathfrak{G}\mathfrak{B}\} = \mathbf{P}\{\mathfrak{G}\mathfrak{G}\} = 1/4$$

- Event probabilities via additivity:

$$\mathbf{P}(A) = \mathbf{P}\{\mathfrak{B}\mathfrak{B}\} = 1/4$$

$$\mathbf{P}(B) = \mathbf{P}\{\mathfrak{B}\mathfrak{B}\} + \mathbf{P}\{\mathfrak{B}\mathfrak{G}\} + \mathbf{P}\{\mathfrak{G}\mathfrak{B}\} = 3/4$$

The chance of A
 (without side information)
 $\mathbf{P}(A) = 1/4$

Conditioned on (at least)
 one boy



The chance of A
 (given that B occurs)
 $\frac{\mathbf{P}(A \cap B)}{\mathbf{P}(B)} = \frac{1/4}{3/4} = \frac{1}{3}$

