


The law of large numbers

What does Chebyshev say about a sum of independent variables?

What does Chebyshev say about a sum of independent variables?

Chebyshev's inequality

$$\mathbf{P}\{|X - \mathbf{E}(X)| > \tau\} \leq \frac{\text{Var}(X)}{\tau^2}$$




What does Chebyshev say about a sum of independent variables?

Chebyshev's inequality

$$\mathbf{P}\{|X - \mathbf{E}(X)| > \tau\} \leq \frac{\text{Var}(X)}{\tau^2}$$


$X \longrightarrow S_n$



What does Chebyshev say about a sum of independent variables?

Chebyshev's inequality

$$\mathbf{P}\{|X - \mathbf{E}(X)| > \tau\} \leq \frac{\text{Var}(X)}{\tau^2} \quad X \xrightarrow{\quad} S_n \quad \mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$



What does Chebyshev say about a sum of independent variables?

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

Repeated independent trials

$$X, X_1, \dots, X_n, \dots, \mathbf{E}(X) = \mu, \text{Var}(X) = \sigma^2$$

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

Repeated independent trials

$$X, X_1, \dots, X_n, \dots, \mathbf{E}(X) = \mu, \text{Var}(X) = \sigma^2$$

$$S_n = X_1 + \dots + X_n$$

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

Repeated independent trials

$$X, X_1, \dots, X_n, \dots, \mathbf{E}(X) = \mu, \text{Var}(X) = \sigma^2$$

$$S_n = X_1 + \dots + X_n$$

additivity

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

Repeated independent trials

$$X, X_1, \dots, X_n, \dots, \mathbf{E}(X) = \mu, \text{Var}(X) = \sigma^2$$

$$S_n = X_1 + \dots + X_n$$

additivity

$$\mathbf{E}(S_n) = \mathbf{E}(X_1) + \dots + \mathbf{E}(X_n) = n\mu$$

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

Repeated independent trials

$$X, X_1, \dots, X_n, \dots, \mathbf{E}(X) = \mu, \text{Var}(X) = \sigma^2$$

$$S_n = X_1 + \dots + X_n$$

additivity

$$\mathbf{E}(S_n) = \mathbf{E}(X_1) + \dots + \mathbf{E}(X_n) = n\mu$$

$$\text{Var}(S_n) = \text{Var}(X_1) + \dots + \text{Var}(X_n) = n\sigma^2$$

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

Repeated independent trials

$$X, X_1, \dots, X_n, \dots, \mathbf{E}(X) = \mu, \text{Var}(X) = \sigma^2$$

$$S_n = X_1 + \dots + X_n$$

additivity

$$\mathbf{E}(S_n) = \mathbf{E}(X_1) + \dots + \mathbf{E}(X_n) = n\mu$$

$$\text{Var}(S_n) = \text{Var}(X_1) + \dots + \text{Var}(X_n) = n\sigma^2$$

$$\sqrt{\text{Var}(S_n)} = \sqrt{n} \sigma$$

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

Repeated independent trials

$$X, X_1, \dots, X_n, \dots, \mathbf{E}(X) = \mu, \text{Var}(X) = \sigma^2$$

$$S_n = X_1 + \dots + X_n$$

additivity

$$\mathbf{E}(S_n) = \mathbf{E}(X_1) + \dots + \mathbf{E}(X_n) = n\mu$$

$$\text{Var}(S_n) = \text{Var}(X_1) + \dots + \text{Var}(X_n) = n\sigma^2$$

$$\sqrt{\text{Var}(S_n)} = \sqrt{n} \sigma$$

The importance of viewing things in the proper scale:

The expectation increases linearly with n ,

but the standard deviation increases only as the square-root of n .

The deviation from the mean

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

$$\mathbf{E}(S_n) = n\mu \qquad \text{Var}(S_n) = n\sigma^2$$

The deviation from the mean

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

$$\mathbf{E}(S_n) = n\mu \qquad \text{Var}(S_n) = n\sigma^2$$

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon\right\}$$

The deviation from the mean

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

$$\mathbf{E}(S_n) = n\mu \qquad \text{Var}(S_n) = n\sigma^2$$

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon\right\} = \mathbf{P}\{|S_n - n\mu| > n\epsilon\}$$

The deviation from the mean

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

$$\mathbf{E}(S_n) = n\mu \qquad \text{Var}(S_n) = n\sigma^2$$

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon\right\} = \mathbf{P}\left\{|S_n - \overbrace{n\mu}^{\mathbf{E}(S_n)}| > \overbrace{n\epsilon}^{\tau}\right\}$$

The deviation from the mean

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

$$\mathbf{E}(S_n) = n\mu \qquad \text{Var}(S_n) = n\sigma^2$$

$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon\right\} &= \mathbf{P}\left\{|S_n - \overbrace{n\mu}^{\mathbf{E}(S_n)}| > \overbrace{n\epsilon}^{\tau}\right\} \\ &\leq \frac{\text{Var}(S_n)}{(n\epsilon)^2} \end{aligned}$$

The deviation from the mean

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

$$\mathbf{E}(S_n) = n\mu \qquad \text{Var}(S_n) = n\sigma^2$$

$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon\right\} &= \mathbf{P}\left\{|S_n - \overbrace{n\mu}^{\mathbf{E}(S_n)}| > \overbrace{n\epsilon}^{\tau}\right\} \\ &\leq \frac{\text{Var}(S_n)}{(n\epsilon)^2} \\ &= \frac{n\sigma^2}{n^2\epsilon^2} \end{aligned}$$

The deviation from the mean

$$\mathbf{P}\{|S_n - \mathbf{E}(S_n)| > \tau\} \leq \frac{\text{Var}(S_n)}{\tau^2}$$

$$\mathbf{E}(S_n) = n\mu \qquad \text{Var}(S_n) = n\sigma^2$$

$$\begin{aligned} \mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon\right\} &= \mathbf{P}\left\{|S_n - \overbrace{n\mu}^{\mathbf{E}(S_n)}| > \overbrace{n\epsilon}^{\tau}\right\} \\ &\leq \frac{\text{Var}(S_n)}{(n\epsilon)^2} \\ &= \frac{n\sigma^2}{n^2\epsilon^2} \\ &= \frac{\sigma^2}{n\epsilon^2} \end{aligned}$$

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon\right\} \leq \frac{\sigma^2}{n\epsilon^2}$$

The law of large numbers

Suppose $X, X_1, \dots, X_n, \dots$ represent repeated independent trials with $E(X) = \mu$ and $\text{Var}(X) = \sigma^2$, and suppose $\epsilon > 0$ is any fixed small, positive number. Then the partial sums $S_n = X_1 + \dots + X_n$ are increasingly concentrated at μ and satisfy the asymptotic relation

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon\right\} \leq \frac{\sigma^2}{n\epsilon^2} \rightarrow 0 \quad (n \rightarrow \infty).$$

The law of large numbers

Suppose $X, X_1, \dots, X_n, \dots$ represent repeated independent trials with $E(X) = \mu$ and $\text{Var}(X) = \sigma^2$, and suppose $\epsilon > 0$ is any fixed small, positive number. Then the partial sums $S_n = X_1 + \dots + X_n$ are increasingly concentrated at μ and satisfy the asymptotic relation

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon\right\} \leq \frac{\sigma^2}{n\epsilon^2} \rightarrow 0 \quad (n \rightarrow \infty).$$

Slogan

The probability that S_n/n deviates from its expected value μ *by even a small amount* is small provided the sample size n is sufficiently large.