Erasmus School of Economics

# **MOOC** Econometrics

Lecture 4.1 on Endogeneity: Motivation

Dennis Fok

**Erasmus University Rotterdam** 



## Interpretation of parameters

Given the estimates (y: flights, x: insurances)

$$y = 10,000 + .25x + e$$

**Correct:** 4,000 insurances sold  $\rightarrow$  expected number of flights =  $10,000 + .25 \times 4,000 = 11,000$ 

- High x tends to go together with high y.
- The identified correlation yields adequate predictions.

**Incorrect:** Selling 4,000 additional insurances causes  $.25 \times 4,000 = 1,000$  additional flights

- The regression does not identify a causal impact!
- A third variable (*travel demand*) affects y (*flights*) and x (*insurances*).

# Motivating example

We want to explain

- Number of flights at an airport per month (y) using
- Number of travel insurances made in previous month (x)

Suppose OLS yields

$$y = 10,000 + .25x + e$$

#### Test

How should we interpret the obtained coefficients? What does the estimate .25 really mean?

Ezafus

Lecture 4.1, Slide 2 of 12, Erasmus School of Economics

# **Endogeneity**

OLS requires some assumptions:

- explanatory variables should be exogenous
- violation of this: endogeneity.

In this set of lectures, you will learn to:

- Understand/recognize endogeneity.
- 2 Know the consequences of endogeneity.
- Stimate parameters under endogeneity.
- Mow the intuition of the new estimator.
- **6** Argue/test assumptions underlying this new estimator.

# Stochastic vs. non-stochastic regressors

Standard assumptions for linear model ( $y = X\beta + \varepsilon$ ) include A2 Explanatory variables are *non-stochastic* 

#### Implications:

- Obtain new data: X stays constant (and y changes)
- Need "controlled experiment"
- OLS estimator b converges to true coefficient  $\beta$  for  $n \to \infty$  (OLS is consistent)

Erofus

Lecture 4.1, Slide 5 of 12, Erasmus School of Economics

## Other examples of endogeneity - Omitted variables

• True model is

$$y = X_1 \beta_1 + X_2 \beta_2 + \eta$$

but we ignore  $X_2$  and perform OLS on

$$y = X_1\beta_1 + \varepsilon$$

- We have:  $\varepsilon = X_2\beta_2 + \eta$
- $X_1$  correlated with  $\varepsilon$  ( $X_1$  is endogenous) if
  - $\triangleright$   $X_1$  correlated with  $X_2$  and
  - $\beta_2 \neq 0$

Derivation:

$$\mathsf{Cov}(X_1, \varepsilon) = \mathsf{Cov}(X_1, X_2 \beta_2 + \eta)$$

$$= \mathsf{Cov}(X_1, X_2) \beta_2 + \underbrace{\mathsf{Cov}(X_1, \eta)}_{=0}$$

Economic models

#### In economics:

- Controlled (or natural) experiments are rare
- New data with same X cannot be obtained
- Explanatory variables are stochastic!

#### If X stochastic:

- new data set  $\rightarrow$  new X values
- X can be correlated with other variables
- If X correlated with  $\varepsilon$ 
  - ▶ *X* is endogenous
  - ▶ There is another variable that affects *y* and *X*
  - ▶ OLS does not properly estimate  $\beta$  (inconsistent)
- ullet If X uncorrelated with arepsilon
  - ▶ *X* is exogenous
  - ▶ OLS consistent

Ezafus,

Lecture 4.1, Slide 6 of 12, Erasmus School of Economics

# Omitted variable – Example

Model student's grade using attendance at lectures.

#### Test

Which omitted factor would lead to endogeneity of attendance?

Three possible omitted factors:

- Difficulty of exam
  - NO: not correlated with attendance.
- Motivation of the students?
  - YES: correlates with attendance and affects grade.
- Compulsory attendance yes/no?

 $\underline{\mathsf{NO}} \mathrm{:}$  does not directly impact the grade

Erafus

## Other examples - Strategic behavior

Other examples – Measurement errors

Consider a model explaining demand using price.

Strategic price setting:

- Sets high price when high demand is expected
- Price and sales positively correlated
- 3 Price will be endogenous in regression of demand on price.

Ezafus,

Lecture 4.1, Slide 9 of 12, Erasmus School of Economics

• y (eg. salary) depends on  $x^*$  (eg. intelligence)

- $x^*$  (intelligence) difficult to observe
- $x = x^* + \text{measurement error}$ : noisy measurement (eg. IQ score)
- measurement error: x is endogenous in  $y = \alpha + \beta x + \varepsilon$

(zafus

Lecture 4.1, Slide 10 of 12, Erasmus School of Economic

# Summary & what's next?

- Endogeneity is a common problem
- OLS is not useful under endogeneity

#### Upcoming topics:

- How to solve for endogeneity?
- How to test for endogeneity?

## TRAINING EXERCISE 4.1

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).

Ezafus,

Erasmus School of Economics

# **MOOC** Econometrics


Lecture 4.2 on Endogeneity: Consequences

Dennis Fok

**Erasmus University Rotterdam** 

# Ezafus

# Simulated example, $y = 1 + 2x^* + u$



# **Endogeneity**

- Common problem in economics
  - Omitted variables
  - Strategic behavior
  - Measurement errors
  - $\rightarrow X$  is correlated with  $\varepsilon$
- Endogeneity violates the basic assumptions
- $\rightarrow$  How bad is this?

Erafus

Lecture 4.2, Slide 2 of 12, Erasmus School of Economics

# Measurement error example

Under measurement error (and endogeneity in general):

• we obtain the wrong coefficients!

#### Test

Can we say anything about the direction of the bias?

Erafus,

## Direction of bias in the measurement error case

OLS is "biased towards zero"

 $\rightarrow \, \mathsf{OLS} \,\, \mathsf{underestimates} \,\, \mathsf{true} \,\, \mathsf{effect}$ 

#### Intuitively:

- x-values on the *left* likely have negative measurement errors
- x-values on the *right* likely have positive measurement errors

Measurement errors "stretch" the scatter in the horizontal direction  $\rightarrow$  a flatter regression line

Erofus

Lecture 4.2, Slide 5 of 12, Erasmus School of Economics

## Consistency: formal argumentation

If X is endogenous:

- If *n* grows the OLS estimator converges to the wrong value.
  - $\rightarrow$  OLS is inconsistent

Consider the standard model  $y = X\beta + \varepsilon$  and the OLS estimator


$$b = (X'X)^{-1}X'y = (X'X)^{-1}X'(X\beta + \varepsilon)$$
$$= (X'X)^{-1}X'X\beta + (X'X)^{-1}X'\varepsilon$$
$$= \beta + (X'X)^{-1}X'\varepsilon$$

So, b can be split into

- **1** True parameter value  $\beta$
- ② Random deviation  $(X'X)^{-1}X'\varepsilon$

# - Zafus

## Distribution of estimator for different n, true value= 2



Lecture 4.2, Slide 6 of 12, Erasmus School of Economics

## Asymptotic properties

What happens to *b* as  $n \to \infty$ ?

Recall:  $b = \beta + (X'X)^{-1}X'\varepsilon$ 

- ullet  $\beta$  is constant
- Elements of (X'X) and  $X'\varepsilon$  are sums over observations:

$$X'X = \begin{pmatrix} \sum_{i=1}^{n} x_{1i}^{2} & \sum_{i=1}^{n} x_{1i}x_{2i} & \dots & \sum_{i=1}^{n} x_{1i}x_{ki} \\ \sum_{i=1}^{n} x_{1i}x_{2i} & \sum_{i=1}^{n} x_{2i}^{2} & \dots & \sum_{i=1}^{n} x_{2i}x_{ki} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{i=1}^{n} x_{ki}x_{1i} & \sum_{i=1}^{n} x_{ki}x_{2i} & \dots & \sum_{i=1}^{n} x_{ki}^{2} \end{pmatrix}, X'\varepsilon = \begin{pmatrix} \sum_{i=1}^{n} x_{1i}\varepsilon_{i} \\ \sum_{i=1}^{n} x_{2i}\varepsilon_{i} \\ \vdots \\ \sum_{i=1}^{n} x_{ki}\varepsilon_{i} \end{pmatrix}$$

 $\rightarrow$  these diverge as  $n \rightarrow \infty$ 

# Asymptotic properties

Rewrite  $b = \beta + (\frac{1}{n}X'X)^{-1}(\frac{1}{n}X'\varepsilon)$ 

- $(\frac{1}{n}X'X)$  is an average  $\rightarrow$  in general converges to, say, Q
- $(\frac{1}{n}X'\varepsilon)$  also converges in general

## Consistency result:

b converges to  $\beta$  as  $n \to \infty$  if

- $\bullet$   $\frac{1}{n}X'X$  converges to Q, and
- $Q^{-1}$  exists, and
- **3**  $\frac{1}{n}X'\varepsilon$  converges to 0
  - ▶ No correlation between X and  $\varepsilon$  (for large n)
  - ► *X* is exogenous

X endogenous: b does not converge to  $\beta$ !

Ezafus,

Lecture 4.2, Slide 9 of 12, Erasmus School of Economics

# OLS in presence of endogeneity

### If X endogenous

- X correlated with  $\varepsilon$
- ullet OLS estimator for eta is not consistent
- Even with in infinite amount of data: OLS does not give useful estimates

## Small sample properties

So far we discussed what happens for  $n \to \infty$ 

#### Test

Why can't we derive the bias?

To obtain the bias

need to evaluate

$$E[b] = E[(X'X)^{-1}X'y] = E[(X'X)^{-1}X'(X\beta + \varepsilon)]$$
$$= E[\beta + (X'X)^{-1}X'\varepsilon] = \beta + \underbrace{E[(X'X)^{-1}X'\varepsilon]}_{=?}.$$


- X is stochastic
- cannot simplify final expectation (without further assumptions)

Ezafus,


Lecture 4.2, Slide 10 of 12, Erasmus School of Economics

## **TRAINING EXERCISE 4.2**

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).



"Solving endogeneity": Graphical representation



What have we so far?

- Endogeneity is a common problem
- Endogeneity causes OLS to be inconsistent
- Estimation requires another estimation technique

Leafus,

Lecture 4.3, Slide 2 of 12, Erasmus School of Economics

## Instrumental variable estimation


- Z variables are instruments if
  - Z and X are correlated
  - ightharpoonup Z does not correlate with  $\varepsilon$
- ullet Correlation between instruments and y is only due to X

$$Cov(Z, y) = Cov(Z, X\beta + \varepsilon) = Cov(Z, X\beta) + \underbrace{Cov(Z, \varepsilon)}_{=0}$$
  
=  $Cov(Z, X)\beta$ 

• Use instruments to estimate  $\beta$ 

# "Solving endogeneity": Graphical representation

- lacktriangle Use Z to decompose X in explained and unexplained part
- 2 Effect size of explained part on y equals  $\beta$
- Unexplained part is added to error term



Endogeneity is solved as

- X unexplained not correlated with X explained
- X explained is exogenous



Lecture 4.3, Slide 5 of 12, Erasmus School of Economics

# **Properties 2SLS**

- Variance of  $b_{2SLS}$ : Var $[b_{2SLS}] = \sigma^2 (X'H_ZX)^{-1}$
- Estimating  $\sigma^2$ :
  - $\hat{\sigma}^2 = \frac{1}{n-k} (y Xb_{2SLS})'(y Xb_{2SLS})$
  - ▶ Do not use residuals (or reported standard errors) of second stage regression!

Derivation of variance (use  $Var[\varepsilon] = \sigma^2 I$ ):

$$b_{2SLS} = (X'H_{Z}X)^{-1}X'H_{Z}y = (X'H_{Z}X)^{-1}X'H_{Z}(X\beta + \varepsilon)$$

$$= \beta + (X'H_{Z}X)^{-1}X'H_{Z}\varepsilon$$

$$Var[b_{2SLS}] = Var[(X'H_{Z}X)^{-1}X'H_{Z}\varepsilon]$$

$$= (X'H_{Z}X)^{-1}X'H_{Z}Var[\varepsilon] ((X'H_{Z}X)^{-1}X'H_{Z})'$$

$$= (X'H_{Z}X)^{-1}X'H_{Z}(\sigma^{2}I)H'_{Z}X(X'H_{Z}X)^{-1}$$

$$= \sigma^{2}(X'H_{Z}X)^{-1}X'H_{Z}H'_{Z}X(X'H_{Z}X)^{-1} = \sigma^{2}(X'H_{Z}X)^{-1}$$

Lecture 4.3, Slide 7 of 12, Erasmus School of Economics

### 2SLS in matrix notation

Given model

$$y = X\beta + \varepsilon$$
,  $Var[\varepsilon] = \sigma^2 I$ 

and instruments Z

**1** Regress X on Z to get explained part:

▶ Model:  $X = Z\gamma + \eta$ 

▶ OLS estimate:  $(Z'Z)^{-1}Z'X$ 

Fitted value:  $\hat{X} = \underbrace{Z(Z'Z)^{-1}Z'}_{H_Z}X = H_ZX$ 

**2** Regress y on  $\hat{X}$ :

$$b_{2SLS} = (\hat{X}'\hat{X})^{-1}\hat{X}'y$$
  
=  $(X'H'_ZH_ZX)^{-1}X'H'_Zy$   
=  $(X'H_ZX)^{-1}X'H_Zy$ 

Use:  $H_Z = H'_Z = H'_Z H_Z$ 



Lecture 4.3, Slide 6 of 12, Erasmus School of Economics

## Properties of 2SLS

- 2SLS is consistent if (when  $n \to \infty$ )
  - ▶ Z and  $\varepsilon$  not correlated:  $\frac{1}{n}Z'\varepsilon \to 0$
  - ▶ Z not multicollinear:  $\frac{1}{n}Z'Z \rightarrow Q_{ZZ}$ , and  $Q_{ZZ}$  invertible
  - ▶ X and Z sufficiently correlated:  $\frac{1}{n}X'Z \rightarrow Q_{XZ}$ , and  $Q_{ZZ}$  rank k

Sketch of proof:

$$b_{2SLS} = \beta + (X'H_{Z}X)^{-1}X'H_{Z}\varepsilon = \beta + (X'Z(Z'Z)^{-1}Z'X)^{-1}X'Z(Z'Z)^{-1}Z'\varepsilon$$

$$= \beta + \underbrace{(\frac{1}{n}X'Z(\frac{1}{n}Z'Z)^{-1}\frac{1}{n}Z'X)^{-1}}_{(Q_{XZ}Q_{ZZ}^{-1}Q'_{XZ})^{-1}}\underbrace{\frac{1}{n}X'Z(\frac{1}{n}Z'Z)^{-1}}_{Q_{XZ}Q_{ZZ}^{-1}}\underbrace{\frac{1}{n}Z'\varepsilon}_{0}$$

## Finding instruments

#### What are good instruments?

- All exogenous variables in X (incl. constant)
- Other instruments are always needed:
  - ► At least one for every endogenous variable
  - ▶ Want: strong correlation between Z and X
  - ▶ Need: no correlation between Z and  $\varepsilon$

Ezafus,

Lecture 4.3, Slide 9 of 12, Erasmus School of Economics

## **Summary**

### If X is in fact exogenous

- OLS and 2SLS both consistent
- Variance OLS smaller than variance 2SLS!
- $\rightarrow$  Use OLS

### If X is endogenous

- 2SLS is consistent
- OLS inconsistent
- $\rightarrow$  Use 2SLS

# **Examples of instruments**

Explain obtained grade using attendance:

Potential instruments:

- Travel time home to university
- Policy change to obligatory attendance

#### Test

What variable would be an instrument for price when modeling consumer sales of ice cream using sales =  $\alpha + \beta$ price +  $\varepsilon$ ?

Potential instruments?


- Prices of raw materials (valid)
- **2** Competitor prices (direct influence on sales, so part of  $\varepsilon$ )
- **3** Outside temperature (direct influence on sales, so part of  $\varepsilon$ )

Lecture 4.3, Slide 10 of 12, Erasmus School of Economics

## **TRAINING EXERCISE 4.3**

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).

- Erafus



## Testing the validity of instruments

Valid instruments satisfy three conditions

- There are enough instruments
  - $\rightarrow$  Easy! Just count.
- - $\rightarrow$  Check significance of instruments in first stage regression
- **3** Instruments are not correlated with  $\varepsilon$ 
  - $\rightarrow$  Perform Sargan test

- Eafing

## Outline

#### Given

- Model  $y = X\beta + \varepsilon$
- Instruments *Z*

Two important things to test

- 2 satisfies assumptions for instruments?
- X exogenous or endogenous?

Erafus

Lecture 4.4, Slide 2 of 1, Erasmus School of Economics

## Test correlation Z vs. X

- X<sub>1</sub> potentially endogenous variables
- $X_2$  exogenous variables
- $Z = (Z^*, X_2)$  instruments

First-stage regression: apply OLS to  $X_1 = Z^*\gamma_1 + X_2\gamma_2 + \eta$ 

#### Test

Why does 2SLS require  $\gamma_1 \neq 0$ ?

If  $\gamma_1 \approx 0$ :

- $\hat{X}_1 \approx X_2 \hat{\gamma}_2$ 
  - $ightarrow \hat{X}_1$  almost perfectly correlated with  $X_2$
- (Extremely) large estimation uncertainty

Test for sufficient correlation:

• Test  $H_0: \gamma_1 = 0$  in first-stage regression.

(Zafus

# Sargan test

Ingredients:

• Model:  $y = X\beta + \varepsilon$ 

• Explanatory variables:  $X = (X_1, X_2)$  $X_1$  (endogenous),  $X_2$  (exogenous)

• Instruments:  $Z = (Z^*, X_2)$ 

Null hypothesis ( $H_0$ ): Correlation Z and  $\varepsilon$  equals 0

Test procedure:

• Rewrite to  $H_0: \delta = 0$  in

$$\varepsilon = Z\delta + \xi$$

ullet  $\varepsilon$  cannot be observed

ightarrow Estimate arepsilon using 2SLS

Erafus

Lecture 4.4, Slide 5 of 1, Erasmus School of Economics

# Notes on the Sargan test

- Test only works when there are "too many" instruments (m > k)
- At least k of the instruments should be valid
- Test cannot indicate which instruments are invalid!

# Sargan test

Procedure:

- **1** Use Z to obtain 2SLS estimator  $b_{2SLS}$  for  $\beta$
- 2 Calculate  $e_{2SLS} = y Xb_{2SLS}$
- **3** Regress  $e_{2SLS}$  on Z
- $nR^2 \approx \chi^2(m-k)$  under  $H_0$  (valid instruments)
  - ► *m* instruments in *Z*
  - ▶ k explanatory variables in X

#### Test

The Sargan test requires m > k. What happens when m = k?

Ezafus,

Lecture 4.4, Slide 6 of 1, Erasmus School of Economics

# Testing for exogeneity of variables – Hausman test

#### Intuition:

- Use the instruments to split potentially endogenous variables into
  - a guaranteed exogenous part
  - 2 a potentially endogenous part
- Check whether the endogenous and exogenous part affect *y* differently.

# Hausman test – procedure

### Ingredients:

• Explanatory variables:  $X = (X_1, X_2)$ 

• Potentially endogenous:  $X_1$  ( $k_1$  variables)

• Exogenous variables: X<sub>2</sub> (k<sub>2</sub> variables)

• Instruments: Z

Null hypothesis  $(H_0)$ :  $X_1$  is exogenous

### Formal procedure:

**2** Regress  $X_1$  on  $Z \rightarrow$  calculate residuals V

 $\odot$  Regress e on X and V

•  $nR^2 \approx \chi^2(k_1)$  under  $H_0$  of exogeneity

Ezafus,

Lecture 4.4, Slide 9 of 1, Erasmus School of Economics

## TRAINING EXERCISE 4.4

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).

(zafus

Lecture 4.4, Slide 10 of 1, Erasmus School of Economics

Erasmus School of Economics

# **MOOC** Econometrics


Lecture 4.5 on Endogeneity: Application

Dennis Fok

**Erasmus University Rotterdam** 

# Ezafus,

# Correlation of GPA with participation



# **Application**

#### Setting:

- Online learning platform
- Grade Point Average (GPA) in MOOC on engineering
- $\bullet$  Impact of preparatory mathematics course
  - $\rightarrow$  participation is voluntary!

#### Data statistics:

- 1000 learners
- 48.8% male
- 33.7% participated in prep course
- Average GPA 5.94 (on 10 point scale)

(Zafins

Lecture 4.5, Slide 2 of 15, Erasmus School of Economics

# Correlation vs. regression

## Seems positive impact

- How large?
- Significant?
- Correction for male vs. female?
- $\rightarrow$  Need econometric model!

## **OLS** estimation

Regress GPA on

Constant

Gender: dummy variable (male=1, female=0)

3 Participation: dummy variable (yes=1, no=0)

Dependent variable: GPA

Sample size: 1000

	Coefficient	Standard error	t-statistic
Constant	5.77	0.034	169.87
Gender	-0.21	0.044	-4.82
<b>Participation</b>	0.82	0.047	17.59
$R^2$	0.24		

Ezafus,

Lecture 4.5, Slide 5 of 15, Erasmus School of Economics

# Over- or underestimation by OLS?

If prep course participation is endogenous

- OLS is inconsistent
- OLS does not estimate causal effect of prep course

#### Test

What omitted factor would lead OLS to <u>over</u>estimate the impact of the preparatory course?

Ezafus,

## Discussion of OLS

Should we trust the OLS estimates?

- → No, participation likely endogenous!
- Learners self-select for prep course
- Omitted factors (characteristics of learners) relate to this selection
- Same characteristics may relate to GPA

Ezafus

Lecture 4.5, Slide 6 of 15, Erasmus School of Economics

# Over- or underestimation by OLS?

#### Overestimation

Omitted factor: Motivation
 High motivation → Get high GPA & Take course

#### Underestimation

Omitted factor: Mathematics level
 High level → Get high GPA & Do not take course

#### Net effect:

- Difficult to judge
- Depends on importance of effects
- Also depends on other variables (age?)

(zafus

## Consistent estimation

- Use two-stage least squares (2SLS)
- Need instruments!

#### Test

What variable can you think of that qualifies as instrument for participation?

Leafus

Lecture 4.5, Slide 9 of 15, Erasmus School of Economics

#### Instruments

Finding instruments

• be creative! ... and lucky

Here

- Learners get email invitation for prep course
- Platform email problem: some did not get email
- Variable

$$\mathsf{Email} = \begin{cases} 0 & \text{if email not received} \\ 1 & \text{if email received} \end{cases}$$

is perfect instrument if

- ▶ Email problem is random
- ► Invitation affects participation

## Instruments

Instruments should...

- relate to prep course participation
- not affect GPA

Many learner specific variables, such as

- Intelligence (IQ-score)
- Number of MOOCs followed before
- Age of learner

are likely not valid!

→ All will impact performance directly!

Erafus

Lecture 4.5, Slide 10 of 15, Erasmus School of Economics

# First-stage regression

Explain participation using all instruments (constant, gender, email)

Dependent variable: Participation Sample size: 1000

		Coefficient	Standard error	t-statistic
	Constant	0.10	0.023	4.41
	Gender	0.05	0.027	1.80
	Email	0.41	0.027	<u>15.35</u>
	$R^2$	0.20		

ightarrow Email affects participation significantly

## 2SLS estimation

Dependent variable: GPA

Sample size: 1000

Instruments used: Constant, Gender, Email

	Coefficient	Standard error	t-statistic
Constant	5.95	0.048	123.54
Gender	-0.17	0.048	-3.59
<b>Participation</b>	0.24	0.115	2.09
$R^2$	0.13		

- Prep course still has significant positive impact
- Effect size decreased (from 0.82 (OLS) to 0.24 (2SLS))
- 2SLS increases variance
  - ▶ Only acceptable when Participation is endogenous
  - ▶ Perform Hausman test

Erafus

Lecture 4.5, Slide 13 of 15, Erasmus School of Economics

## **TRAINING EXERCISE 4.5**

- Train yourself by making the training exercise (see the website).
- After making this exercise, check your answers by studying the webcast solution (also available on the website).

Ezafus,

Lecture 4.5, Slide 15 of 15, Erasmus School of Economics

# Hausman test ( $H_0$ : Participation is exogenous)

Dependent variable: Residuals from OLS

Sample size: 1000

	Coefficient	Standard error	t-statistic
Constant	0.18	0.044	4.02
Gender	0.04	0.044	0.93
Participation	-0.58	0.105	-5.55
First-stage residuals $(v)$	0.72	0.117	6.17
$R^2$	0.0368		

- Test-statistic:  $nR^2 = 1000 \times 0.0368 = 36.8$
- Reject  $H_0$  (critical value from  $\chi^2(1)$ : 3.8)
- Participation is endogenous
- 2SLS is needed

Erafus

Lecture 4.5, Slide 14 of 15, Erasmus School of Economics