A theorem of de Moivre and Laplace



The setting: $X_1, X_2, ..., X_n, ... \sim Bernoulli(p)$

The setting: $X_1, X_2, ..., X_n, ... \sim Bernoulli(p)$ $S_n = X_1 + \cdots + X_n \sim Binomial(n, p)$

The setting: $X_1, X_2, \ldots, X_n, \ldots \sim \text{Bernoulli}(p)$ $S_n = X_1 + \cdots + X_n \sim \text{Binomial}(n, p)$ $S_n^* = \frac{S_n - E(S_n)}{\sqrt{\text{Var}(S_n)}}$

The setting: $X_1, X_2, ..., X_n, ... \sim Bernoulli(p)$

$$S_n = X_1 + \cdots + X_n \sim \text{Binomial}(n, p)$$

$$S_n^* = \frac{S_n - E(S_n)}{\sqrt{Var(S_n)}} = \frac{S_n - np}{\sqrt{np(1-p)}}$$

The setting:
$$X_1, X_2, ..., X_n, ... \sim \text{Bernoulli}(p)$$

$$S_n = X_1 + \dots + X_n \sim \text{Binomial}(n, p)$$

$$S_n^* = \frac{S_n - E(S_n)}{\sqrt{\text{Var}(S_n)}} = \frac{S_n - np}{\sqrt{np(1-p)}}$$

The de Moivre-Laplace limit theorem

Suppose a < b. Then, asymptotically, as $n \to \infty$, we have

$$\mathbf{P}\{a < S_n^* \le b\} \to \int_a^b \phi(x) \, dx$$

The setting:
$$X_1, X_2, ..., X_n, ... \sim \text{Bernoulli}(p)$$


$$S_n = X_1 + ... + X_n \sim \text{Binomial}(n, p)$$

$$S_n^* = \frac{S_n - E(S_n)}{\sqrt{\text{Var}(S_n)}} = \frac{S_n - np}{\sqrt{np(1-p)}}$$

The de Moivre-Laplace limit theorem

Suppose a < b. Then, asymptotically, as $n \to \infty$, we have

$$\mathbf{P}\{a < S_n^* \le b\} \to \int_0^b \varphi(x) \, dx$$



The setting:
$$X_1, X_2, ..., X_n, ... \sim \text{Bernoulli}(p)$$


$$S_n = X_1 + \cdots + X_n \sim \text{Binomial}(n, p)$$

$$S_n^* = \frac{S_n - E(S_n)}{\sqrt{Var(S_n)}} = \frac{S_n - np}{\sqrt{np(1-p)}}$$

The de Moivre-Laplace limit theorem

Suppose a < b. Then, asymptotically, as $n \to \infty$, we have

$$\mathbf{P}\{a < S_n^* \le b\} \to \int_a^b \Phi(x) \, dx = \Phi(b) - \Phi(a)$$





Slogan

Binomial probabilities (viewed in the proper scale) are governed approximately by the area under the bell curve.