

The ancient art of sieves





Boole's inequality

Boole's inequality

$$\mathbf{P}\left(\bigcup_j A_j\right) \leq \sum_j \mathbf{P}(A_j)$$

Boole's inequality

$$\mathbf{P}\left(\bigcup_j A_j\right) \leq \sum_j \mathbf{P}(A_j)$$

Two regions where the bound becomes interesting:

Boole's inequality

$$\mathbf{P}\left(\bigcup_j A_j\right) \leq \sum_j \mathbf{P}(A_j)$$

Two regions where the bound becomes interesting:

- ✧ The sum on the right is small (but not zero).

Boole's inequality

$$\mathbf{P}\left(\bigcup_j A_j\right) \leq \sum_j \mathbf{P}(A_j)$$

Two regions where the bound becomes interesting:

- ✧ The sum on the right is small (but not zero).
- ✧ The sum on the right is large (but not one).

Boole's inequality

$$\mathbf{P}\left(\bigcup_j A_j\right) \leq \sum_j \mathbf{P}(A_j)$$

Two regions where the bound becomes interesting:

- ✧ The sum on the right is small (but not zero).
- ✧ The sum on the right is large (but not one).

Some **colourful** terminology:

- ✧ **Bad** events: A_j
- ✧ **Good** events: $(A_j)^c$

Boole's inequality

$$\mathbf{P}\left(\bigcup_j A_j\right) \leq \sum_j \mathbf{P}(A_j)$$

Two regions where the bound becomes interesting:

- ✧ The sum on the right is small (but not zero).
- ✧ The sum on the right is large (but not one).

Some **colourful** terminology:

- ✧ **Bad** events: A_j
- ✧ **Good** events: $(A_j)^c$

$\bigcup_j A_j :=$ a bad event occurs

Boole's inequality

$$\mathbf{P}\left(\bigcup_j A_j\right) \leq \sum_j \mathbf{P}(A_j)$$

Two regions where the bound becomes interesting:

- ✧ The sum on the right is small (but not zero).
- ✧ The sum on the right is large (but not one).

Some **colourful** terminology:

- ✧ **Bad** events: A_j
- ✧ **Good** events: $(A_j)^c$

$\bigcup_j A_j :=$ a bad event occurs

$\bigcap_j A_j^c :=$ no bad event occurs

When Boole's bound is near zero

Some colourful terminology:

- ❖ `Bad' events: A_j
- ❖ `Good' events: $(A_j)^c$

$$\mathbf{P}\left(\bigcup_j A_j\right) \leq \sum_j \mathbf{P}(A_j)$$

When Boole's bound is near zero

Some colourful terminology:

- ✧ `Bad' events: A_j
- ✧ `Good' events: $(A_j)^c$

$$\mathbf{P}\left(\bigcup_j A_j\right) \leq \sum_j \mathbf{P}(A_j)$$

If $\sum_j \mathbf{P}(A_j) < \epsilon$ then $\mathbf{P}\left(\bigcap_j A_j^c\right) = 1 - \mathbf{P}\left(\bigcup_j A_j\right) > 1 - \epsilon$.

When Boole's bound is near zero

Some colourful terminology:

- ✧ `Bad' events: A_j
- ✧ `Good' events: $(A_j)^c$

$$\mathbf{P}\left(\bigcup_j A_j\right) \leq \sum_j \mathbf{P}(A_j)$$

If $\sum_j \mathbf{P}(A_j) < \epsilon$ then $\mathbf{P}\left(\bigcap_j A_j^c\right) = 1 - \mathbf{P}\left(\bigcup_j A_j\right) > 1 - \epsilon$.

Slogan

If Boole's bound is near zero then *most* outcomes are good.

When Boole's bound is less than one

Some colourful terminology:

- ❖ `Bad' events: A_j
- ❖ `Good' events: $(A_j)^c$

$$\mathbf{P}\left(\bigcup_j A_j\right) \leq \sum_j \mathbf{P}(A_j)$$

When Boole's bound is less than one

Some colourful terminology:

- ✧ `Bad' events: A_j
- ✧ `Good' events: $(A_j)^c$

$$\mathbf{P}\left(\bigcup_j A_j\right) \leq \sum_j \mathbf{P}(A_j)$$

If $\sum_j \mathbf{P}(A_j) < 1$ then $\mathbf{P}\left(\bigcap_j A_j^c\right) = 1 - \mathbf{P}\left(\bigcup_j A_j\right) > 0$.

When Boole's bound is less than one

Some colourful terminology:

- ✧ `Bad' events: A_j
- ✧ `Good' events: $(A_j)^c$

$$\mathbf{P}\left(\bigcup_j A_j\right) \leq \sum_j \mathbf{P}(A_j)$$

If $\sum_j \mathbf{P}(A_j) < 1$ then $\mathbf{P}\left(\bigcap_j A_j^c\right) = 1 - \mathbf{P}\left(\bigcup_j A_j\right) > 0$.

Slogan: Boole's sieve

If Boole's bound is less than one then there *exist* good outcomes.

Embedding a cube in a two-coloured sphere
