

Welcome back! If you found this question useful,
don't forget to vote both the question and the answers up.

close this message

Showing a Normal and a Chi square are independent

Asked 3 years, 7 months ago Modified 1 year, 7 months ago Viewed 1k times



0



Student's t distribution is defined as the ratio of a standard normally distributed random variable and the square root of a Chi-square distributed random variable divided by its degrees of freedom, given that they are independent. In formulas one can write $\frac{Z}{\sqrt{\frac{U}{df}}}$, where

Z is $N(0, 1)$ and U is χ_{df}^2 .

In showing that this statement is true, I arrived at the point in which I have $\frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2$ and $\frac{\bar{X}-\mu}{\frac{\sigma}{\sqrt{n}}} \sim N(0, 1)$. Then, following the definition, we would have that

$$\frac{\frac{\bar{X}-\mu}{\frac{\sigma}{\sqrt{n}}}}{\sqrt{\frac{\frac{(n-1)S^2}{\sigma^2}}{n-1}}}$$

is distributed as a t_{n-1} . But I am stuck at how to prove that these two random variables are independent between them. We covered a result about independence in the case of two Chi-square random variables and I thought of seeing the standard Normal as the square of a Chi-square random variable but I am afraid of it being mathematically sacrilegious.

Do you have any hint?

distributions

self-study

normal-distribution

t-distribution

chi-squared-distribution

Share Cite Edit Follow Flag

edited Feb 8, 2021 at 16:50



kjetil b halvorsen ♦

67.1k

29

148

494

asked Feb 6, 2019 at 23:36



PhDing

2,520

6

33

60

2 Answers

Sorted by:

Highest score (default)



This is a brute force solution requiring just multivariable calculus.

3

It suffices to prove that the sample mean

Welcome back! If you found this question useful,
don't forget to vote both the question and the answers up.

close this message

and the sample variance



$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

are independent. Thus, it suffices to prove that the sample mean \bar{X} is independent of the vector

$$(X_1 - \bar{X}, \dots, X_n - \bar{X}).$$

Moreover, since

$$\begin{aligned} \sum_{i=1}^n (X_i - \bar{X}) &= \sum_{i=1}^n X_i - \sum_{i=1}^n \bar{X} \\ &= n\bar{X} - n\bar{X} \\ &= 0, \end{aligned}$$

and hence

$$X_1 - \bar{X} = - \sum_{i=2}^n (X_i - \bar{X}),$$

it follows that $X_1 - \bar{X}$ can be recovered from just knowing $(X_2 - \bar{X}, \dots, X_n - \bar{X})$.

Thus, it suffices to prove that the sample mean \bar{X} is independent from

$$(X_2 - \bar{X}, \dots, X_n - \bar{X}).$$

Now consider the joint density

$$\begin{aligned} f_{(X_1, \dots, X_n)}(x_1, \dots, x_n) &= (2\pi\sigma^2)^{-n/2} \exp\left(-\sum_{i=1}^n \frac{1}{2} \left(\frac{x_i - \mu}{\sigma}\right)^2\right) \\ &= (2\pi\sigma^2)^{-n/2} \exp\left(-\sum_{i=1}^n \frac{1}{2} \left(\frac{x_i - \bar{x}}{\sigma}\right)^2 - \frac{n}{2} \left(\frac{\bar{x} - \mu}{\sigma}\right)^2\right) \\ &= \underbrace{(2\pi\sigma^2)^{-n/2}}_{\text{constant}} \underbrace{\exp\left(-\sum_{i=1}^n \frac{1}{2} \left(\frac{x_i - \bar{x}}{\sigma}\right)^2\right)}_{\text{depends only on } (x_2 - \bar{x}, \dots, x_n - \bar{x})} \underbrace{\exp\left(-\frac{n}{2} \left(\frac{\bar{x} - \mu}{\sigma}\right)^2\right)}_{\text{depends only on } \bar{x}} \end{aligned}$$

To get from (X_1, \dots, X_n) to $(\bar{X}, X_2 - \bar{X}, \dots, X_n - \bar{X})$, consider the diffeomorphism $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ given by

$$T(x_1, \dots, x_n) = (\bar{x}, x_2 - \bar{x}, \dots, x_n - \bar{x}).$$

Welcome back! If you found this question useful,
don't forget to vote both the question and the answers up.

[close this message](#)

which is also clearly differentiable). Up to transpose, the Jacobian matrix of T is

$$DT(x_1, \dots, x_n) = \begin{bmatrix} 1/n & 1/n & 1/n & \cdots & 1/n \\ -1/n & (n-1)/n & -1/n & \cdots & -1/n \\ -1/n & -1/n & (n-1)/n & \cdots & -1/n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -1/n & -1/n & -1/n & \cdots & (n-1)/n. \end{bmatrix},$$

which doesn't depend on x_1, \dots, x_n . Thus, the determinant of DT is some constant C . Now the joint density of $(\bar{X}, X_2 - \bar{X}, \dots, X_n - \bar{X})$ satisfies

$$f_{(\bar{X}, X_2 - \bar{X}, \dots, X_n - \bar{X})}(y_1, \dots, y_n) = |C| f_{(X_1, \dots, X_n)}(T^{-1}(y_1, \dots, y_n))$$

which factors as a function of y_1 times a function of (y_2, \dots, y_n) by what was shown above.

Therefore, \bar{X} and $(X_2 - \bar{X}, \dots, X_n - \bar{X})$ are independent.

Share Cite Edit Follow Flag

edited Feb 7, 2019 at 2:02

answered Feb 7, 2019 at 1:13



Artem Mavrin

3,577 2 16 27



I'll provide a hint to your self-study question: A corollary of a classic statistical theorem states that if $\mathbf{x} \sim N_p(\boldsymbol{\mu}, \sigma^2 \mathbf{I})$, then \mathbf{Bx} and $\mathbf{x}'\mathbf{Ax}$ are independent if and only if \mathbf{BA} is equal to the zero matrix. So, perhaps you could write the numerator as \mathbf{Bx} and the denominator as $\mathbf{x}'\mathbf{Ax}$ and work from there?



Share Cite Edit Follow Flag

answered Feb 7, 2019 at 0:53



StatsStudent

10.5k 4 39 70



I knew a slightly different result but this seems to be the one I need! Thanks! – [PhDing](#) Feb 7, 2019 at 4:23



If this hint worked for you, please accept the answer. Thank you! – [StatsStudent](#) Feb 7, 2019 at 15:10



Sure, I'll try later! – [PhDing](#) Feb 7, 2019 at 15:59



I am fine with the denominator, which can be rewritten as $\frac{X - \mu e'}{\sigma} \frac{ee'}{n-1} \frac{X - \mu e}{\sigma}$, and has the required form. I am still stuck with the numerator because it is already a Normal, then the only thing I can think of is multiplying by an identity matrix but this is not working. – [PhDing](#) Feb 7, 2019 at 20:23



Think about redefining a new variable Z_i . Does that help? – [StatsStudent](#) Feb 7, 2019 at 22:20

Welcome back! If you found this question useful,
don't forget to vote both the question and the answers up.
[close this message](#)
