

Advice from a magician — a view from a sample point

Events A_1, A_2, \dots, A_n

Advice from a magician — a view from a sample point

Events A_1, A_2, \dots, A_n

$$X_j = \begin{cases} 1 & \text{if } A_j \text{ occurs,} \\ 0 & \text{otherwise.} \end{cases}$$

Advice from a magician — a view from a sample point

Events A_1, A_2, \dots, A_n

$$X_j = \begin{cases} 1 & \text{if } A_j \text{ occurs,} \\ 0 & \text{otherwise.} \end{cases}$$

$$S_k := \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} \mathbf{P}(A_{j_1} \cap A_{j_2} \cap \dots \cap A_{j_k}) \quad (1 \leq k \leq n)$$

Advice from a magician — a view from a sample point

Events A_1, A_2, \dots, A_n

$$X_j = \begin{cases} 1 & \text{if } A_j \text{ occurs,} \\ 0 & \text{otherwise.} \end{cases} \quad S_k := \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} \mathbf{P}(A_{j_1} \cap A_{j_2} \cap \dots \cap A_{j_k}) \quad (1 \leq k \leq n)$$

Inclusion–exclusion: The distribution of the number of occurrences of A_1, \dots, A_n is given by

$$\mathbf{P}\{X_1 + \dots + X_n = k\} = \binom{k}{k} S_k - \binom{k+1}{k} S_{k+1} + \dots + (-1)^j \binom{k+j}{k} S_{k+j} + \dots + (-1)^{n-k} S_n$$

Advice from a magician — a view from a sample point

Events A_1, A_2, \dots, A_n

$$X_j = \begin{cases} 1 & \text{if } A_j \text{ occurs,} \\ 0 & \text{otherwise.} \end{cases} \quad S_k := \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} \mathbf{P}(A_{j_1} \cap A_{j_2} \cap \dots \cap A_{j_k}) \quad (1 \leq k \leq n)$$

Inclusion–exclusion: The distribution of the number of occurrences of A_1, \dots, A_n is given by

$$\mathbf{P}\{X_1 + \dots + X_n = k\} = \binom{k}{k} S_k - \binom{k+1}{k} S_{k+1} + \dots + (-1)^j \binom{k+j}{k} S_{k+j} + \dots + (-1)^{n-k} S_n$$

Fix any k and select any sample point ω .

How much does ω contribute to the left and to the right of the equation?

Advice from a magician — a view from a sample point

Events A_1, A_2, \dots, A_n

$$X_j = \begin{cases} 1 & \text{if } A_j \text{ occurs,} \\ 0 & \text{otherwise.} \end{cases} \qquad S_k := \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} \mathbf{P}(A_{j_1} \cap A_{j_2} \cap \dots \cap A_{j_k}) \quad (1 \leq k \leq n)$$

Inclusion–exclusion: The distribution of the number of occurrences of A_1, \dots, A_n is given by

$$\mathbf{P}\{X_1 + \dots + X_n = k\} = \binom{k}{k} S_k - \binom{k+1}{k} S_{k+1} + \dots + (-1)^j \binom{k+j}{k} S_{k+j} + \dots + (-1)^{n-k} S_n$$

Fix any k and select any sample point ω .

How much does ω contribute to the left and to the right of the equation?

The selected sample point ω will lie in a certain number, say, l of the sets A_j :

Advice from a magician — a view from a sample point

Events A_1, A_2, \dots, A_n

$$X_j = \begin{cases} 1 & \text{if } A_j \text{ occurs,} \\ 0 & \text{otherwise.} \end{cases} \quad S_k := \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} \mathbf{P}(A_{j_1} \cap A_{j_2} \cap \dots \cap A_{j_k}) \quad (1 \leq k \leq n)$$

Inclusion–exclusion: The distribution of the number of occurrences of A_1, \dots, A_n is given by

$$\mathbf{P}\{X_1 + \dots + X_n = k\} = \binom{k}{k} S_k - \binom{k+1}{k} S_{k+1} + \dots + (-1)^j \binom{k+j}{k} S_{k+j} + \dots + (-1)^{n-k} S_n$$

Fix any k and select any sample point ω .

How much does ω contribute to the left and to the right of the equation?

The selected sample point ω will lie in a certain number, say, l of the sets A_j :

- * Identify the sets: suppose ω is in $A_{j_1}, A_{j_2}, \dots, A_{j_l}$.

Advice from a magician — a view from a sample point

Events A_1, A_2, \dots, A_n

$$X_j = \begin{cases} 1 & \text{if } A_j \text{ occurs,} \\ 0 & \text{otherwise.} \end{cases} \quad S_k := \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} \mathbf{P}(A_{j_1} \cap A_{j_2} \cap \dots \cap A_{j_k}) \quad (1 \leq k \leq n)$$

Inclusion–exclusion: The distribution of the number of occurrences of A_1, \dots, A_n is given by

$$\mathbf{P}\{X_1 + \dots + X_n = k\} = \binom{k}{k} S_k - \binom{k+1}{k} S_{k+1} + \dots + (-1)^j \binom{k+j}{k} S_{k+j} + \dots + (-1)^{n-k} S_n$$

Fix any k and select any sample point ω .

How much does ω contribute to the left and to the right of the equation?

The selected sample point ω will lie in a certain number, say, l of the sets A_j :

- ✦ Identify the sets: suppose ω is in $A_{j_1}, A_{j_2}, \dots, A_{j_l}$.
- ✦ Then $X_1 + X_2 + \dots + X_n = l$.

Advice from a magician — a view from a sample point

Events A_1, A_2, \dots, A_n

$$X_j = \begin{cases} 1 & \text{if } A_j \text{ occurs,} \\ 0 & \text{otherwise.} \end{cases} \quad S_k := \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} \mathbf{P}(A_{j_1} \cap A_{j_2} \cap \dots \cap A_{j_k}) \quad (1 \leq k \leq n)$$

Inclusion–exclusion: The distribution of the number of occurrences of A_1, \dots, A_n is given by

$$\mathbf{P}\{X_1 + \dots + X_n = k\} = \binom{k}{k} S_k - \binom{k+1}{k} S_{k+1} + \dots + (-1)^j \binom{k+j}{k} S_{k+j} + \dots + (-1)^{n-k} S_n$$

Fix any k and select any sample point ω .

How much does ω contribute to the left and to the right of the equation?

The selected sample point ω will lie in a certain number, say, l of the sets A_j :

- ✦ Identify the sets: suppose ω is in $A_{j_1}, A_{j_2}, \dots, A_{j_l}$.
- ✦ Then $X_1 + X_2 + \dots + X_n = l$.
- ✦ Three cases: (1) $l < k$. (2) $l = k$. (3) $l > k$.