

Physics takes a gamble!

The science behind Monte Carlo methods

---



# A chance-driven computation

---

**Given:** a bounded function  $f(\mathbf{x}) = f(x_1, \dots, x_D)$  on the unit cube  $[0, 1]^D$  with  $|f(\mathbf{x})| \leq 1$



**Compute:**  $J = \int \cdots \int_{[0,1]^D} f(\mathbf{x}) \, d\mathbf{x}$

**Select**  $\mathbf{X} = (X_1, \dots, X_D)$  **at random** from the D-dimensional cube  $[0, 1]^D$

This means:  $X_1, \dots, X_D$  are independent and are each uniformly distributed in the unit interval.


$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

**Evaluate**  $Y = f(\mathbf{X})$





$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$






$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

$$\mathbb{E}(f(\mathbf{X})) = J$$

$$\text{Var}(f(\mathbf{X})) \leq 1$$






$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

$$\mathbf{E}(f(\mathbf{X})) = J$$

$$\text{Var}(f(\mathbf{X})) \leq 1$$




**Generate** an independent, random sample of points in the unit cube  $[0, 1]^D$ :  $\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \dots, \mathbf{X}^{(n)}$



$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

$$\mathbf{E}(f(\mathbf{X})) = J$$

$$\text{Var}(f(\mathbf{X})) \leq 1$$



**Generate** an independent, random sample of points in the unit cube  $[0, 1]^D$ :  $\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \dots, \mathbf{X}^{(n)}$


**Evaluate:**  $Y_1 = f(\mathbf{X}^{(1)}), Y_2 = f(\mathbf{X}^{(2)}), \dots, Y_n = f(\mathbf{X}^{(n)})$



$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

$$\mathbb{E}(f(\mathbf{X})) = J$$

$$\text{Var}(f(\mathbf{X})) \leq 1$$



**Generate** an independent, random sample of points in the unit cube  $[0, 1]^D$ :  $\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \dots, \mathbf{X}^{(n)}$

**Evaluate:**  $Y_1 = f(\mathbf{X}^{(1)}), Y_2 = f(\mathbf{X}^{(2)}), \dots, Y_n = f(\mathbf{X}^{(n)})$


What do we know about these values?



$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

$$\mathbb{E}(f(\mathbf{X})) = \mathbf{J}$$

$$\text{Var}(f(\mathbf{X})) \leq 1$$



**Generate** an independent, random sample of points in the unit cube  $[0, 1]^D$ :  $\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \dots, \mathbf{X}^{(n)}$

**Evaluate:**  $Y_1 = f(\mathbf{X}^{(1)}), Y_2 = f(\mathbf{X}^{(2)}), \dots, Y_n = f(\mathbf{X}^{(n)})$

What do we know about these values?


The induced sample  $Y_1, \dots, Y_n$  constitutes a sequence of repeated independent trials with common expectation  $\mathbf{J}$  and variance bounded above by 1.



$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

$$\mathbf{E}(f(\mathbf{X})) = \mathbf{J}$$

$$\text{Var}(f(\mathbf{X})) \leq 1$$



**Generate** an independent, random sample of points in the unit cube  $[0, 1]^D$ :  $\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \dots, \mathbf{X}^{(n)}$

**Evaluate:**  $Y_1 = f(\mathbf{X}^{(1)}), Y_2 = f(\mathbf{X}^{(2)}), \dots, Y_n = f(\mathbf{X}^{(n)})$

What do we know about these values?

The induced sample  $Y_1, \dots, Y_n$  constitutes a sequence of repeated independent trials with common expectation  $\mathbf{J}$  and variance bounded above by 1.


The law of large numbers beckons!



$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

$$\mathbb{E}(f(\mathbf{X})) = J$$

$$\text{Var}(f(\mathbf{X})) \leq 1$$



**Generate** an independent, random sample of points in the unit cube  $[0, 1]^D$ :  $\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \dots, \mathbf{X}^{(n)}$

**Evaluate:**  $Y_1 = f(\mathbf{X}^{(1)}), Y_2 = f(\mathbf{X}^{(2)}), \dots, Y_n = f(\mathbf{X}^{(n)})$

What do we know about these values?

The induced sample  $Y_1, \dots, Y_n$  constitutes a sequence of repeated independent trials with common expectation  $J$  and variance bounded above by 1.

The law of large numbers beckons!


**Form:**  $S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$



$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

$$\mathbf{E}(f(\mathbf{X})) = \mathbf{J}$$

$$\text{Var}(f(\mathbf{X})) \leq 1$$



**Generate** an independent, random sample of points in the unit cube  $[0, 1]^D$ :  $\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \dots, \mathbf{X}^{(n)}$

**Evaluate:**  $Y_1 = f(\mathbf{X}^{(1)}), Y_2 = f(\mathbf{X}^{(2)}), \dots, Y_n = f(\mathbf{X}^{(n)})$

What do we know about these values?

The induced sample  $Y_1, \dots, Y_n$  constitutes a sequence of repeated independent trials with common expectation  $\mathbf{J}$  and variance bounded above by 1.

The law of large numbers beckons!

**Form:**  $S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$


$$\mathbf{E}(S_n) = \mathbf{E}(Y_1) + \cdots + \mathbf{E}(Y_n)$$



$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

$$\mathbf{E}(f(\mathbf{X})) = \mathbf{J}$$

$$\text{Var}(f(\mathbf{X})) \leq 1$$



**Generate** an independent, random sample of points in the unit cube  $[0, 1]^D$ :  $\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \dots, \mathbf{X}^{(n)}$

**Evaluate:**  $Y_1 = f(\mathbf{X}^{(1)}), Y_2 = f(\mathbf{X}^{(2)}), \dots, Y_n = f(\mathbf{X}^{(n)})$

What do we know about these values?

The induced sample  $Y_1, \dots, Y_n$  constitutes a sequence of repeated independent trials with common expectation  $\mathbf{J}$  and variance bounded above by 1.

The law of large numbers beckons!

**Form:**  $S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$


$$\mathbf{E}(S_n) = \mathbf{E}(Y_1) + \cdots + \mathbf{E}(Y_n) = n\mathbf{J}$$



$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

$$\mathbf{E}(f(\mathbf{X})) = \mathbf{J}$$

$$\text{Var}(f(\mathbf{X})) \leq 1$$



**Generate** an independent, random sample of points in the unit cube  $[0, 1]^D$ :  $\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \dots, \mathbf{X}^{(n)}$

**Evaluate:**  $Y_1 = f(\mathbf{X}^{(1)}), Y_2 = f(\mathbf{X}^{(2)}), \dots, Y_n = f(\mathbf{X}^{(n)})$

What do we know about these values?

The induced sample  $Y_1, \dots, Y_n$  constitutes a sequence of repeated independent trials with common expectation  $\mathbf{J}$  and variance bounded above by 1.

The law of large numbers beckons!

**Form:**  $S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$

$$\mathbf{E}(S_n) = \mathbf{E}(Y_1) + \cdots + \mathbf{E}(Y_n) = n\mathbf{J}$$


$$\text{Var}(S_n) = \text{Var}(Y_1) + \cdots + \text{Var}(Y_n)$$



$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

$$\mathbf{E}(f(\mathbf{X})) = \mathbf{J}$$

$$\text{Var}(f(\mathbf{X})) \leq 1$$



**Generate** an independent, random sample of points in the unit cube  $[0, 1]^D$ :  $\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \dots, \mathbf{X}^{(n)}$

**Evaluate:**  $Y_1 = f(\mathbf{X}^{(1)}), Y_2 = f(\mathbf{X}^{(2)}), \dots, Y_n = f(\mathbf{X}^{(n)})$

What do we know about these values?

The induced sample  $Y_1, \dots, Y_n$  constitutes a sequence of repeated independent trials with common expectation  $\mathbf{J}$  and variance bounded above by 1.

The law of large numbers beckons!

**Form:**  $S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$


$$\mathbf{E}(S_n) = \mathbf{E}(Y_1) + \cdots + \mathbf{E}(Y_n) = n\mathbf{J}$$

$$\text{Var}(S_n) = \text{Var}(Y_1) + \cdots + \text{Var}(Y_n) \leq n \cdot 1 = n$$



$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} \mathbf{E}(f(\mathbf{X})) &= J \\ \text{Var}(f(\mathbf{X})) &\leq 1 \end{aligned}$$



**Generate** an independent, random sample of points in the unit cube  $[0, 1]^D$ :  $\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \dots, \mathbf{X}^{(n)}$

**Evaluate:**  $Y_1 = f(\mathbf{X}^{(1)}), Y_2 = f(\mathbf{X}^{(2)}), \dots, Y_n = f(\mathbf{X}^{(n)})$

What do we know about these values?

The induced sample  $Y_1, \dots, Y_n$  constitutes a sequence of repeated independent trials with common expectation  $J$  and variance bounded above by 1.

The law of large numbers beckons!

**Form:**  $S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$

$$\mathbf{E}(S_n) = \mathbf{E}(Y_1) + \cdots + \mathbf{E}(Y_n) = nJ$$


$$\text{Var}(S_n) = \text{Var}(Y_1) + \cdots + \text{Var}(Y_n) \leq n \cdot 1 = n$$

**Estimate** the unknown  $J$  by the sample mean  $S_n/n$



$$\mathbf{X} \sim p(\mathbf{x}) = u(x_1) \times \cdots \times u(x_D) = \begin{cases} 1 & \text{if } \mathbf{x} \in [0, 1]^D \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} \mathbf{E}(f(\mathbf{X})) &= J \\ \text{Var}(f(\mathbf{X})) &\leq 1 \end{aligned}$$



**Generate** an independent, random sample of points in the unit cube  $[0, 1]^D$ :  $\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \dots, \mathbf{X}^{(n)}$

**Evaluate:**  $Y_1 = f(\mathbf{X}^{(1)}), Y_2 = f(\mathbf{X}^{(2)}), \dots, Y_n = f(\mathbf{X}^{(n)})$

What do we know about these values?

The induced sample  $Y_1, \dots, Y_n$  constitutes a sequence of repeated independent trials with common expectation  $J$  and variance bounded above by 1.

The law of large numbers beckons!

**Form:**  $S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$


$$\mathbf{E}(S_n) = \mathbf{E}(Y_1) + \cdots + \mathbf{E}(Y_n) = nJ$$

$$\text{Var}(S_n) = \text{Var}(Y_1) + \cdots + \text{Var}(Y_n) \leq n \cdot 1 = n$$

Monte Carlo method

**Estimate** the unknown  $J$  by the sample mean  $S_n/n$






$$S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$$

$$\mathbf{E}(S_n) = n\mathbf{J}$$

$$\text{Var}(S_n) \leq n$$






$$S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$$

$$\mathbf{E}(S_n) = n\mathbf{J}$$

$$\text{Var}(S_n) \leq n$$

How well does the sample mean  $S_n/n$  estimate the unknown integral  $\mathbf{J}$ ?





$$S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$$


$$\mathbf{E}(S_n) = n\mathbf{J}$$

$$\text{Var}(S_n) \leq n$$

How well does the sample mean  $S_n/n$  estimate the unknown integral  $\mathbf{J}$ ?

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mathbf{J}\right| > \epsilon\right\}$$





$$S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$$


$$\mathbf{E}(S_n) = n\mathbf{J}$$

$$\text{Var}(S_n) \leq n$$

How well does the sample mean  $S_n/n$  estimate the unknown integral  $\mathbf{J}$ ?

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mathbf{J}\right| > \epsilon\right\} = \mathbf{P}\{|S_n - n\mathbf{J}| > n\epsilon\}$$





$$S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$$


$$\mathbf{E}(S_n) = nJ$$

$$\text{Var}(S_n) \leq n$$

How well does the sample mean  $S_n/n$  estimate the unknown integral  $J$ ?

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - J\right| > \epsilon\right\} = \mathbf{P}\{|S_n - nJ| > n\epsilon\} \leq \frac{\text{Var } S_n}{(n\epsilon)^2}$$





$$S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$$


$$\mathbf{E}(S_n) = nJ$$

$$\text{Var}(S_n) \leq n$$

How well does the sample mean  $S_n/n$  estimate the unknown integral  $J$ ?

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - J\right| > \epsilon\right\} = \mathbf{P}\{|S_n - nJ| > n\epsilon\} \leq \frac{\text{Var } S_n}{(n\epsilon)^2} \leq \frac{n}{n^2\epsilon^2}$$





$$S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$$


$$\mathbf{E}(S_n) = nJ$$

$$\text{Var}(S_n) \leq n$$

How well does the sample mean  $S_n/n$  estimate the unknown integral  $J$ ?

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - J\right| > \epsilon\right\} = \mathbf{P}\{|S_n - nJ| > n\epsilon\} \leq \frac{\text{Var } S_n}{(n\epsilon)^2} \leq \frac{n}{n^2\epsilon^2} = \frac{1}{n\epsilon^2}$$





$$S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$$

$$\mathbf{E}(S_n) = nJ$$

$$\text{Var}(S_n) \leq n$$


How well does the sample mean  $S_n/n$  estimate the unknown integral  $J$ ?

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - J\right| > \epsilon\right\} = \mathbf{P}\{|S_n - nJ| > n\epsilon\} \leq \frac{\text{Var } S_n}{(n\epsilon)^2} \leq \frac{n}{n^2\epsilon^2} = \frac{1}{n\epsilon^2}$$

The sample mean  $S_n/n$  estimate the unknown integral  $J$  with an absolute error of no more than  $\epsilon$  and a confidence of at least  $1 - \delta$  if the sample size  $n$  satisfies

$$\frac{1}{n\epsilon^2} \leq \delta \quad \text{—or—} \quad n \geq \frac{1}{\epsilon^2\delta}$$





$$S_n = Y_1 + \cdots + Y_n = f(\mathbf{X}^{(1)}) + \cdots + f(\mathbf{X}^{(n)})$$

$$\mathbf{E}(S_n) = nJ$$

$$\text{Var}(S_n) \leq n$$

How well does the sample mean  $S_n/n$  estimate the unknown integral  $J$ ?

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - J\right| > \epsilon\right\} = \mathbf{P}\{|S_n - nJ| > n\epsilon\} \leq \frac{\text{Var } S_n}{(n\epsilon)^2} \leq \frac{n}{n^2\epsilon^2} = \frac{1}{n\epsilon^2}$$

The sample mean  $S_n/n$  estimate the unknown integral  $J$  with an absolute error of no more than  $\epsilon$  and a confidence of at least  $1 - \delta$  if the sample size  $n$  satisfies

$$\frac{1}{n\epsilon^2} \leq \delta \quad \text{—or—} \quad n \geq \frac{1}{\epsilon^2\delta}$$

To illustrate: if  $n = 10^6$  then the estimate error will be less than 1% ( $\epsilon = 0.01$ ) and the confidence in excess of 99% ( $\delta = 0.01$ )!