

# The ballot problem

W. A. Whitworth, *Messenger of Math*, 1878

---

In a run-off election between two candidates for class prefect, Jane gets 6 votes and Bob gets 4 votes. The ballots are counted in random order. What is the chance that Jane leads Bob at every step of the count?



# The ballot problem

W. A. Whitworth, *Messenger of Math*, 1878

In a run-off election between two candidates for class prefect, Jane gets 6 votes and Bob gets 4 votes. The ballots are counted in random order. What is the chance that Jane leads Bob at every step of the count?

Ballot order	J	B	J	B	B	J	J	J	B	J
Jane's running total	1	1	2	2	2	3	4	5	5	6
Bob's running total	0	1	1	2	3	3	3	3	4	4



# The ballot problem

W. A. Whitworth, *Messenger of Math*, 1878

In a run-off election between two candidates for class prefect, Jane gets 6 votes and Bob gets 4 votes. The ballots are counted in random order. What is the chance that Jane leads Bob at every step of the count?

Ballot order	J	B	J	B	B	J	J	J	B	J
Jane's running total	1	1	2	2	2	3	4	5	5	6
Bob's running total	0	1	1	2	3	3	3	3	4	4



# The ballot problem

W. A. Whitworth, *Messenger of Math*, 1878

5149

In a run-off election between two candidates for class prefect, Jane gets 6 votes and Bob gets 4 votes. The ballots are counted in random order. What is the chance that Jane leads Bob at every step of the count?

Ballot order	J	B	J	B	B	J	J	J	B	J
Jane's running total	1	1	2	2	2	3	4	5	5	6
Bob's running total	0	1	1	2	3	3	3	3	4	4



# The ballot problem

W. A. Whitworth, *Messenger of Math*, 1878

$n$   
51

$m$   
49

In a run-off election between two candidates for class prefect, Jane gets 6 votes and Bob gets 4 votes. The ballots are counted in random order. What is the chance that Jane leads Bob at every step of the count?

Ballot order	J	B	J	B	B	J	J	J	B	J
Jane's running total	1	1	2	2	2	3	4	5	5	6
Bob's running total	0	1	1	2	3	3	3	3	4	4



An useful idea for sequential problems: condition on a step (first, last)

---

1	2	3						n+m-1	n+m
---	---	---	--	--	--	--	--	-------	-----



An useful idea for sequential problems: condition on a step (first, last)

---

1	2	3						$n+m-1$	$n+m$
---	---	---	--	--	--	--	--	---------	-------

- *Sample space*: all sequences of  $n$  Js and  $m$  Bs.



An useful idea for sequential problems: condition on a step (first, last)

---

1	2	3						$n+m-1$	$n+m$
---	---	---	--	--	--	--	--	---------	-------

- *Sample space*: all sequences of  $n$  Js and  $m$  Bs.
- *Probability measure*: combinatorial, all atoms have equal probability.



An useful idea for sequential problems: condition on a step (first, last)

---

1	2	3						$n+m-1$	$n+m$
---	---	---	--	--	--	--	--	---------	-------

- *Sample space*: all sequences of  $n$  Js and  $m$  Bs.
- *Probability measure*: combinatorial, all atoms have equal probability.
- *Events of interest*:



An useful idea for sequential problems: condition on a step (first, last)

---


1	2	3						$n+m-1$	$n+m$
---	---	---	--	--	--	--	--	---------	-------

- *Sample space*: all sequences of  $n$  Js and  $m$  Bs.
- *Probability measure*: combinatorial, all atoms have equal probability.
- *Events of interest*:
  - $H :=$  Jane leads at each step of the count.



An useful idea for sequential problems: condition on a step (first, last)

---




- *Sample space*: all sequences of  $n$  Js and  $m$  Bs.
- *Probability measure*: combinatorial, all atoms have equal probability.
- *Events of interest*:
  - $H :=$  Jane leads at each step of the count.



An useful idea for sequential problems: condition on a step (first, last)

---




- *Sample space:* all sequences of  $n$  Js and  $m$  Bs.
- *Probability measure:* combinatorial, all atoms have equal probability.
- *Events of interest:*
  - $H :=$  Jane leads at each step of the count.



An useful idea for sequential problems: condition on a step (first, last)

---




- *Sample space*: all sequences of  $n$  Js and  $m$  Bs.
- *Probability measure*: combinatorial, all atoms have equal probability.
- *Events of interest*:
  - $H :=$  Jane leads at each step of the count.



An useful idea for sequential problems: condition on a step (first, last)

---




- *Sample space*: all sequences of  $n$  Js and  $m$  Bs.
- *Probability measure*: combinatorial, all atoms have equal probability.
- *Events of interest*:
  - $H :=$  Jane leads at each step of the count.
  - $A :=$  Jane gets the last ballot.



# An useful idea for sequential problems: condition on a step (first, last)

Ballots for: (Jane, Bob)




- *Sample space:* all sequences of  $n$  Js and  $m$  Bs.
- *Probability measure:* combinatorial, all atoms have equal probability.
- *Events of interest:*
  - $H :=$  Jane leads at each step of the count.
  - $A :=$  Jane gets the last ballot.



# An useful idea for sequential problems: condition on a step (first, last)

Ballots for: (Jane, Bob)




$\mathbf{P(H)} =: P_{n,m}$

- *Sample space:* all sequences of  $n$  Js and  $m$  Bs.
- *Probability measure:* combinatorial, all atoms have equal probability.
- *Events of interest:*
  - $H :=$  Jane leads at each step of the count.
  - $A :=$  Jane gets the last ballot.



# An useful idea for sequential problems: condition on a step (first, last)

Ballots for: (Jane, Bob)



- *Sample space:* all sequences of  $n$  Js and  $m$  Bs.
- *Probability measure:* combinatorial, all atoms have equal probability.
- *Events of interest:*
  - $H :=$  Jane leads at each step of the count.
  - $A :=$  Jane gets the last ballot.


$$\mathbf{P}(H) =: P_{n,m}$$

$$\mathbf{P}(H \mid A) = P_{n-1,m}$$



# An useful idea for sequential problems: condition on a step (first, last)

Ballots for: (Jane, Bob)



- *Sample space:* all sequences of  $n$  Js and  $m$  Bs.
- *Probability measure:* combinatorial, all atoms have equal probability.
- *Events of interest:*
  - $H :=$  Jane leads at each step of the count.
  - $A :=$  Jane gets the last ballot.

$$\mathbf{P}(H) =: P_{n,m}$$


$$\mathbf{P}(H \mid A) = P_{n-1,m}$$

$$\mathbf{P}(H \mid A^c) = P_{n,m-1}$$



# An useful idea for sequential problems: condition on a step (first, last)

Ballots for: (Jane, Bob)



- *Sample space*: all sequences of  $n$  Js and  $m$  Bs.
- *Probability measure*: combinatorial, all atoms have equal probability.
- *Events of interest*:
  - $H$  := Jane leads at each step of the count.
  - $A$  := Jane gets the last ballot.

$$\mathbf{P}(H) =: P_{n,m}$$

$$\mathbf{P}(H \mid A) = P_{n-1,m}$$


$$\mathbf{P}(H \mid A^c) = P_{n,m-1}$$

$$\mathbf{P}(A) = \frac{\binom{n+m-1}{m}}{\binom{n+m}{m}} = \frac{n}{n+m}$$



# An useful idea for sequential problems: condition on a step (first, last)

Ballots for: (Jane, Bob)



- *Sample space*: all sequences of  $n$  Js and  $m$  Bs.
- *Probability measure*: combinatorial, all atoms have equal probability.
- *Events of interest*:
  - $H$  := Jane leads at each step of the count.
  - $A$  := Jane gets the last ballot.

$$\mathbf{P}(H) =: P_{n,m}$$

$$\mathbf{P}(H \mid A) = P_{n-1,m}$$

$$\mathbf{P}(H \mid A^c) = P_{n,m-1}$$

$$\mathbf{P}(A) = \frac{\binom{n+m-1}{m}}{\binom{n+m}{m}} = \frac{n}{n+m}$$

$$\mathbf{P}(A^c) = 1 - \mathbf{P}(A) = \frac{m}{n+m}$$