

# Quo vadis?

---



# Chebyshev and after: first steps

---



# Chebyshev and after: first steps

---

Repeated independent trials  $X_1, \dots, X_n, \dots$  with finite expectation  $\mu$



# Chebyshev and after: first steps

---

Repeated independent trials  $X_1, \dots, X_n, \dots$  with finite expectation  $\mu$   
(but no assumption of a finite variance)



# Chebyshev and after: first steps

---

Repeated independent trials  $X_1, \dots, X_n, \dots$  with finite expectation  $\mu$   
(but no assumption of a finite variance)

Sequence of sample means:  $\frac{S_n}{n} = \frac{X_1 + \dots + X_n}{n}$



# Chebyshev and after: first steps

---

Repeated independent trials  $X_1, \dots, X_n, \dots$  with finite expectation  $\mu$   
(but no assumption of a finite variance)

Sequence of sample means:  $\frac{S_n}{n} = \frac{X_1 + \dots + X_n}{n}$

The **weak** law of large numbers  
(Khinchin, 1929)



# Chebyshev and after: first steps

---

Repeated independent trials  $X_1, \dots, X_n, \dots$  with finite expectation  $\mu$   
(but no assumption of a finite variance)

Sequence of sample means:  $\frac{S_n}{n} = \frac{X_1 + \dots + X_n}{n}$

The **weak** law of large numbers  
(Khinchin, 1929)

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon\right\} \rightarrow 0 \quad (n \rightarrow \infty)$$



# Sample paths and a new question

---



# Sample paths and a new question

---

**Sample path:** trace the sample means in one sequence of tosses of a fair coin  $\frac{S_1}{1}, \frac{S_2}{2}, \dots, \frac{S_n}{n}, \dots$



# Sample paths and a new question

---

**Sample path:** trace the sample means in one sequence of tosses of a fair coin  $\frac{S_1}{1}, \frac{S_2}{2}, \dots, \frac{S_n}{n}, \dots$


How does the sample path behave?



# Sample paths and a new question

**Sample path:** trace the sample means in one sequence of tosses of a fair coin  $\frac{S_1}{1}, \frac{S_2}{2}, \dots, \frac{S_n}{n}, \dots$

How does the sample path behave?






# Sample paths and a new question

**Sample path:** trace the sample means in one sequence of tosses of a fair coin  $\frac{S_1}{1}, \frac{S_2}{2}, \dots, \frac{S_n}{n}, \dots$

How does the sample path behave?



What are the chances that the sample mean **repeatedly** drifts away from 0.5 by more than 10%?  
By more than  $\epsilon$ ?



# The laws of large numbers

---

Repeated independent trials  $X_1, \dots, X_n, \dots$  with finite expectation  $\mu$

(but no assumption of a finite variance)

Sequence of sample means:  $\frac{S_n}{n} = \frac{X_1 + \dots + X_n}{n}$

The **weak** law of large numbers  
(Khinchin, 1929)

$$\mathbf{P}\left\{\left|\frac{S_n}{n} - \mu\right| > \epsilon\right\} \rightarrow 0 \quad (n \rightarrow \infty)$$



# The laws of large numbers

---

Repeated independent trials  $X_1, \dots, X_n, \dots$  with finite expectation  $\mu$

(but no assumption of a finite variance)

Sequence of sample means:  $\frac{S_n}{n} = \frac{X_1 + \dots + X_n}{n}$

Can the events  $\left\{ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right\}$  occur *infinitely often (i.o.)*?

The **weak** law of large numbers  
(Khinchin, 1929)

$$\mathbf{P} \left\{ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right\} \rightarrow 0 \quad (n \rightarrow \infty)$$



# The laws of large numbers

---

Repeated independent trials  $X_1, \dots, X_n, \dots$  with finite expectation  $\mu$

(but no assumption of a finite variance)

Sequence of sample means:  $\frac{S_n}{n} = \frac{X_1 + \dots + X_n}{n}$

Can the events  $\left\{ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right\}$  occur *infinitely often (i.o.)*?

The **weak** law of large numbers  
(Khinchin, 1929)

$$\mathbf{P} \left\{ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right\} \rightarrow 0 \quad (n \rightarrow \infty)$$

The **strong** law of large numbers  
(Kolmogorov, 1933)

$$\mathbf{P} \left\{ \left| \frac{S_n}{n} - \mu \right| > \epsilon \text{ i.o.} \right\} = 0$$