

Test your understanding

Relative ranks

Relative ranks

- ❖ Four athletes p, q, r , and s run a race and are ranked in order of finish. Suppose all outcomes are equally likely.

Relative ranks

- ❖ Four athletes p, q, r , and s run a race and are ranked in order of finish. Suppose all outcomes are equally likely.
 - ❖ *Sample space* Ω : the set of $4! = 24$ permutations of (p, q, r, s) .

Relative ranks

- ❖ Four athletes p, q, r , and s run a race and are ranked in order of finish. Suppose all outcomes are equally likely.
 - ❖ *Sample space* Ω : the set of $4! = 24$ permutations of (p, q, r, s) .
 - ❖ *Events*:

Relative ranks

- ✧ Four athletes p, q, r , and s run a race and are ranked in order of finish. Suppose all outcomes are equally likely.
 - ✧ *Sample space* Ω : the set of $4! = 24$ permutations of (p, q, r, s) .
 - ✧ *Events*:
 - ✧ $A := p$ is ahead of q .

Relative ranks

- ✧ Four athletes p, q, r , and s run a race and are ranked in order of finish. Suppose all outcomes are equally likely.
 - ✧ *Sample space* Ω : the set of $4! = 24$ permutations of (p, q, r, s) .
 - ✧ *Events*:
 - ✧ $A := p$ is ahead of q .
 - ✧ $B := r$ is ahead of s .

Relative ranks

- ❖ Four athletes p, q, r , and s run a race and are ranked in order of finish. Suppose all outcomes are equally likely.
 - ❖ *Sample space* Ω : the set of $4! = 24$ permutations of (p, q, r, s) .
 - ❖ *Events*:
 - ❖ $A := p$ is ahead of q .
 - ❖ $B := r$ is ahead of s .
 - ❖ *Probability measure*: Combinatorial setting with mass function assigning equal probability $1/24$ to each atom.

Relative ranks

- ❖ Four athletes p, q, r , and s run a race and are ranked in order of finish. Suppose all outcomes are equally likely.
 - ❖ *Sample space* Ω : the set of $4! = 24$ permutations of (p, q, r, s) .
 - ❖ *Events*:
 - ❖ $A := p$ is ahead of q .
 - ❖ $B := r$ is ahead of s .
 - ❖ *Probability measure*: Combinatorial setting with mass function assigning equal probability $1/24$ to each atom.
- ❖ Are A and B independent events?

Relative ranks

- ❖ Four athletes p, q, r , and s run a race and are ranked in order of finish. Suppose all outcomes are equally likely.

$$\mathbf{P}(A \cap B) \stackrel{?}{=} \mathbf{P}(A) \mathbf{P}(B)$$

- ❖ *Sample space* Ω : the set of $4! = 24$ permutations of (p, q, r, s) .
 - ❖ *Events*:
 - ❖ $A := p$ is ahead of q .
 - ❖ $B := r$ is ahead of s .
 - ❖ *Probability measure*: Combinatorial setting with mass function assigning equal probability $1/24$ to each atom.
- ❖ Are A and B independent events?

Relative ranks

- ❖ Four athletes p, q, r , and s run a race and are ranked in order of finish. Suppose all outcomes are equally likely.

- ❖ *Sample space* Ω : the set of $4! = 24$ permutations of (p, q, r, s) .

- ❖ *Events*:

- ❖ $A := p$ is ahead of q .

- ❖ $B := r$ is ahead of s .

- ❖ *Probability measure*: Combinatorial setting with mass function assigning equal probability $1/24$ to each atom.

- ❖ Are A and B independent events?

$$\mathbf{P}(A \cap B) \stackrel{?}{=} \mathbf{P}(A) \mathbf{P}(B)$$

#ways q can be placed to the right of p #ways r and s can be arranged in the remaining two positions

$$\mathbf{P}(A) = \mathbf{P}(B) = \frac{(3 + 2 + 1)2!}{4!} = \frac{1}{2}$$

Relative ranks

- ❖ Four athletes p, q, r , and s run a race and are ranked in order of finish. Suppose all outcomes are equally likely.

- ❖ *Sample space* Ω : the set of $4! = 24$ permutations of (p, q, r, s) .

- ❖ *Events*:

- ❖ $A := p$ is ahead of q .

- ❖ $B := r$ is ahead of s .

- ❖ *Probability measure*: Combinatorial setting with mass function assigning equal probability $1/24$ to each atom.

- ❖ Are A and B independent events?

$$\mathbf{P}(A \cap B) \stackrel{?}{=} \mathbf{P}(A) \mathbf{P}(B)$$

#ways q can be placed to the right of p #ways r and s can be arranged in the remaining two positions

$$\mathbf{P}(A) = \mathbf{P}(B) = \frac{(3 + 2 + 1)2!}{4!} = \frac{1}{2}$$

$$\mathbf{P}(A \cap B) = \frac{3 + 2 + 1}{4!} = \frac{1}{4}$$

Relative ranks

- ❖ Four athletes p, q, r , and s run a race and are ranked in order of finish. Suppose all outcomes are equally likely.

- ❖ *Sample space* Ω : the set of $4! = 24$ permutations of (p, q, r, s) .

- ❖ *Events*:

- ❖ $A := p$ is ahead of q .

- ❖ $B := r$ is ahead of s .

- ❖ *Probability measure*: Combinatorial setting with mass function assigning equal probability $1/24$ to each atom.

- ❖ Are A and B independent events?

$$\mathbf{P}(A \cap B) \stackrel{?}{=} \mathbf{P}(A) \mathbf{P}(B)$$

#ways q can be placed to the right of p #ways r and s can be arranged in the remaining two positions

$$\mathbf{P}(A) = \mathbf{P}(B) = \frac{(3 + 2 + 1)2!}{4!} = \frac{1}{2}$$

$$\mathbf{P}(A \cap B) = \frac{3 + 2 + 1}{4!} = \frac{1}{4} = \frac{1}{2} \cdot \frac{1}{2} = \mathbf{P}(A) \mathbf{P}(B)$$

Relative ranks

- ❖ Four athletes p, q, r , and s run a race and are ranked in order of finish. Suppose all outcomes are equally likely.

- ❖ *Sample space* Ω : the set of $4! = 24$ permutations of (p, q, r, s) .

- ❖ *Events*:

- ❖ $A := p$ is ahead of q .

- ❖ $B := r$ is ahead of s .

- ❖ *Probability measure*: Combinatorial setting with mass function assigning equal probability $1/24$ to each atom.

- ❖ Are A and B independent events?

$$\mathbf{P}(A \cap B) \stackrel{?}{=} \mathbf{P}(A) \mathbf{P}(B)$$

#ways q can be placed to the right of p #ways r and s can be arranged in the remaining two positions

$$\mathbf{P}(A) = \mathbf{P}(B) = \frac{(3 + 2 + 1)2!}{4!} = \frac{1}{2}$$

$$\mathbf{P}(A \cap B) = \frac{3 + 2 + 1}{4!} = \frac{1}{4} = \frac{1}{2} \cdot \frac{1}{2} = \mathbf{P}(A) \mathbf{P}(B)$$

A and B are independent.