


A chi-squared test

The principle:

Compare Burt's IQ data with a random sample generated from a normal distribution with mean 100 and variance 225.

A hypothesised normal distribution of IQs with mean 100 and variance 225

Random sample: X_1, \dots, X_n



$$S_n^{(1)} \sim \text{Binomial}(n, p_1)$$

$$S_n^{(2)} \sim \text{Binomial}(n, p_2)$$

.....

$$S_n^{(7)} \sim \text{Binomial}(n, p_7)$$

$$S_n^{(8)} \sim \text{Binomial}(n, p_8)$$

IQ bins j	$70 <$	70–80	80–90	90–100	100–110	110–120	120–130	>130
	1	2	3	4	5	6	7	8
Probabilities p_j	0.0228	0.0685	0.1613	0.2475	0.2475	0.1613	0.0685	0.0228

The key observation:

We anticipate that each of the values $S_n^{(k)}$ will take values near its expectation np_k and, viewed in the proper scale, will be approximately normally distributed around that point.

The key observation:

We anticipate that each of the values $S_n^{(k)}$ will take values near its expectation np_k and, viewed in the proper scale, will be approximately normally distributed around that point.

The genesis of a statistical test:

What principles should guide the construction of a test to check whether the values $S_n^{(k)}$ are appropriately close to their expectations np_k ?

The key observation:

We anticipate that each of the values $S_n^{(k)}$ will take values near its expectation np_k and, viewed in the proper scale, will be approximately normally distributed around that point.

The genesis of a statistical test:

What principles should guide the construction of a test to check whether the values $S_n^{(k)}$ are appropriately close to their expectations np_k ?

Two maxims of T. W. Körner:

- ✧ The test should be easy to apply.
- ✧ The method should not ignore certain pieces of data.