

# Test your understanding

---



# Test your understanding

---

- ❖ For what value of constant  $C$  is the function  $p(k) = C2^{-k}$  defined on the non-negative integers  $k = 0, 1, 2, \dots$  a distribution?



# Test your understanding

---

- ❖ For what value of constant  $C$  is the function  $p(k) = C2^{-k}$  defined on the non-negative integers  $k = 0, 1, 2, \dots$  a distribution?

Geometric series:  $\sum_{k=0}^{\infty} 2^{-k} = 1 + 2^{-1} + 2^{-2} + \dots = \frac{1}{1 - 2^{-1}} = 2$



# Test your understanding

---

- ❖ For what value of constant  $C$  is the function  $p(k) = C2^{-k}$  defined on the non-negative integers  $k = 0, 1, 2, \dots$  a distribution?

- $C$  is a positive normalisation constant:  $C = 1/2$ .
- This is the geometric distribution with parameter  $1/2$ .

Geometric series:  $\sum_{k=0}^{\infty} 2^{-k} = 1 + 2^{-1} + 2^{-2} + \dots = \frac{1}{1 - 2^{-1}} = 2$



# Test your understanding

---

- ❖ For what value of constant  $C$  is the function  $p(k) = C2^{-k}$  defined on the non-negative integers  $k = 0, 1, 2, \dots$  a distribution?

- $C$  is a positive normalisation constant:  $C = 1/2$ .
- This is the geometric distribution with parameter  $1/2$ .

- ❖ For what value of constant  $C$  is the function  $p(k) = Ce^{-|k|}$  defined as  $k$  varies over all integers a distribution?

Geometric series:  $\sum_{k=0}^{\infty} 2^{-k} = 1 + 2^{-1} + 2^{-2} + \dots = \frac{1}{1 - 2^{-1}} = 2$



# Test your understanding

---

- ❖ For what value of constant  $C$  is the function  $p(k) = C2^{-k}$  defined on the non-negative integers  $k = 0, 1, 2, \dots$  a distribution?

- $C$  is a positive normalisation constant:  $C = 1/2$ .
- This is the geometric distribution with parameter  $1/2$ .

Geometric series:  $\sum_{k=0}^{\infty} 2^{-k} = 1 + 2^{-1} + 2^{-2} + \dots = \frac{1}{1 - 2^{-1}} = 2$

- ❖ For what value of constant  $C$  is the function  $p(k) = Ce^{-|k|}$  defined as  $k$  varies over all integers a distribution?

$$\sum_{k=-\infty}^{\infty} e^{-|k|} = 1 + 2(e^{-1} + e^{-2} + \dots) = 1 + \frac{2e^{-1}}{1 - e^{-1}} = \frac{1 + e^{-1}}{1 - e^{-1}} = \frac{e + 1}{e - 1}$$



# Test your understanding

---

- ❖ For what value of constant  $C$  is the function  $p(k) = C2^{-k}$  defined on the non-negative integers  $k = 0, 1, 2, \dots$  a distribution?

- $C$  is a positive normalisation constant:  $C = 1/2$ .
- This is the geometric distribution with parameter  $1/2$ .

Geometric series:  $\sum_{k=0}^{\infty} 2^{-k} = 1 + 2^{-1} + 2^{-2} + \dots = \frac{1}{1 - 2^{-1}} = 2$

- ❖ For what value of constant  $C$  is the function  $p(k) = Ce^{-|k|}$  defined as  $k$  varies over all integers a distribution?

- $C = (e - 1)/(e + 1)$ .

$$\sum_{k=-\infty}^{\infty} e^{-|k|} = 1 + 2(e^{-1} + e^{-2} + \dots) = 1 + \frac{2e^{-1}}{1 - e^{-1}} = \frac{1 + e^{-1}}{1 - e^{-1}} = \frac{e + 1}{e - 1}$$



# Test your understanding

---

- ❖ For what value of constant  $C$  is the function  $p(k) = C2^{-k}$  defined on the non-negative integers  $k = 0, 1, 2, \dots$  a distribution?

- $C$  is a positive normalisation constant:  $C = 1/2$ .
- This is the geometric distribution with parameter  $1/2$ .

Geometric series:  $\sum_{k=0}^{\infty} 2^{-k} = 1 + 2^{-1} + 2^{-2} + \dots = \frac{1}{1 - 2^{-1}} = 2$

- ❖ For what value of constant  $C$  is the function  $p(k) = Ce^{-|k|}$  defined as  $k$  varies over all integers a distribution?

- $C = (e - 1)/(e + 1)$ .

$$\sum_{k=-\infty}^{\infty} e^{-|k|} = 1 + 2(e^{-1} + e^{-2} + \dots) = 1 + \frac{2e^{-1}}{1 - e^{-1}} = \frac{1 + e^{-1}}{1 - e^{-1}} = \frac{e + 1}{e - 1}$$

- ❖ Can the function  $p(k) = Ce^{-|k|} \cos \pi k$  defined as  $k$  varies over all integers be a distribution for any choice of constant  $C$ ?



# Test your understanding

---

- ❖ For what value of constant  $C$  is the function  $p(k) = C2^{-k}$  defined on the non-negative integers  $k = 0, 1, 2, \dots$  a distribution?

Geometric series:  $\sum_{k=0}^{\infty} 2^{-k} = 1 + 2^{-1} + 2^{-2} + \dots = \frac{1}{1 - 2^{-1}} = 2$

- $C$  is a positive normalisation constant:  $C = 1/2$ .
- This is the geometric distribution with parameter  $1/2$ .

- ❖ For what value of constant  $C$  is the function  $p(k) = Ce^{-|k|}$  defined as  $k$  varies over all integers a distribution?

$$\sum_{k=-\infty}^{\infty} e^{-|k|} = 1 + 2(e^{-1} + e^{-2} + \dots) = 1 + \frac{2e^{-1}}{1 - e^{-1}} = \frac{1 + e^{-1}}{1 - e^{-1}} = \frac{e + 1}{e - 1}$$

- $C = (e - 1)/(e + 1)$ .

- ❖ Can the function  $p(k) = Ce^{-|k|} \cos \pi k$  defined as  $k$  varies over all integers be a distribution for any choice of constant  $C$ ?

$$\cos \pi k = \begin{cases} -1 & \text{if } k \text{ is odd,} \\ +1 & \text{if } k \text{ is even.} \end{cases}$$



# Test your understanding

---

- ❖ For what value of constant  $C$  is the function  $p(k) = C2^{-k}$  defined on the non-negative integers  $k = 0, 1, 2, \dots$  a distribution?

Geometric series:  $\sum_{k=0}^{\infty} 2^{-k} = 1 + 2^{-1} + 2^{-2} + \dots = \frac{1}{1 - 2^{-1}} = 2$

- $C$  is a positive normalisation constant:  $C = 1/2$ .
- This is the geometric distribution with parameter  $1/2$ .

- ❖ For what value of constant  $C$  is the function  $p(k) = Ce^{-|k|}$  defined as  $k$  varies over all integers a distribution?

$$\sum_{k=-\infty}^{\infty} e^{-|k|} = 1 + 2(e^{-1} + e^{-2} + \dots) = 1 + \frac{2e^{-1}}{1 - e^{-1}} = \frac{1 + e^{-1}}{1 - e^{-1}} = \frac{e + 1}{e - 1}$$

- $C = (e - 1)/(e + 1)$ .

- ❖ Can the function  $p(k) = Ce^{-|k|} \cos \pi k$  defined as  $k$  varies over all integers be a distribution for any choice of constant  $C$ ?

$$\cos \pi k = \begin{cases} -1 & \text{if } k \text{ is odd,} \\ +1 & \text{if } k \text{ is even.} \end{cases}$$

- No. Positivity will be violated for any choice of  $C$  as  $p(k)$  alternates sign for odd and even values of  $k$ .