

# Families

---



# Families

---

- ❖ A family is known to have three children. Make the appropriate assumptions for the probability space.



# Families

---

- ❖ A family is known to have three children. Make the appropriate assumptions for the probability space.
  - ❖ *Sample space:*  $\Omega = \{bbb, bbg, bgb, bgg, gbb, gbg, ggb, ggg\}$ .



# Families

---

- ❖ A family is known to have three children. Make the appropriate assumptions for the probability space.
  - ❖ *Sample space:*  $\Omega = \{bbb, bbg, bgb, bgg, gbb, gbg, ggb, ggg\}$ .
  - ❖ *Events:*



# Families

---

- ❖ A family is known to have three children. Make the appropriate assumptions for the probability space.
  - ❖ *Sample space:*  $\Omega = \{bbb, bbg, bgb, bgg, gbb, gb g, ggb, ggg\}$ .
  - ❖ *Events:*
    - ❖  $A :=$  The family has at most one girl  $= \{bbb, bbg, bgb, gbb\}$ .



# Families

---

- ❖ A family is known to have three children. Make the appropriate assumptions for the probability space.
  - ❖ *Sample space:*  $\Omega = \{bbb, bbg, bgb, bgg, gbb, gb g, ggb, ggg\}$ .
  - ❖ *Events:*
    - ❖  $A :=$  The family has at most one girl  $= \{bbb, bbg, bgb, gbb\}$ .
    - ❖  $B :=$  The family has children of both sexes  $= \{bbg, bgb, bgg, gbb, gb g, ggb\}$ ;  $B^c = \{bbb, ggg\}$ .



# Families

---

- \* A family is known to have three children. Make the appropriate assumptions for the probability space.
  - \* *Sample space:*  $\Omega = \{bbb, bbg, bgb, bgg, gbb, gb g, ggb, ggg\}$ .
  - \* *Events:*
    - \*  $A :=$  The family has at most one girl  $= \{bbb, bbg, bgb, gbb\}$ .
    - \*  $B :=$  The family has children of both sexes  $= \{bbg, bgb, bgg, gbb, gb g, ggb\}$ ;  $B^c = \{bbb, ggg\}$ .
  - \* *Probability measure:* Combinatorial setting with mass function assigning equal probability  $1/8$  to each atom.



# Families

---

- \* A family is known to have three children. Make the appropriate assumptions for the probability space.
  - \* *Sample space:*  $\Omega = \{bbb, bbg, bgb, bgg, gbb, gb g, ggb, ggg\}$ .
  - \* *Events:*
    - \*  $A :=$  The family has at most one girl  $= \{bbb, bbg, bgb, gbb\}$ .
    - \*  $B :=$  The family has children of both sexes  $= \{bbg, bgb, bgg, gbb, gb g, ggb\}$ ;  $B^c = \{bbb, ggg\}$ .
  - \* *Probability measure:* Combinatorial setting with mass function assigning equal probability  $1/8$  to each atom.
- \* Are A and B independent events?



# Families

---

- \* A family is known to have three children. Make the appropriate assumptions for the probability space.
  - \* *Sample space:*  $\Omega = \{bbb, bbg, bgb, bgg, gbb, gbg, ggb, ggg\}$ .
  - \* *Events:*
    - \*  $A :=$  The family has at most one girl  $= \{bbb, bbg, bgb, gbb\}$ .
    - \*  $B :=$  The family has children of both sexes  $= \{bbg, bgb, bgg, gbb, gbg, ggb\}$ ;  $B^c = \{bbb, ggg\}$ .
  - \* *Probability measure:* Combinatorial setting with mass function assigning equal probability  $1/8$  to each atom.
- \* Are A and B independent events?

#outcomes with no girls ←      → #outcomes with one girl

$$\mathbf{P}(A) = \frac{1 + 3}{8} = \frac{1}{2}$$



# Families

---

- \* A family is known to have three children. Make the appropriate assumptions for the probability space.
  - \* *Sample space:*  $\Omega = \{bbb, bbg, bgb, bgg, gbb, gb g, ggb, ggg\}$ .
  - \* *Events:*
    - \*  $A :=$  The family has at most one girl  $= \{bbb, bbg, bgb, gbb\}$ .
    - \*  $B :=$  The family has children of both sexes  $= \{bbg, bgb, bgg, gbb, gb g, ggb\}$ ;  $B^c = \{bbb, ggg\}$ .
  - \* *Probability measure:* Combinatorial setting with mass function assigning equal probability  $1/8$  to each atom.
- \* Are  $A$  and  $B$  independent events?

#outcomes with no girls      #outcomes with one girl

$$\mathbf{P}(A) = \frac{1 + 3}{8} = \frac{1}{2}$$

$$\mathbf{P}(B^c) = \frac{2}{8} = \frac{1}{4}$$



# Families

---

- \* A family is known to have three children. Make the appropriate assumptions for the probability space.
- \* *Sample space:*  $\Omega = \{bbb, bbg, bgb, bgg, gbb, gbg, ggb, ggg\}$ .
- \* *Events:*
  - \*  $A :=$  The family has at most one girl  $= \{bbb, bbg, bgb, gbb\}$ .
  - \*  $B :=$  The family has children of both sexes  $= \{bbg, bgb, bgg, gbb, gbg, ggb\}$ ;  $B^c = \{bbb, ggg\}$ .
- \* *Probability measure:* Combinatorial setting with mass function assigning equal probability  $1/8$  to each atom.
- \* Are A and B independent events?

#outcomes with no girls ←      #outcomes with one girl →

$$\mathbf{P}(A) = \frac{1 + 3}{8} = \frac{1}{2}$$

$$\mathbf{P}(B^c) = \frac{2}{8} = \frac{1}{4}$$

$$\mathbf{P}(B) = 1 - \mathbf{P}(B^c) = 1 - \frac{1}{4} = \frac{3}{4}$$



# Families

---

- \* A family is known to have three children. Make the appropriate assumptions for the probability space.
- \* *Sample space:*  $\Omega = \{bbb, bbg, bgb, bgg, gbb, gbg, ggb, ggg\}$ .
- \* *Events:*
  - \*  $A :=$  The family has at most one girl  $= \{bbb, bbg, bgb, gbb\}$ .
  - \*  $B :=$  The family has children of both sexes  $= \{bbg, bgb, bgg, gbb, gbg, ggb\}$ ;  $B^c = \{bbb, ggg\}$ .
- \* *Probability measure:* Combinatorial setting with mass function assigning equal probability  $1/8$  to each atom.
- \* Are  $A$  and  $B$  independent events?

#outcomes with no girls      #outcomes with one girl

$$\mathbf{P}(A) = \frac{1 + 3}{8} = \frac{1}{2}$$

$$\mathbf{P}(B^c) = \frac{2}{8} = \frac{1}{4}$$

$$\mathbf{P}(B) = 1 - \mathbf{P}(B^c) = 1 - \frac{1}{4} = \frac{3}{4}$$

$$\mathbf{P}(A \cap B) = \frac{3}{8}$$



# Families

---

- \* A family is known to have three children. Make the appropriate assumptions for the probability space.
  - \* *Sample space:*  $\Omega = \{bbb, bbg, bgb, bgg, gbb, gb g, ggb, ggg\}$ .
  - \* *Events:*
    - \*  $A :=$  The family has at most one girl  $= \{bbb, bbg, bgb, gbb\}$ .
    - \*  $B :=$  The family has children of both sexes  $= \{bbg, bgb, bgg, gbb, gb g, ggb\}$ ;  $B^c = \{bbb, ggg\}$ .
  - \* *Probability measure:* Combinatorial setting with mass function assigning equal probability  $1/8$  to each atom.
- \* Are A and B independent events?

#outcomes with no girls      #outcomes with one girl

$$\mathbf{P}(A) = \frac{1 + 3}{8} = \frac{1}{2}$$

$$\mathbf{P}(B^c) = \frac{2}{8} = \frac{1}{4}$$

$$\mathbf{P}(B) = 1 - \mathbf{P}(B^c) = 1 - \frac{1}{4} = \frac{3}{4}$$

$$\mathbf{P}(A \cap B) = \frac{3}{8} = \frac{1}{2} \cdot \frac{3}{4} = \mathbf{P}(A) \mathbf{P}(B)$$



# Families

---

- \* A family is known to have three children. Make the appropriate assumptions for the probability space.
  - \* *Sample space:*  $\Omega = \{bbb, bbg, bgb, bgg, gbb, gbg, ggb, ggg\}$ .
  - \* *Events:*
    - \*  $A :=$  The family has at most one girl  $= \{bbb, bbg, bgb, gbb\}$ .
    - \*  $B :=$  The family has children of both sexes  $= \{bbg, bgb, bgg, gbb, gbg, ggb\}$ ;  $B^c = \{bbb, ggg\}$ .
  - \* *Probability measure:* Combinatorial setting with mass function assigning equal probability  $1/8$  to each atom.
- \* Are A and B independent events?

#outcomes with no girls ←      #outcomes with one girl →

$$\mathbf{P}(A) = \frac{1 + 3}{8} = \frac{1}{2}$$

$$\mathbf{P}(B^c) = \frac{2}{8} = \frac{1}{4}$$

$$\mathbf{P}(B) = 1 - \mathbf{P}(B^c) = 1 - \frac{1}{4} = \frac{3}{4}$$

$$\mathbf{P}(A \cap B) = \frac{3}{8} = \frac{1}{2} \cdot \frac{3}{4} = \mathbf{P}(A) \mathbf{P}(B)$$

A and B are independent.



# Families

---

- ❖ A family is known to have three children. Make the appropriate assumptions for the probability space.

- ❖ *Sample space:*  $\Omega = \{bbb, bbg, bgb, bgg, gbb, gbg, ggb, ggg\}$ .

- ❖ *Events:*

- ❖  $A :=$  The family has at most one girl  $= \{bbb, bbg, bgb, gbb\}$ .

- ❖  $B :=$  The family has children of both sexes  
 $= \{bbg, bgb, bgg, gbb, gbg, ggb\}$ ;  $B^c = \{bbb, ggg\}$ .

- ❖ *Probability measure:* Combinatorial setting with mass function assigning equal probability  $1/8$  to each atom.

- ❖ Are A and B independent events?

- ❖ What if the family has  $n$  children?

#outcomes with no girls ←      #outcomes with one girl →

$$\mathbf{P}(A) = \frac{1 + 3}{8} = \frac{1}{2}$$

$$\mathbf{P}(B^c) = \frac{2}{8} = \frac{1}{4}$$

$$\mathbf{P}(B) = 1 - \mathbf{P}(B^c) = 1 - \frac{1}{4} = \frac{3}{4}$$

$$\mathbf{P}(A \cap B) = \frac{3}{8} = \frac{1}{2} \cdot \frac{3}{4} = \mathbf{P}(A) \mathbf{P}(B)$$

A and B are independent.



# Families, reprise

---

- \* A family is known to have  $n$  children. Make the appropriate assumptions for the probability space.

- \* *Sample space*  $\Omega$ : all  $n$ -tuples of  $b$ 's and  $g$ 's.

- \* *Events*:

- \*  $A :=$  The family has at most one girl.

- \*  $B :=$  The family has children of both sexes;  $B^c :=$  the family has only boys or only girls.

- \* *Probability measure*: Combinatorial setting with mass function assigning equal probability  $1/2^n$  to each atom.

- \* Are  $A$  and  $B$  independent events?

#outcomes with no girls ←      → #outcomes with one girl

$$\mathbf{P}(A) = \frac{1 + 3}{8} = \frac{1}{2}$$

$$\mathbf{P}(B^c) = \frac{2}{8} = \frac{1}{4}$$

$$\mathbf{P}(B) = 1 - \mathbf{P}(B^c) = 1 - \frac{1}{4} = \frac{3}{4}$$

$$\mathbf{P}(A \cap B) = \frac{3}{8} = \frac{1}{2} \cdot \frac{3}{4} = \mathbf{P}(A) \mathbf{P}(B)$$



# Families, reprise

---

- \* A family is known to have  $n$  children. Make the appropriate assumptions for the probability space.

- \* *Sample space*  $\Omega$ : all  $n$ -tuples of  $b$ 's and  $g$ 's.

- \* *Events*:

- \*  $A :=$  The family has at most one girl.

- \*  $B :=$  The family has children of both sexes;  $B^c :=$  the family has only boys or only girls.

- \* *Probability measure*: Combinatorial setting with mass function assigning equal probability  $1/2^n$  to each atom.

- \* Are  $A$  and  $B$  independent events?

#outcomes with no girls ←      → #outcomes with one girl

$$\mathbf{P}(A) = \frac{1+n}{2^n}$$

$$\mathbf{P}(B^c) = \frac{2}{8} = \frac{1}{4}$$

$$\mathbf{P}(B) = 1 - \mathbf{P}(B^c) = 1 - \frac{1}{4} = \frac{3}{4}$$

$$\mathbf{P}(A \cap B) = \frac{3}{8} = \frac{1}{2} \cdot \frac{3}{4} = \mathbf{P}(A) \mathbf{P}(B)$$



# Families, reprise

---

- \* A family is known to have  $n$  children. Make the appropriate assumptions for the probability space.

- \* *Sample space*  $\Omega$ : all  $n$ -tuples of  $b$ 's and  $g$ 's.

- \* *Events*:

- \*  $A :=$  The family has at most one girl.

- \*  $B :=$  The family has children of both sexes;  $B^c :=$  the family has only boys or only girls.

- \* *Probability measure*: Combinatorial setting with mass function assigning equal probability  $1/2^n$  to each atom.

- \* Are  $A$  and  $B$  independent events?

#outcomes with no girls ←      → #outcomes with one girl

$$\mathbf{P}(A) = \frac{1 + n}{2^n}$$

$$\mathbf{P}(B^c) = \frac{2}{2^n} = \frac{1}{2^{n-1}} \quad \mathbf{P}(B) = 1 - \mathbf{P}(B^c) = 1 - \frac{1}{4} = \frac{3}{4}$$

$$\mathbf{P}(A \cap B) = \frac{3}{8} = \frac{1}{2} \cdot \frac{3}{4} = \mathbf{P}(A) \mathbf{P}(B)$$



# Families, reprise

---

- \* A family is known to have  $n$  children. Make the appropriate assumptions for the probability space.

- \* *Sample space*  $\Omega$ : all  $n$ -tuples of  $b$ 's and  $g$ 's.

- \* *Events*:

- \*  $A :=$  The family has at most one girl.

- \*  $B :=$  The family has children of both sexes;  $B^c :=$  the family has only boys or only girls.

- \* *Probability measure*: Combinatorial setting with mass function assigning equal probability  $1/2^n$  to each atom.

- \* Are  $A$  and  $B$  independent events?

#outcomes with no girls ←      → #outcomes with one girl

$$\mathbf{P}(A) = \frac{1 + n}{2^n}$$

$$\mathbf{P}(B^c) = \frac{2}{2^n} = \frac{1}{2^{n-1}} \quad \mathbf{P}(B) = 1 - \mathbf{P}(B^c) = 1 - \frac{1}{2^{n-1}}$$

$$\mathbf{P}(A \cap B) = \frac{3}{8} = \frac{1}{2} \cdot \frac{3}{4} = \mathbf{P}(A) \mathbf{P}(B)$$



# Families, reprise

---

- \* A family is known to have  $n$  children. Make the appropriate assumptions for the probability space.

- \* *Sample space*  $\Omega$ : all  $n$ -tuples of  $b$ 's and  $g$ 's.

- \* *Events*:

- \*  $A :=$  The family has at most one girl.

- \*  $B :=$  The family has children of both sexes;  $B^c :=$  the family has only boys or only girls.

- \* *Probability measure*: Combinatorial setting with mass function assigning equal probability  $1/2^n$  to each atom.

- \* Are  $A$  and  $B$  independent events?

#outcomes with no girls  $\leftarrow$   $\rightarrow$  #outcomes with one girl

$$\mathbf{P}(A) = \frac{1 + n}{2^n}$$

$$\mathbf{P}(B^c) = \frac{2}{2^n} = \frac{1}{2^{n-1}} \quad \mathbf{P}(B) = 1 - \mathbf{P}(B^c) = 1 - \frac{1}{2^{n-1}}$$

$$\mathbf{P}(A \cap B) = \frac{n}{2^n}$$



# Families, reprise

---

- \* A family is known to have  $n$  children. Make the appropriate assumptions for the probability space.

- \* *Sample space*  $\Omega$ : all  $n$ -tuples of  $b$ 's and  $g$ 's.

- \* *Events*:

- \*  $A :=$  The family has at most one girl.

- \*  $B :=$  The family has children of both sexes;  $B^c :=$  the family has only boys or only girls.

- \* *Probability measure*: Combinatorial setting with mass function assigning equal probability  $1/2^n$  to each atom.

- \* Are  $A$  and  $B$  independent events?

#outcomes with no girls ←      → #outcomes with one girl

$$\mathbf{P}(A) = \frac{1+n}{2^n}$$

$$\mathbf{P}(B^c) = \frac{2}{2^n} = \frac{1}{2^{n-1}} \quad \mathbf{P}(B) = 1 - \mathbf{P}(B^c) = 1 - \frac{1}{2^{n-1}}$$

$$\mathbf{P}(A \cap B) = \frac{n}{2^n}$$

Checking independence:

$$\frac{n}{2^n} \stackrel{?}{=} \frac{1+n}{2^n} \left( 1 - \frac{1}{2^{n-1}} \right)$$



# Families, reprise

- \* A family is known to have  $n$  children. Make the appropriate assumptions for the probability space.

- \* *Sample space*  $\Omega$ : all  $n$ -tuples of  $b$ 's and  $g$ 's.

- \* *Events*:

- \*  $A :=$  The family has at most one girl.

- \*  $B :=$  The family has children of both sexes;  $B^c :=$  the family has only boys or only girls.

- \* *Probability measure*: Combinatorial setting with mass function assigning equal probability  $1/2^n$  to each atom.

- \* Are  $A$  and  $B$  independent events?

#outcomes with no girls ←      → #outcomes with one girl

$$\mathbf{P}(A) = \frac{1+n}{2^n}$$

$$\mathbf{P}(B^c) = \frac{2}{2^n} = \frac{1}{2^{n-1}} \quad \mathbf{P}(B) = 1 - \mathbf{P}(B^c) = 1 - \frac{1}{2^{n-1}}$$

$$\mathbf{P}(A \cap B) = \frac{n}{2^n}$$

Checking independence:

$$\frac{n}{2^n} \stackrel{?}{=} \frac{1+n}{2^n} \left(1 - \frac{1}{2^{n-1}}\right)$$

$$n \stackrel{?}{=} 1+n - \frac{1+n}{2^{n-1}}$$



# Families, reprise

- \* A family is known to have  $n$  children. Make the appropriate assumptions for the probability space.

- \* *Sample space*  $\Omega$ : all  $n$ -tuples of  $b$ 's and  $g$ 's.

- \* *Events*:

- \*  $A :=$  The family has at most one girl.
- \*  $B :=$  The family has children of both sexes;  $B^c :=$  the family has only boys or only girls.

- \* *Probability measure*: Combinatorial setting with mass function assigning equal probability  $1/2^n$  to each atom.

- \* Are  $A$  and  $B$  independent events?

#outcomes with no girls ←      → #outcomes with one girl

$$\mathbf{P}(A) = \frac{1+n}{2^n}$$

$$\mathbf{P}(B^c) = \frac{2}{2^n} = \frac{1}{2^{n-1}} \quad \mathbf{P}(B) = 1 - \mathbf{P}(B^c) = 1 - \frac{1}{2^{n-1}}$$

$$\mathbf{P}(A \cap B) = \frac{n}{2^n}$$

Checking independence:

$$\frac{n}{2^n} \stackrel{?}{=} \frac{1+n}{2^n} \left(1 - \frac{1}{2^{n-1}}\right)$$

$$n \stackrel{?}{=} 1+n - \frac{1+n}{2^{n-1}}$$

$$2^{n-1} \stackrel{?}{=} 1+n$$



# Families, reprise

- \* A family is known to have  $n$  children. Make the appropriate assumptions for the probability space.
  - \* *Sample space*  $\Omega$ : all  $n$ -tuples of  $b$ 's and  $g$ 's.
  - \* *Events*:
    - \*  $A :=$  The family has at most one girl.
    - \*  $B :=$  The family has children of both sexes;  $B^c :=$  the family has only boys or only girls.
  - \* *Probability measure*: Combinatorial setting with mass function assigning equal probability  $1/2^n$  to each atom.
- \* Are  $A$  and  $B$  independent events?

#outcomes with no girls

#outcomes with one girl

$$P(A) = \frac{1 + n}{2^n}$$

$$P(B^c) = \frac{2}{2^n} = \frac{1}{2^{n-1}}$$

$$P(B) = 1 - P(B^c) = 1 - \frac{1}{2^{n-1}}$$

$$P(A \cap B) = \frac{n}{2^n}$$

n	1 + n	2 <sup>n-1</sup>
2	3	2
3	4	4
4	5	8
5	6	16
6	7	32

Checking independence:

$$\frac{n}{2^n} \stackrel{?}{=} \frac{1 + n}{2^n} \left( 1 - \frac{1}{2^{n-1}} \right)$$

$$n \stackrel{?}{=} 1 + n - \frac{1 + n}{2^{n-1}}$$

$$2^{n-1} \stackrel{?}{=} 1 + n$$



# Families, reprise

- \* A family is known to have  $n$  children. Make the appropriate assumptions for the probability space.
- \* *Sample space*  $\Omega$ : all  $n$ -tuples of  $b$ 's and  $g$ 's.
- \* *Events*:
  - \*  $A :=$  The family has at most one girl.
  - \*  $B :=$  The family has children of both sexes;  $B^c :=$  the family has only boys or only girls.
- \* *Probability measure*: Combinatorial setting with mass function assigning equal probability  $1/2^n$  to each atom.
- \* Are  $A$  and  $B$  independent events?

$A$  and  $B$  are independent if, and only if,  $n = 3$ .

#outcomes with no girls ←      → #outcomes with one girl

$$P(A) = \frac{1 + n}{2^n}$$

$$P(B^c) = \frac{2}{2^n} = \frac{1}{2^{n-1}} \qquad P(B) = 1 - P(B^c) = 1 - \frac{1}{2^{n-1}}$$

$$P(A \cap B) = \frac{n}{2^n}$$

$n$	$1 + n$	$2^{n-1}$
2	3	2
3	4	4
4	5	8
5	6	16
6	7	32

Checking independence:

$$\frac{n}{2^n} \stackrel{?}{=} \frac{1 + n}{2^n} \left( 1 - \frac{1}{2^{n-1}} \right)$$

$$n \stackrel{?}{=} 1 + n - \frac{1 + n}{2^{n-1}}$$

$$2^{n-1} \stackrel{?}{=} 1 + n$$