

# F-distribution

From Wikipedia, the free encyclopedia

In probability theory and statistics, the ***F*-distribution** is a continuous probability distribution.<sup>[1][2][3][4]</sup> It is also known as **Snedecor's *F* distribution** or the **Fisher–Snedecor distribution** (after R. A. Fisher and George W. Snedecor). The *F*-distribution arises frequently as the null distribution of a test statistic, most notably in the analysis of variance; see *F*-test.


## Contents

- 1 Definition
- 2 Characterization
- 3 Generalization
- 4 Related distributions and properties
- 5 See also
- 6 References
- 7 External links

## Definition

If a random variable *X* has an *F*-distribution with parameters *d*<sub>1</sub> and *d*<sub>2</sub>, we write *X* ~ F(*d*<sub>1</sub>, *d*<sub>2</sub>). Then the probability density function for *X* is given by

### Fisher-Snedecor



<b>Variance</b>	$\frac{2 d_2^2 (d_1 + d_2 - 2)}{d_1 (d_2 - 2)^2 (d_2 - 4)}$ for $d_2 > 4$
<b>Skewness</b>	$\frac{(2d_1 + d_2 - 2) \sqrt{8(d_2 - 4)}}{(d_2 - 6) \sqrt{d_1 (d_1 + d_2 - 2)}}$ for $d_2 > 6$
<b>Ex. kurtosis</b>	<i>see text</i>
<b>MGF</b>	<i>does not exist, raw moments defined in text and in <sup>[1]</sup>[2]</i>
<b>CF</b>	<i>see text</i>

$$f(x; d_1, d_2) = \frac{\sqrt{\frac{(d_1 x)^{d_1} d_2^{d_2}}{(d_1 x + d_2)^{d_1 + d_2}}}}{x B\left(\frac{d_1}{2}, \frac{d_2}{2}\right)} = \frac{1}{B\left(\frac{d_1}{2}, \frac{d_2}{2}\right)} \left(\frac{d_1}{d_2}\right)^{\frac{d_1}{2}} x^{\frac{d_1}{2}-1} \left(1 + \frac{d_1}{d_2} x\right)^{-\frac{d_1+d_2}{2}}$$

for real  $x \geq 0$ . Here  $B$  is the beta function. In many applications, the parameters  $d_1$  and  $d_2$  are positive integers, but the distribution is well-defined for positive real values of these parameters.

The cumulative distribution function is

$$F(x; d_1, d_2) = I_{\frac{d_1 x}{d_1 x + d_2}}\left(\frac{d_1}{2}, \frac{d_2}{2}\right),$$

where  $I$  is the regularized incomplete beta function.

The expectation, variance, and other details about the  $F(d_1, d_2)$  are given in the sidebar; for  $d_2 > 8$ , the excess kurtosis is

$$\gamma_2 = 12 \frac{d_1(5d_2 - 22)(d_1 + d_2 - 2) + (d_2 - 4)(d_2 - 2)^2}{d_1(d_2 - 6)(d_2 - 8)(d_1 + d_2 - 2)}.$$

The  $k$ -th moment of an  $F(d_1, d_2)$  distribution exists and is finite only when  $2k < d_2$  and it is equal to <sup>[5]</sup>

$$\mu_X(k) = \left(\frac{d_2}{d_1}\right)^k \frac{\Gamma\left(\frac{d_1}{2} + k\right) \Gamma\left(\frac{d_2}{2} - k\right)}{\Gamma\left(\frac{d_1}{2}\right) \Gamma\left(\frac{d_2}{2}\right)}$$

The  $F$ -distribution is a particular parametrization of the beta prime distribution, which is also called the beta distribution of the second kind.

The characteristic function is listed incorrectly in many standard references (e.g., <sup>[2]</sup>). The correct expression <sup>[6]</sup> is

$$\varphi_{d_1, d_2}^F(s) = \frac{\Gamma(\frac{d_1+d_2}{2})}{\Gamma(\frac{d_2}{2})} U\left(\frac{d_1}{2}, 1 - \frac{d_2}{2}, -\frac{d_2}{d_1} s\right)$$

where  $U(a, b, z)$  is the confluent hypergeometric function of the second kind.

## Characterization

A random variate of the  $F$ -distribution with parameters  $d_1$  and  $d_2$  arises as the ratio of two appropriately scaled chi-squared variates:<sup>[7]</sup>

$$X = \frac{U_1/d_1}{U_2/d_2}$$

where

- $U_1$  and  $U_2$  have chi-squared distributions with  $d_1$  and  $d_2$  degrees of freedom respectively, and
- $U_1$  and  $U_2$  are independent.

In instances where the  $F$ -distribution is used, for example in the analysis of variance, independence of  $U_1$  and  $U_2$  might be demonstrated by applying Cochran's theorem.

Equivalently, the random variable of the  $F$ -distribution may also be written

$$X = \frac{s_1^2}{\sigma_1^2} / \frac{s_2^2}{\sigma_2^2}$$

where  $s_1^2$  and  $s_2^2$  are the sums of squares  $S_1^2$  and  $S_2^2$  from two normal processes with variances  $\sigma_1^2$  and  $\sigma_2^2$  divided by the corresponding number of  $\chi^2$  degrees of freedom,  $d_1$  and  $d_2$  respectively.<sup>[citation needed]</sup>

In a frequentist context, a scaled  $F$ -distribution therefore gives the probability  $p(s_1^2/s_2^2 \mid \sigma_1^2, \sigma_2^2)$ , with the  $F$ -distribution itself, without any scaling, applying where  $\sigma_1^2$  is being taken equal to  $\sigma_2^2$ . This is the context in which the  $F$ -distribution most generally appears in  $F$ -tests: where the null hypothesis is that two independent normal variances are equal, and the observed sums of some appropriately selected squares are then examined to see whether their ratio is significantly incompatible with this null hypothesis.

The quantity  $X$  has the same distribution in Bayesian statistics, if an uninformative rescaling-invariant Jeffreys prior is taken for the prior probabilities of  $\sigma_1^2$  and  $\sigma_2^2$ .<sup>[8]</sup> In this context, a scaled  $F$ -distribution thus gives the posterior probability  $p(\sigma_2^2/\sigma_1^2 \mid s_1^2, s_2^2)$ , where now the observed sums  $s_1^2$  and  $s_2^2$  are what are taken as known.

## Generalization

A generalization of the (central)  $F$ -distribution is the noncentral  $F$ -distribution.

## Related distributions and properties

- If  $X \sim \chi_{d_1}^2$  and  $Y \sim \chi_{d_2}^2$  are independent, then  $\frac{X/d_1}{Y/d_2} \sim F(d_1, d_2)$
- If  $X \sim \text{Beta}(d_1/2, d_2/2)$  (Beta distribution) then  $\frac{d_2 X}{d_1(1-X)} \sim F(d_1, d_2)$
- Equivalently, if  $X \sim F(d_1, d_2)$ , then  $\frac{d_1 X/d_2}{1 + d_1 X/d_2} \sim \text{Beta}(d_1/2, d_2/2)$ .
- If  $X \sim F(d_1, d_2)$  then  $Y = \lim_{d_2 \rightarrow \infty} d_1 X$  has the chi-squared distribution  $\chi_{d_1}^2$
- $F(d_1, d_2)$  is equivalent to the scaled Hotelling's T-squared distribution  $\frac{d_2}{d_1(d_1 + d_2 - 1)} T^2(d_1, d_1 + d_2 - 1)$ .
- If  $X \sim F(d_1, d_2)$  then  $X^{-1} \sim F(d_2, d_1)$ .
- If  $X \sim t(n)$  then

$$\begin{aligned} X^2 &\sim F(1, n) \\ X^{-2} &\sim F(n, 1) \end{aligned}$$

- $F$ -distribution is a special case of type 6 Pearson distribution
- If  $X$  and  $Y$  are independent, with  $X, Y \sim \text{Laplace}(\mu, b)$  then

$$\frac{|X-\mu|}{|Y-\mu|} \sim F(2, 2)$$

- If  $X \sim F(n, m)$  then  $\frac{\log X}{2} \sim \text{FisherZ}(n, m)$  (Fisher's z-distribution)
- The noncentral  $F$ -distribution simplifies to the  $F$ -distribution if  $\lambda = 0$ .
- The doubly noncentral  $F$ -distribution simplifies to the  $F$ -distribution if  $\lambda_1 = \lambda_2 = 0$
- If  $Q_X(p)$  is the quantile  $p$  for  $X \sim F(d_1, d_2)$  and  $Q_Y(1-p)$  is the quantile  $1-p$  for  $Y \sim F(d_2, d_1)$ , then

$$Q_X(p) = \frac{1}{Q_Y(1-p)}.$$

## See also

- Chi-squared distribution
- Chow test
- Gamma distribution
- Hotelling's T-squared distribution
- Student's t-distribution
- Wilks' lambda distribution
- Wishart distribution

## References

- <sup>a</sup> <sup>b</sup> Johnson, Norman Lloyd; Samuel Kotz, N. Balakrishnan (1995). *Continuous Univariate Distributions*,

*Volume 2 (Second Edition, Section 27)*. Wiley. ISBN 0-471-58494-0.

2. ^ *a b c* Abramowitz, Milton; Stegun, Irene A., eds. (1965), "Chapter 26" ([http://www.math.sfu.ca/~cbm/aands/page\\_946.htm](http://www.math.sfu.ca/~cbm/aands/page_946.htm)), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, New York: Dover, p. 946, ISBN 978-0486612720, MR 0167642 (<http://www.ams.org/mathscinet-getitem?mr=0167642>).
3. ^ NIST (2006). Engineering Statistics Handbook – F Distribution (<http://www.itl.nist.gov/div898/handbook/eda/section3/eda3665.htm>)
4. ^ Mood, Alexander; Franklin A. Graybill, Duane C. Boes (1974). *Introduction to the Theory of Statistics (Third Edition, pp. 246–249)*. McGraw-Hill. ISBN 0-07-042864-6.
5. ^ Taboga, Marco. "The F distribution" ([http://www.statlect.com/F\\_distribution.htm](http://www.statlect.com/F_distribution.htm)).
6. ^ Phillips, P. C. B. (1982) "The true characteristic function of the F distribution," *Biometrika*, 69: 261–264 JSTOR 2335882 (<http://www.jstor.org/stable/2335882>)
7. ^ M.H. DeGroot (1986), *Probability and Statistics* (2nd Ed), Addison-Wesley. ISBN 0-201-11366-X, p. 500
8. ^ G.E.P. Box and G.C. Tiao (1973), *Bayesian Inference in Statistical Analysis*, Addison-Wesley. p.110

## External links

- Table of critical values of the *F*-distribution (<http://www.itl.nist.gov/div898/handbook/eda/section3/eda3673.htm>)
- Earliest Uses of Some of the Words of Mathematics: entry on *F*-distribution contains a brief history (<http://jeff560.tripod.com/f.html>)
- Free calculator for *F*-testing (<http://www.waterlog.info/f-test.htm>)

Retrieved from "<http://en.wikipedia.org/w/index.php?title=F-distribution&oldid=599359656>"

Categories: Continuous distributions | Analysis of variance | Probability distributions

- 
- This page was last modified on 12 March 2014 at 23:26.
  - Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy.
- Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.