



An R Introduction to Statistics

[HOME](#)[DOWNLOAD](#)[SALES](#)[EBOOK](#)[SITE MAP](#)

Two-Tailed Test of Population Proportion

The null hypothesis of the **two-tailed test about population proportion** can be expressed as follows:

$$p = p_0$$

where p_0 is a hypothesized value of the true population proportion p .

Let us define the test statistic z in terms of the sample proportion and the sample size:

$$z = \frac{\bar{p} - p_0}{\sqrt{p_0(1 - p_0)/n}}$$

Then the null hypothesis of the two-tailed test is to be *rejected* if $z \leq -z_{\alpha/2}$ or $z \geq z_{\alpha/2}$, where $z_{\alpha/2}$ is the $100(1 - \alpha)$ percentile of the **standard normal distribution**.

Problem

Suppose a coin toss turns up 12 heads out of 20 trials. At .05 significance level, can one reject the null hypothesis that the coin toss is fair?

Solution

The null hypothesis is that $p = 0.5$. We begin with computing the test statistic.

```
> pbar = 12/20          # sample proportion
> p0 = .5               # hypothesized value
> n = 20                # sample size
> z = (pbar-p0)/sqrt(p0*(1-p0)/n)
> z                     # test statistic
[1] 0.89443
```

We then compute the critical values at .05 significance level.

```
> alpha = .05
> z.half.alpha = qnorm(1-alpha/2)
> c(-z.half.alpha, z.half.alpha)
[1] -1.9600 1.9600
```

Answer

The test statistic 0.89443 lies between the critical values -1.9600 and 1.9600. Hence, at .05 significance level, we do *not* reject the null hypothesis that the coin toss is fair.

Alternative Solution 1

Instead of using the critical value, we apply the `pnorm` function to compute the two-tailed **p-value** of the test statistic. It doubles the *upper* tail p-value as the sample proportion is *greater* than the hypothesized value. Since it turns out to be greater than the .05 significance level, we do not reject the null hypothesis that $p = 0.5$.

```
> pval = 2 * pnorm(z, lower.tail=FALSE) # upper tail
> pval                                  # two-tailed p-value
[1] 0.37109
```

Alternative Solution 2

Search this site:

Search

R Tutorial eBook



R Tutorials

[R Introduction](#)[Elementary Statistics with R](#)[Qualitative Data](#)[Quantitative Data](#)[Numerical Measures](#)[Probability Distributions](#)[Interval Estimation](#)[Hypothesis Testing](#)[Lower Tail Test of
Population Mean with
Known Variance](#)[Upper Tail Test of
Population Mean with
Known Variance](#)[Two-Tailed Test of
Population Mean with
Known Variance](#)[Lower Tail Test of
Population Mean with
Unknown Variance](#)[Upper Tail Test of
Population Mean with
Unknown Variance](#)[Two-Tailed Test of
Population Mean with
Unknown Variance](#)[Lower Tail Test of](#)

We apply the `prop.test` function to compute the p-value directly. The Yates continuity correction is disabled for pedagogical reasons.

```
> prop.test(12, 20, p=0.5, correct=FALSE)

      1-sample proportions test without continuity
      correction

data:  12 out of 20, null probability 0.5
X-squared = 0.8, df = 1, p-value = 0.3711
alternative hypothesis: true p is not equal to 0.5
95 percent confidence interval:
 0.38658 0.78119
sample estimates:
      p 
 0.6
```

< Upper Tail Test of Population Proportion

up

Type II Error >

Tags: [Elementary Statistics with R](#) [hypothesis testing](#) [normal distribution](#) [p-value](#) [population proportion](#) [pnorm](#) [prop.test](#) [qnorm](#) [sqrt](#)

Population Proportion
Upper Tail Test of Population Proportion
Two-Tailed Test of Population Proportion
Type II Error
Inference About Two Populations
Goodness of Fit
Analysis of Variance
Non-parametric Methods
Simple Linear Regression
Multiple Linear Regression
Logistic Regression
GPU Computing with R

Recent Articles

- [Installing CUDA Toolkit 7.5 on Fedora 21 Linux](#)
September 10, 2015
- [Installing CUDA Toolkit 7.5 on Ubuntu 14.04 Linux](#)
September 10, 2015
- [Hierarchical Linear Model](#)
July 22, 2013
- [Bayesian Classification with Gaussian Process](#)
January 6, 2013