

What lessons can we deduce?

What lessons can we deduce?

When the sample space is **discrete**:

$$\Omega = \{ \omega_k : k \geq 1 \} \quad \text{or} \quad \{ \omega_k : k = 0, \pm 1, \pm 2, \dots \}$$

What lessons can we deduce?

When the sample space is **discrete**:

$$\Omega = \{ \omega_k : k \geq 1 \} \quad \text{or} \quad \{ \omega_k : k = 0, \pm 1, \pm 2, \dots \}$$

In many discrete applications it will be convenient to rename the sample points and identify them with integers: $\omega_k = k$, $\Omega = \mathbb{Z}$.

What lessons can we deduce?

When the sample space is **discrete**:

$$\Omega = \{ \omega_k : k \geq 1 \} \quad \text{or} \quad \{ \omega_k : k = 0, \pm 1, \pm 2, \dots \}$$

In many discrete applications it will be convenient to rename the sample points and identify them with integers: $\omega_k = k$, $\Omega = \mathbb{Z}$.

- **Atoms** (singleton sets consisting of single sample points) carry probabilistic mass called the **atomic probabilities**:

$$\mathbf{P}\{\omega_k\} := p(k) \geq 0, \quad \sum_k p(k) = 1$$

What lessons can we deduce?

When the sample space is **discrete**:

$$\Omega = \{ \omega_k : k \geq 1 \} \quad \text{or} \quad \{ \omega_k : k = 0, \pm 1, \pm 2, \dots \}$$

In many discrete applications it will be convenient to rename the sample points and identify them with integers: $\omega_k = k$, $\Omega = \mathbb{Z}$.

- **Atoms** (singleton sets consisting of single sample points) carry probabilistic mass called the **atomic probabilities**:

$$\overset{\text{Positivity}}{\mathbf{P}\{\omega_k\} := p(k) \geq 0,} \quad \overset{\text{Normalisation}}{\sum_k p(k) = 1}$$

What lessons can we deduce?

When the sample space is **discrete**:

$$\Omega = \{ \omega_k : k \geq 1 \} \quad \text{or} \quad \{ \omega_k : k = 0, \pm 1, \pm 2, \dots \}$$

In many discrete applications it will be convenient to rename the sample points and identify them with integers: $\omega_k = k$, $\Omega = \mathbb{Z}$.

- **Atoms** (singleton sets consisting of single sample points) carry probabilistic mass called the **atomic probabilities**:

$$\overset{\text{Positivity}}{\mathbf{P}\{\omega_k\} := p(k) \geq 0,} \quad \overset{\text{Normalisation}}{\sum_k p(k) = 1}$$

- Event probabilities are obtained by summing over atomic probabilities for the constituent atoms:

$$A = \{\omega_{k_1}, \omega_{k_2}, \dots\} \subseteq \Omega$$
$$\mathbf{P}(A) = \sum_{\omega \in A} \mathbf{P}\{\omega\} = p(k_1) + p(k_2) + \dots$$

What lessons can we deduce?

When the sample space is **discrete**:

$$\Omega = \{ \omega_k : k \geq 1 \} \quad \text{or} \quad \{ \omega_k : k = 0, \pm 1, \pm 2, \dots \}$$

In many discrete applications it will be convenient to rename the sample points and identify them with integers: $\omega_k = k$, $\Omega = \mathbb{Z}$.

- **Atoms** (singleton sets consisting of single sample points) carry probabilistic mass called the **atomic probabilities**:

$$\overset{\text{Positivity}}{\mathbf{P}\{\omega_k\} := p(k) \geq 0,} \quad \overset{\text{Normalisation}}{\sum_k p(k) = 1}$$

- Event probabilities are obtained by summing over atomic probabilities for the constituent atoms:

$$A = \{ \omega_{k_1}, \omega_{k_2}, \dots \} \subseteq \Omega$$
$$\mathbf{P}(A) = \sum_{\omega \in A} \mathbf{P}\{\omega\} = p(k_1) + p(k_2) + \dots$$

What lessons can we deduce?

When the sample space is **discrete**:

$$\Omega = \{ \omega_k : k \geq 1 \} \quad \text{or} \quad \{ \omega_k : k = 0, \pm 1, \pm 2, \dots \}$$

In many discrete applications it will be convenient to rename the sample points and identify them with integers: $\omega_k = k$, $\Omega = \mathbb{Z}$.

- **Atoms** (singleton sets consisting of single sample points) carry probabilistic mass called the **atomic probabilities**:

$$\overset{\text{Positivity}}{\mathbf{P}\{\omega_k\} := p(k) \geq 0,} \quad \overset{\text{Normalisation}}{\sum_k p(k) = 1}$$

- Event probabilities are obtained by summing over atomic probabilities for the constituent atoms:

$$A = \{ \omega_{k_1}, \omega_{k_2}, \dots \} \subseteq \Omega$$
$$\mathbf{P}(A) = \sum_{\omega \in A} \mathbf{P}\{\omega\} = p(k_1) + p(k_2) + \dots$$

What lessons can we deduce?

When the sample space is **discrete**:

$$\Omega = \{ \omega_k : k \geq 1 \} \quad \text{or} \quad \{ \omega_k : k = 0, \pm 1, \pm 2, \dots \}$$

In many discrete applications it will be convenient to rename the sample points and identify them with integers: $\omega_k = k$, $\Omega = \mathbb{Z}$.

- **Atoms** (singleton sets consisting of single sample points) carry probabilistic mass called the **atomic probabilities**:

Positivity

$$\mathbf{P}\{\omega_k\} := p(k) \geq 0,$$

Normalisation

$$\sum_k p(k) = 1$$

- Event probabilities are obtained by summing over atomic probabilities for the constituent atoms:

$$A = \{ \omega_{k_1}, \omega_{k_2}, \dots \} \subseteq \Omega$$

$$\mathbf{P}(A) = \sum_{\omega \in A} \mathbf{P}\{\omega\} = p(k_1) + p(k_2) + \dots$$

Additivity

Characterising chance in discrete spaces

Characterising chance in discrete spaces

- ❖ **Atoms** (singleton events) $\{\omega_k\}$ carry mass in discrete spaces.

Characterising chance in discrete spaces

- ❖ **Atoms** (singleton events) $\{\omega_k\}$ carry mass in discrete spaces.
- ❖ The atomic probabilities $p(k) := \mathbf{P}\{\omega_k\}$ determine a function called the **mass function** or **distribution**. This function satisfies *positivity*, $p(k) \geq 0$, and is properly *normalised*, $\sum_k p(k) = 1$.

Characterising chance in discrete spaces

- ❖ **Atoms** (singleton events) $\{\omega_k\}$ carry mass in discrete spaces.
- ❖ The atomic probabilities $p(k) := \mathbf{P}\{\omega_k\}$ determine a function called the **mass function** or **distribution**. This function satisfies *positivity*, $p(k) \geq 0$, and is properly *normalised*, $\sum_k p(k) = 1$.
- ❖ The probability measure $\mathbf{P}(\cdot)$ is determined from the distribution $p(\cdot)$ via *additivity*:

$$\mathbf{P}\{\omega_{k_1}, \omega_{k_2}, \omega_{k_3}, \dots\} = p(k_1) + p(k_2) + p(k_3) + \dots.$$

Any *honest* **mass function** $p(k)$ induces a discrete probability measure.
All we have to do is verify that it is *positive* and properly *normalised*.