

[< Previous](#)

 





[Next >](#)

12.5.2 The Rayleigh Quotient Iteration

 Bookmark this page

12.5.2 The Rayleigh Quotient Iteration

Reading Assignment

0 points possible (ungraded)
Read Unit 12.5.2 of the notes. [\[LINK\]](#)

☒ Done

✓

Submit

✓ Correct

Homework 12.5.2.1

1/1 point (ungraded)
The script in `RayleighQuotientIterationScript.m` illustrates how shifting the matrix by the Rayleigh Quotient can greatly improve how fast the Shifted Inverse Power Method, starting with a random vector, computes an eigenvector. It could be that the random vector is close to an eigenvector associated with any of the eigenvalues, in which case the method will start converging towards an eigenvector associated with that eigenvalue. Pay close attention to how many digit are accurate from one iteration to the next.

To try it out, in the Command Window type

>> RayleighQuotientIterationScript
input a vector of eigenvalues. e.g.: [4; 3; 2; 1]
[4; 3; 2; 1]

☒ done/skip

✓

Submit

i Answers are displayed within the problem



edX

- About
- Affiliates
- edX for Business
- Open edX
- Careers
- News

Legal

- Terms of Service & Honor Code
- Privacy Policy
- Accessibility Policy
- Trademark Policy
- Sitemap
- Cookie Policy
- Your Privacy Choices

Connect

- Idea Hub
- Contact Us
- Help Center
- Security
- Media Kit



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)