

<u>Help</u>

sandipan_dey >

Next >

Progress <u>Dates</u> <u>Calendar</u> **Discussion** <u>Notes</u> <u>Course</u>

☆ Course / Unit 3: Optimization / Lecture 11: Lagrange Multipliers



You are taking "Exam (Timed, No Correctness Feedback)" as a timed exam. Show more



22:49:28





☐ Bookmark this page

Previous



Explore

Optimization along the boundary



Start of transcript. Skip to the end.

PROFESSOR: Let's see what people are thinking.

So we know that the maximum is on the boundary,

so I'm just going to move the cursor-

move the little-- the cursor along the boundary.

And you guys will tell me, keep

0:00 / 0:00

2.0x

X

" CC

aoina or stop there

Video

Download video file

Transcripts

Download SubRip (.srt) file Download Text (.txt) file

We can start to think about optimization by first zooming in on the function $f\left(x,y
ight)$ and using the linear approximation. Say we start at (x,y) and move to a point $(x+\Delta x,y+\Delta y)$. We want to know if doing this causes the change in $m{f}$ to be positive, negative, or zero. The linear approximation is given by


$$f(x + \Delta x, y + \Delta y) \approx f(x, y) + \underbrace{f_x(x, y) \Delta x + f_y(x, y) \Delta y}_{\text{change in } f}$$
 (4.155)

where we have highlighted the change in f as the part of the linear approximation that deviates from f(x,y).

We can write the change in $m{f}$ as the following dot product

change in
$$f = f_x(x, y) \Delta x + f_y(x, y) \Delta y = \langle f_x, f_y \rangle \cdot \langle \Delta x, \Delta y \rangle = |\langle f_x, f_y \rangle| |\langle \Delta x, \Delta y \rangle| \cos \theta$$
 (4.156)

where heta is the angle between the gradient $\langle f_x, f_y
angle$ and $\langle \Delta x, \Delta y
angle$. An illustration of this is shown below.



From the equation for the change in $m{f}$, we can conclude the following:


• If $heta<rac{\pi}{2}$, then $\cos heta>0$ and therefore the change in f is positive.

⊞ Calculator

Hide Notes

- If $heta>rac{\pi}{2}$, then $\cos heta<0$ and therefore the change in f is negative.
- If $heta=rac{\pi}{2}$, then $\cos heta=0$ and therefore the change in f is zero.

What does this mean when we are looking at the change of a function $f\left(x,y
ight)$ as we move along some curve?



- ullet If you move in a direction along a curve that makes an angle less than $\pi/2$ with the gradient of a f, the function $oldsymbol{f}$ is increasing as you move along the curve in that direction.
- ullet If you move in a direction along a curve that makes an angle greater than $\pi/2$ with the gradient of f, the function \boldsymbol{f} decreasing as you move along the curve in that direction.
- ullet If you move in a direction along a curve that makes an angle equal to $\pi/2$ with the gradient of f, the function fis neither increasing or decreasing.

Therefore the function f is as its largest when we reach a point along the curve such that the curve and the gradient are perpendicular to each other. At this point, the function does not increase as you move in either direction. In all other cases, there is a direction you can move that will increase f.

Key point: The maximum of f along a boundary curve occurs when the gradient ∇f is normal (perpendicular) to the boundary.


The argument we gave was setup for a general function f(x,y) moving along a direction specified by a curve C. None of this argument was specific to our example, so this holds more generally.

4. Optimization and linear approximation

Topic: Unit 3: Optimization / 4. Optimization and linear approximation

Hide Discussion

Add a Post



© All Rights Reserved



edX

About

<u>Affiliates</u>

edX for Business

Open edX

Careers

<u>News</u>

Legal

Terms of Service & Honor Code

Privacy Policy

Accessibility Policy

Trademark Policy

<u>Sitemap</u>

Connect

Blog

Contact Us

Help Center

Media Kit

Donate



















© 2021 edX Inc. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>