

[Course](#)

[Progress](#)

[Dates](#)

[Discussion](#)

[Syllabus](#)

[Outline](#)

[laff routines](#)

[Community](#)

[🏠](#) [Course](#) / [Week 12: Eigenvalues and Eigenvectors](#) / [12.2 Getting Started](#)



< Previous	 	 	 	 		Next >
-------------------------------	---	---	---	---	---	---------------------------

12.2.2 Simple Examples

 Bookmark this page

Week 12 due Dec 29, 2023 10:42 IST Completed

12.2.2 Simple Examples

Video 12.2.2 Part 1

[Start of transcript. Skip to the end.](#)



▶ 0:00 / 0:00

▶ 2.0x

🔊

⌂

CC

“ ”

Dr. Robert van de Geijn: As we have often done in the past we will start by looking at some simple examples.

Go and do this homework.

In the homework what I want you to do is I want you to identify eigenvalues, λ , and corresponding eigenvectors

Video

📄 [Download video file](#)

Transcripts

📄 [Download SubRip \(.srt\) file](#)

📄 [Download Text \(.txt\) file](#)

Reading Assignment

0 points possible (ungraded)
Read Unit 12.2.2 of the notes. [\[LINK\]](#)

☒ Done

✓

Submit

Discussion

Hide Discussion

Topic: Week 12 / 12.2.2 Part 1

Add a Post

Show all posts ▼ by recent activity ▼

There are no posts in this topic yet.

✕

🧮 Calculator

Homework 12.2.2.1

10.0/10.0 points (graded)

Which of the following are eigenpairs (λ, x) of the 2×2 zero matrix:

$$\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} x = \lambda x,$$

where $x \neq 0$.

(Mark all correct answers.)

☐ $(1, \begin{pmatrix} 0 \\ 0 \end{pmatrix})$.

☒ $(0, \begin{pmatrix} 1 \\ 0 \end{pmatrix})$.

☒ $(0, \begin{pmatrix} 0 \\ 1 \end{pmatrix})$.

☒ $(0, \begin{pmatrix} -1 \\ 1 \end{pmatrix})$.

☒ $(0, \begin{pmatrix} 1 \\ 1 \end{pmatrix})$.

☐ $(0, \begin{pmatrix} 0 \\ 0 \end{pmatrix})$.



Answers 2., 3., 4., and 5. are all correct, since for all $Ax = \lambda x$.

Answers 1. and 6. are not correct because the zero vector is not considered an eigenvector.

Submit

i Answers are displayed within the problem

Video 12.2.2 Part 2



so
is you subtract lambda from the
diagonal elements.
And if you then ask the question, when
is this matrix singular?
What you conclude is that lambda is
equal to 0.
And then once you actually plug that in,
you end up with the matrix 0, 0, 0, 0.
And you ask yourself the question, for
what x is this true?
And you end up again with the span of
the unit basis vectors.
So let's move on to something a little bit
more difficult.
Why don't you contemplate wha

Calculator

eigenvalues and eigenvectors.

of this matrix might be?

You will see me in the next video.

Video

Download video file

Transcripts

Download SubRip (.srt) file

Download Text (.txt) file

Homework 12.2.2.2

1/1 point (graded)

Which of the following are eigenpairs (λ, x) of the 2×2 zero matrix:

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} x = \lambda x,$$

where $x \neq 0$.

(Mark all correct answers.)

☐ $(1, \begin{pmatrix} 0 \\ 0 \end{pmatrix})$.

☒ $(1, \begin{pmatrix} 1 \\ 0 \end{pmatrix})$.

☒ $(1, \begin{pmatrix} 0 \\ 1 \end{pmatrix})$.

☒ $(1, \begin{pmatrix} -1 \\ 1 \end{pmatrix})$.

☒ $(1, \begin{pmatrix} 1 \\ 1 \end{pmatrix})$.

☐ $(-1, \begin{pmatrix} 1 \\ -1 \end{pmatrix})$.



Answers 2., 3., 4., and 5. are all correct, since for all $Ax = \lambda x$.

Answer 1. is not correct because the zero vector is not considered an eigenvector. Answer 6. is not correct because it doesn't satisfy $Ax = \lambda x$.

Submit

Answers are displayed within the problem

Video 12.2.2 Part 3

Start of transcript. Skip to the end.

Calculator



Dr. Robert van de Geijn: OK.

So again, we may want to do this by examination.

We know that the identity times x is always equal to x .

And we think of that as the scale of 1 times the vector x .

And what you notice is, therefore, that 1 seems to be an eigenvalue of the



Video

Download video file

Transcripts

Download SubRip (.srt) file

Download Text (.txt) file

Homework 12.2.2.3

3/3 points (graded)

Let $A = \begin{pmatrix} 3 & 0 \\ 0 & -1 \end{pmatrix}$.

- $\begin{pmatrix} 3 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ so that $(3, \begin{pmatrix} 1 \\ 0 \end{pmatrix})$ is an eigenpair.

TRUE

Answer: TRUE

Just multiply it out.

- The set of all eigenvectors associated with eigenvalue **3** is characterized by (mark all that apply):

☒ All vectors $x \neq 0$ that satisfy $Ax = 3x$.

☒ All vectors $x \neq 0$ that satisfy $(A - 3I)x = 0$.

☒ All vectors $x \neq 0$ that satisfy $\begin{pmatrix} 0 & 0 \\ 0 & -4 \end{pmatrix} x = 0$.

☐ $\left\{ \begin{pmatrix} \chi_0 \\ 0 \end{pmatrix} \middle| \chi_0 \text{ is a scalar} \right\}$



True: this is the definition of an eigenvector associated with an eigenvalue.

True: this is an alternate condition.

True: $(A - 3I) = \begin{pmatrix} 0 & 0 \\ 0 & -4 \end{pmatrix}$ in this case.

False: the zero vector is in this set.

Calculator

$$\begin{pmatrix} 3 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = -1 \begin{pmatrix} 0 \\ 1 \end{pmatrix} \text{ so that } (-1, \begin{pmatrix} 0 \\ 1 \end{pmatrix}) \text{ is an eigenpair.}$$

TRUE

✔ Answer: TRUE

True: Just multiply it out.

Submit

Answers are displayed within the problem

Video 12.2.2 Part 4

[Start of transcript. Skip to the end.](#)



Dr. Robert van de Geijn: So this time it might actually be useful to just go ahead and subtract lambda times the identity from the matrix. And then you end up with this right here.

< Previous

Next >

© All Rights Reserved



edX

- About
- Affiliates
- edX for Business
- Open edX
- Careers
- News

Legal

- Terms of Service & Honor Code
- Privacy Policy
- Accessibility Policy
- Trademark Policy
- Sitemap

Calculator

[Cookie Policy](#)
[Your Privacy Choices](#)

Connect

[Idea Hub](#)
[Contact Us](#)
[Help Center](#)
[Security](#)
[Media Kit](#)



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)