



Course &gt; Section... &gt; 2.2 The... &gt; 2.2.2 E...

## 2.2.2 Exploratory Quiz: Computing Average Value

[Bookmark this page](#)

### Question 1: Think About It...

1/1 point (graded)

Let's start with the average value of the function  $M(t)$  describing marlin population. The average value of the function  $M$  over a time period length  $L$  is defined as:

$$\overline{M} = \frac{1}{L} \int_0^L M(t) dt.$$

- Take the differential equation for the rate of change of marlin,  $\frac{dM}{dt} = -cM + dSM$ , and solve for  $M$ . Then substitute this expression for  $M(t)$  into the formula for average value  $\overline{M} = \frac{1}{L} \int_0^L M(t) dt$ .
- Take the differential equation for the rate of change of sardines  $\frac{dS}{dt} = aS - bSM$  and solve for  $M$ . Then substitute this expression for  $M(t)$  into the formula for average value  $\overline{M} = \frac{1}{L} \int_0^L M(t) dt$ .

You now have two possible ways to think about  $\overline{M} = \frac{1}{L} \int_0^L M(t) dt$ . Which integral looks easier to evaluate? Why?

second one, since it contains the variable S only.



Thank you for your response.

#### Explanation

These computations are covered in the next video. The integral from part (b) looks easier because it involves only  $S$  and its derivative  $\frac{dS}{dt}$ . The integral from part (a) involves two variables:  $S$  and the derivative of  $M$ ,  $\frac{dM}{dt}$ .

Submit



**i** Answers are displayed within the problem

[Learn About Verified Certificates](#)

© All Rights Reserved



 English ▼

© 2012–2018 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open edX logos are registered trademarks or trademarks of edX Inc. | 粤ICP备17044299号-2

POWERED BY  
**OPEN**edX®

