



< Previous	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	Next >
------------	---	---	---	---	---	---	---	---	---	---	--------

6. Moving average model

Bookmark this page

Exercises due Nov 10, 2021 17:29 IST Completed

Moving average model

Start of transcript. Skip to the end.

Prof Jegelka: After exploring our first three statistical models, that is white noise, autoregressive models, and random walk models. We'll now look at two other important models, and these are moving average models, and ARMA models.

▶ 0:00 / 0:00

▶ 2.0x

🔊

🔍

CC

🗣️

Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)
[Download Text \(.txt\) file](#)

A time series $\{X_t\}_t$ is a **moving average process** of order q , denoted by **MA** (q), if it can be represented as a weighted moving average

$$\begin{aligned} X_t &= W_t + \theta_1 W_{t-1} + \theta_2 W_{t-2} + \cdots + \theta_q W_{t-q} \\ &= \sum_{h=0}^q \theta_h W_{t-h} \end{aligned}$$

of a white noise series $\{W_t\}_t$.

Marginal mean of moving average model

1/1 point (graded)

What is the marginal mean μ_X of a moving average time series?

0

✔ Answer: 0

Solution:


We have $\mu_X(t) = \mathbf{E}(\sum_{h=0}^q \theta_h W_{t-h}) = 0$.

Submit

You have used 1 of 3 attempts

📘 Answers are displayed within the problem

ACF of moving average model



From the definition of a moving average time series, we find that the autocovariance function is

$$\gamma_X(h) = \text{Cov}\left(\sum_{j=0}^q \theta_j W_{t-j}, \sum_{k=0}^q \theta_k W_{t+h-k}\right)$$
$$= \sum_{j=0}^{q-h} \theta_j \theta_{j+h} \sigma_W^2 \quad \text{for } 0 \leq h \leq q$$

because W_j is uncorrelated with W_k for $j \neq k$.

ACF of moving average model

5/5 points (graded)
Let $X_t = W_t + \frac{1}{2}W_{t-1} + \frac{1}{3}W_{t-2}$ where $W_t \sim \text{WN}(\sigma^2 = 1)$.

Compute the variance $\gamma_X(0)$ and autocovariance $\gamma_X(h)$ for the following gaps $h = 1, 2, 3, 4$.

(Enter an answer accurate to at least 2 decimal places).

Variance $\gamma_X(0) =$

49/36

✔ Answer: 1.361

$\gamma_X(1)$

2/3

✔ Answer: 0.667

$\gamma_X(2)$

1/3

✔ Answer: 0.333

$\gamma_X(3)$

0

✔ Answer: 0

$\gamma_X(4)$

0

✔ Answer: 0

Solution:

$$\begin{aligned}\gamma_X(0) &= \text{Var}\left(W_t + \frac{1}{2}W_{t-1} + \frac{1}{3}W_{t-2}\right) \\ &= \text{Var}(W_t) + \frac{1}{4}\text{Var}(W_{t-1}) + \frac{1}{9}\text{Var}(W_{t-2}) \\ &= 1 + \frac{1}{4} + \frac{1}{9} = \frac{49}{36}\end{aligned}$$

$$\begin{aligned}\gamma_X(1) &= \text{Cov}\left(W_t + \frac{1}{2}W_{t-1} + \frac{1}{3}W_{t-2}, W_{t-1} + \frac{1}{2}W_{t-2} + \frac{1}{3}W_{t-3}\right) \\ &= \frac{1}{2}\text{Cov}(W_{t-1}, W_{t-1}) + \frac{1}{3} \cdot \frac{1}{2}\text{Cov}(W_{t-2}, W_{t-2}) \\ &= \frac{1}{2} + \frac{1}{3} \cdot \frac{1}{2} = \frac{4}{6}\end{aligned}$$

$$\begin{aligned}\gamma_X(2) &= \text{Cov}\left(W_t + \frac{1}{2}W_{t-1} + \frac{1}{3}W_{t-2}, W_{t-2} + \frac{1}{2}W_{t-3} + \frac{1}{3}W_{t-4}\right) \\ &= \frac{1}{3}\text{Cov}(W_{t-2}, W_{t-2}) = \frac{1}{3}\end{aligned}$$

The autocovariance $\gamma_X(h) = 0$ for all $h > 2$. **Remark:** In general, $\gamma_X(h) = 0$ for all $h > q$ because there are no common white noise terms in such terms of an **MA** (q) time series.

Submit

You have used 1 of 3 attempts

ⓘ Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Module 4: Time Series:Time Series: Statistical Models / 6. Moving average model

Add a Post

Show all posts ▼by recent activity ▼

💬 Lecture's aren't very good.

I know the last question can be easily solved because it was covered in the lecture. I even referred to the video but after looking at t...

16 ▼

< Previous

Next >



edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)

Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Media Kit](#)
- [Donate](#)



© 2021 edX Inc. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)