



[Course](#) > [Unit 1: Fourier Series](#) > [Period 2L](#) > [2. Properties of Fourier Series \(of](#)
> [13. Complex Fourier series](#)

Audit Access Expires Jun 24, 2020

You lose all access to this course, including your progress, on Jun 24, 2020.

Upgrade by Jun 7, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

13. Complex Fourier series

Here's an idea: Euler's formula

$$e^{it} = \cos t + i \sin t$$

tells us that complex exponentials can be written as a sum of a sine and a cosine function. This suggests that we might be able to write a Fourier series

$$\frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nt + b_n \sin nt)$$

with real coefficients a_n and b_n as a series of complex exponentials $\sum_{n=-\infty}^{\infty} c_n e^{int}$, for some **complex** coefficients c_n . As it turns out, this is true, that is, we can always write a Fourier series in terms of complex exponentials. Since the two series turn out to be equal, we'll also call the series in terms of complex exponentials a Fourier series.

So let's walk through the process of converting a series of the form



$$\frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nt + b_n \sin nt)$$

to a series of the form $\sum_{n=-\infty}^{\infty} c_n e^{int}$.

Step 1: Rewriting the sum.

Using Euler's formula and the fact that $\sin t$ is an odd function and $\cos t$ is an even function, we notice that

$$\sin t = \frac{i}{2} (e^{-it} - e^{it})$$

and

$$\cos t = \frac{1}{2} (e^{it} + e^{-it}).$$

Then we see that given any Fourier series f , we can write

$$\begin{aligned} f(t) &= \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nt + b_n \sin nt) \\ &= \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(\frac{a_n}{2} (e^{int} + e^{-int}) + \frac{ib_n}{2} (e^{-int} - e^{int}) \right) \\ &= \frac{a_0}{2} + \frac{1}{2} \sum_{n=1}^{\infty} ((a_n - ib_n) e^{int} + (a_n + ib_n) e^{-int}). \end{aligned}$$

Step 2: Defining coefficients.

We can see that f can be written as a sum of complex exponentials. Let's write the coefficients of these exponentials nicely so that we can easily convert back and forth between the two forms. Define $c_0 := a_0/2$. For $n > 0$, define



$$c_n := \frac{a_n - ib_n}{2},$$

and

$$c_{-n} := \bar{c}_n = \frac{a_n + ib_n}{2}.$$

Then we can write f compactly as

$$f(t) = \sum_{n=-\infty}^{\infty} c_n e^{int}.$$

Step 3: Relating coefficients to inner products.

Remark about inner products:

Note that for real valued 2π -periodic functions f and g we define an inner product as

$$\langle f, g \rangle = \int_{-\pi}^{\pi} f \cdot g \, dt.$$

If f and g are complex valued functions, we must define the inner product as

$$\langle f, g \rangle = \int_{-\pi}^{\pi} f \cdot \bar{g} \, dt,$$

where \bar{g} is the complex conjugate of g .

Example 13.1



$$\langle e^{int}, e^{int} \rangle = \int_{-\pi}^{\pi} e^{int} \cdot e^{-int} dt = \int_{-\pi}^{\pi} dt = 2\pi.$$

If $m \neq n$,

$$\begin{aligned} \langle e^{int}, e^{imt} \rangle &= \int_{-\pi}^{\pi} e^{int} \cdot e^{-imt} dt \\ &= \int_{-\pi}^{\pi} e^{i(n-m)t} dt \\ &= \left. \frac{e^{i(n-m)t}}{i(n-m)} \right|_{-\pi}^{\pi} \\ &= \frac{e^{i(n-m)\pi} - e^{-i(n-m)\pi}}{i(n-m)} = \frac{2}{n-m} \sin((n-m)\pi) = 0. \end{aligned}$$

Therefore the e^{int} are orthogonal.

In particular, notice that for $n > 0$, we can compute c_n by the formula

$$c_n = \frac{\langle f, e^{int} \rangle}{\langle e^{int}, e^{int} \rangle}.$$

This definition is in agreement with the definition of $c_n = \frac{a_n - ib_n}{2}$. (You can check this!)

Check worked out

[Show](#)

Key properties of complex Fourier series

- Often integrals involving complex exponentials are actually much easier to compute than integrals involving sines and cosines.



- We can also make sense of Fourier series of complex-valued functions more easily in this setting.

Complex Fourier series concept check

2/2 points (graded)

Let $Sq(t)$ be the 2π -periodic square wave that is equal to 1 for $0 \leq t < \pi$ and equal to -1 for $-\pi \leq t < 0$. For $n \neq 0$, write a general formula for the n th coefficient of the complex Fourier coefficient c_n of $Sq(t)$.

Tip : Type a^b for a^b , type π for π , and i for $i = \sqrt{-1}$.

n even, $c_n =$

✓ Answer: 0

n odd, $c_n =$

✓ Answer: $2/(i*n*\pi)$

Solution:

There are multiple ways to approach this problem, one is to use the coefficients a_n and b_n (which we already have calculated) to write c_n , another (better) approach is to calculate c_n directly; this is the approach we'll take. We see that for $n \neq 0$,

$$\begin{aligned}
 c_n &= \frac{1}{2\pi} \int_{-\pi}^{\pi} Sq(t) e^{-int} dt \\
 &= \frac{1}{2\pi} \int_{-\pi}^0 -e^{-int} dt + \frac{1}{2\pi} \int_0^{\pi} e^{-int} dt \\
 &= -\frac{1}{2\pi} \left(-\frac{1}{in} \right) e^{-int} \Big|_{-\pi}^0 + \frac{1}{2\pi} \left(-\frac{1}{in} \right) e^{-int} \Big|_0^{\pi} \\
 &= \frac{1}{2\pi in} (e^0 - e^{in\pi} - (e^{-in\pi} - e^0)) \\
 &= \frac{1}{2\pi in} (2 - (e^{in\pi} + e^{-in\pi})).
 \end{aligned}$$



Examining the cases where n is even and where n is odd separately, we notice that

$$c_n = \begin{cases} 0, & n \text{ even} \\ \frac{2}{in\pi}, & n \text{ odd} \end{cases}$$

$$= \frac{1 - (-1)^n}{in\pi}.$$

In the case that $n = 0$, we see that

$$c_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} S_q(t) dt = 0.$$

Hence the Fourier series for $S_q(t)$ is

$$S_q(t) = \sum_{n=-\infty}^{\infty} \frac{1 - (-1)^n}{in\pi} e^{int}.$$

Submit

You have used 1 of 5 attempts

i Answers are displayed within the problem

13. Complex Fourier series

Topic: Unit 1: Fourier Series / 13. Complex Fourier series

Hide Discussion

Add a Post

Show all posts ▼

by recent activity ▼

? [Confused about inner products with complex functions](#)

5

? [i m lost at step 2....
i don't seem to understand why the coefficients could be compacted?](#)

9



✓ [Step 3](#)

4

💬 [Different approach - same result](#)

2

[I did something different and I don't know why it happens to be correct. I just took F.S of the \$S_q\(t\)\$ and transfer it into complex F.S where I transfered: \$4/\(n\pi\) \sin n\pi t \rightarrow 2/\(n\pi\)\$](#)

💬 [Inner product when \$n \neq m\$](#)

3

✓ [Confusion regarding the question](#)

3

[The question asks for the \$n\$ th complex coefficients, where \$n\$ is either odd or even. My confusion is that since the coefficient depends on whether \$n\$ is positive or negative, sho...](#)

[Learn About Verified Certificates](#)

© All Rights Reserved

