

[Unit 2: Boundary value problems](#)

[Course](#) > [and PDEs](#)

4. Separation of variables: solving a  
PDE with homogeneous boundary

> [5. The Heat Equation](#) > conditions

## 4. Separation of variables: solving a PDE with homogeneous boundary conditions

### Worked example: homogeneous boundary conditions

both the initial condition and the two boundary conditions.

Handwritten notes on the chalkboard:

- Left section:**

$$\frac{\partial \theta}{\partial t} = \nu \frac{\partial^2 \theta}{\partial x^2}$$

$$\theta(x, 0) = 1$$

$$\theta(0, t) = 0$$

$$\theta(L, t) = 0$$

$$\theta(x, t) = a(x)b(t)$$

$$a''(x) = -\lambda a(x)$$

$$b'(t) = -\lambda b(t)$$
- Middle section:**

$$\frac{1}{b} \frac{db}{dt} = \frac{1}{a} \frac{d^2 a}{dx^2} = -\lambda$$

$$\frac{d^2 a}{dx^2} = -\lambda a$$

$$a(0) = 0$$

$$a(L) = 0$$

$$a(x) = A \sin(n\pi x)$$
- Right section:**

$$\frac{db}{dt} = -\lambda b$$

$$b(t) = B e^{-\lambda t}$$

$$\theta(x, t) = C_n e^{-\lambda t} \sin(n\pi x)$$

$$\theta(x, t) = \sum_{n=1}^{\infty} C_n e^{-\lambda t} \sin(n\pi x)$$

▶ 14:07 / 14:07

▶ 1.50x 🔊 🗖 📄 🗨

## Video

[Download video file](#)

## Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Let's now try to solve the PDE. For simplicity, suppose that  $L = \pi$ ,  $\theta_0 = 1$ , and  $\nu = 1$ . (The general case is similar. In fact, one could reduce to this special case by changes of variable.)



So now we are solving

$$\begin{aligned}\frac{\partial \theta}{\partial t} &= \frac{\partial^2 \theta}{\partial x^2}, & 0 < x < \pi, \ t > 0, \\ \theta(0, t) &= 0 & \text{for } t \geq 0 \\ \theta(\pi, t) &= 0 & \text{for } t \geq 0 \\ \theta(x, 0) &= 1 & \text{for } x \in (0, \pi).\end{aligned}$$

**Idea (separation of variables):** Forget about the initial condition  $\theta(x, 0) = 1$  for now, but look for nonzero solutions of the form

$$\theta(x, t) = v(x) w(t).$$

(Note that we are making a slight change in notation.) Substituting into the PDE gives

$$\begin{aligned}v(x) \dot{w}(t) &= w(t) v''(x) \\ \frac{\dot{w}(t)}{w(t)} &= \frac{v''(x)}{v(x)}.\end{aligned}$$

(at least where  $v(x)$  and  $w(t)$  are nonzero).

The only way for a function of  $x$  to equal to a function of  $t$  is if both functions are the same constant. That is, there is a constant  $\lambda$  such that

$$\frac{v''(x)}{v(x)} = \lambda \quad \text{and} \quad \frac{\dot{w}(t)}{w(t)} = \lambda,$$

or in other words,



$$v''(x) = \lambda v(x) \quad \text{and} \quad \dot{w}(t) = \lambda w(t).$$

Substituting  $\theta(x, t) = v(x) w(t)$  into the first boundary condition  $\theta(0, t) = 0$  gives  $v(0) w(t) = 0$  for all  $t$ , but  $w(t)$  is not the zero function, so this translates into  $v(0) = 0$ . Similarly, the second boundary condition  $\theta(\pi, t) = 0$  translates into  $v(\pi) = 0$ .

We already solved  $v''(x) = \lambda v(x)$  subject to the boundary conditions  $v(0) = 0$  and  $v(\pi) = 0$ : nonzero solutions  $v(x)$  exist only if  $\lambda = -n^2$  for some positive integer  $n$ , and in that case  $v(x)$  is a scalar times  $\sin nx$ .

For  $\lambda = -n^2$ , what is a matching possibility for  $w$ ? Since  $\dot{w} = -n^2 w$ , the function  $w$  is a scalar times  $e^{-n^2 t}$ .

This gives rise to one solution

$$\theta_n(x, t) = e^{-n^2 t} \sin nx$$

for each positive integer  $n$ , to the PDE with boundary conditions. Each such solution is called a **normal mode**.

Because the boundary conditions are homogeneous, we can get other solutions by taking linear combinations that also satisfy the homogeneous boundary conditions:

$$\theta(x, t) = b_1 e^{-t} \sin x + b_2 e^{-4t} \sin 2x + b_3 e^{-9t} \sin 3x + \dots \quad (3.44)$$

This turns out to be the general solution to the PDE  $\frac{\partial \theta}{\partial t} = \nu \frac{\partial^2 \theta}{\partial x^2}$  with the boundary conditions  $\theta(0, t) = 0$  and  $\theta(\pi, t) = 0$ .

#### 4. Separation of variables: solving a PDE with homogeneous boundary conditions

Hide Discussion



Add a Post

Show all posts	by recent activity
<input checked="" type="checkbox"/> <u>3D case</u>	3
<p>Is it possible to generalize this procedure to the 2D or 3D case? I'd like go depper into this after finishing <a href="#">this course</a>.</p>	
? <u>solution form</u>	4
<p>Re: Videos, good idea to show whole development</p>	2
<p>The title says it, I just thought I'd say it's great how in these recitation videos you keep the entire development on the board the whole time. This way you can instantly pause ...</p>	
<input checked="" type="checkbox"/> <u>Can solution to diff. eqn. involving x only be a sum?</u>	3
<p>Hi, at about 8:50 of the video he says that <math>A \sin(nx)</math> is the solution to the differential equation (the one that involves x only). Would it be correct to say that more generally a s...</p>	
<input checked="" type="checkbox"/> <u>Separation of Variables?</u>	2
<p>The assumption that the overall solution to the heat equation can be written as the product of <math>X(x) \cdot T(t)</math> certainly works. How do we know that there isn't a function that is not...</p>	

