

< Previous	 	 		Next >
-------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------	---------------------------

12.4.3 In Preparation for this Week's Enrichment

 Bookmark this page

Week 12 due Dec 29, 2023 10:42 IST

12.4.3 In Preparation for this Week's Enrichment

No video for this unit

Reading Assignment

0 points possible (ungraded)
Read Unit 12.4.3 of the notes. [\[LINK\]](#)

☒ Done

✓

Submit

✓ Correct

Discussion

Topic: Week 12 / 12.4.3

Hide Discussion

Add a Post

Show all posts

by recent activity

☒ [Actual restrictions of the Rayleigh quotient?](#)

3

Homework 12.4.3.1

1/1 point (graded)
Let $A \in \mathbb{R}^{n \times n}$ and x equal an eigenvector of A . Assume that x is real valued as is the eigenvalue λ with $Ax = \lambda x$.

$\lambda = \frac{x^T Ax}{x^T x}$ is the eigenvalue associated with the eigenvector x .

Always ✓ Answer: Always

$Ax = \lambda x$ implies that $x^T Ax = x^T (\lambda x) = \lambda x^T x$. But $x^T x \neq 0$ since x is an eigenvector. Hence $\lambda = x^T Ax / (x^T x)$.

Submit

Answers are displayed within the problem

Homework 12.4.3.2

1/1 point (graded)
Let $A \in \mathbb{R}^{n \times n}$ be nonsingular, $\lambda \in \Lambda(A)$, and $Ax = \lambda x$. Then $A^{-1}x = \frac{1}{\lambda}x$.

TRUE ✓ Answer: TRUE

$Ax = \lambda x$ means that $\frac{1}{\lambda}A^{-1}Ax = \frac{1}{\lambda}A^{-1}\lambda x$ which means that $\frac{1}{\lambda}x = A^{-1}x$.

Calculator

Submit

Answers are displayed within the problem

Homework 12.4.3.3

10/10 points (graded)

Let $A \in \mathbb{R}^{n \times n}$ and $\lambda \in \Lambda(A)$. Then $(\lambda - \mu) \in \Lambda(A - \mu I)$.

TRUE

Answer: TRUE

Let $Ax = \lambda x$ for $x \neq 0$. Then

$$(A - \mu I)x = Ax - \mu Ix = Ax - \mu x = \lambda x - \mu x = (\lambda - \mu)x.$$

Submit

Answers are displayed within the problem

< Previous

Next >

© All Rights Reserved



edX

[About](#)

[Affiliates](#)

[edX for Business](#)

[Open edX](#)

[Careers](#)

[News](#)

Legal

[Terms of Service & Honor Code](#)

[Privacy Policy](#)

[Accessibility Policy](#)

[Trademark Policy](#)

[Sitemap](#)

[Cookie Policy](#)

[Your Privacy Choices](#)

Calculator

Connect

[Idea Hub](#)

[Contact Us](#)

[Help Center](#)

[Security](#)

[Media Kit](#)



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)