

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Lecture 6: Introduction to Hypothesis Testing, and Type 1 and](#)

[Course](#) > [Unit 2 Foundation of Inference](#) > [Type 2 Errors](#)


> 16. Level of a Statistical test

Currently enrolled in **Audit Track** (expires December 25, 2019) [Upgrade \(\\$300\)](#)

16. Level of a Statistical test

Level of a Statistical test

Generating Speech Output



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Testing the Support of a Uniform Variable: Level and Threshold

Generating Speech Output

2/2 points (graded)

As in the problems on the previous page, let $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Unif}[0, \theta]$ for an unknown parameter θ and we designed the statistical test

$$\psi_n = \mathbf{1}(\max_{1 \leq i \leq n} X_i > 1/2)$$

to decide between the null and alternative hypotheses

$$H_0 : \theta \leq 1/2$$

$$H_1 : \theta > 1/2.$$

Let $\alpha_{\psi_n}(\theta)$ and $\beta_{\psi_n}(\theta)$ be the type 1 and type 2 errors respectively.

Recall from lecture that a test ψ has **level** α if

$$\alpha \geq \alpha_{\psi}(\theta) \quad \text{for all } \theta \in \Theta_0,$$

where $\alpha_{\psi} = \mathbf{P}_{\theta}(\psi = 1)$ is the type 1 error. We will often use the word "level" to mean the "smallest" such level, i.e. the least upper bound of the type 1 error, defined as follows:

$$\alpha = \sup_{\theta \in \Theta_0} \alpha_{\psi}(\theta)$$

Here, $\sup_{\theta \in \Theta_0}$ stands for the supremum over all values of θ within Θ_0 . If Θ_0 is a closed (*resp.* closed half-interval), and if $\alpha_{\psi}(\theta)$ is continuous (*resp.* continuous and decreasing as it approaches infinity), then its supremum equals the maximum.

Using the graph of the errors on the previous page, what is the smallest level α of the test ψ_n ?

$\alpha =$

✓ Answer: 0

How should the threshold of the test be changed to increase the smallest level α ? In other words, consider tests of the form

$$\psi_{n,C} = \mathbf{1}(\max_{1 \leq i \leq n} X_i > C)$$

where C is the threshold. In the original test above, $C = 1/2$. What should the value of C be so that the level of $\psi_{n,C}$ is greater than the level of the $\psi_{n,1/2}$?

(Think of how the graph of $\mathbf{P}_\theta(\psi_C)$ changes with the threshold C .)

☐ $C > 1/2$

☒ $C < 1/2$



Solution:

Since the type 1 error $\alpha_{\psi_n}(\theta)$ is constantly zero over Θ_0 , the smallest level of this test ψ is $\alpha = 0$.

To increase the smallest level α from 0, note that $\mathbf{P}_\theta\left(\max_{1 \leq i \leq n} X_i > C\right) = 0$ if and only if $\theta \leq C$. This means the constant zero region of graph of $\mathbf{P}_\theta(\psi_C) = 0$ shifts to the right as C increases from $1/2$, and to the left as C decreases from $1/2$. Since the maximum of type 1 error occurs at the boundary $\theta = 1/2$, this means $C < 1/2$ is required for the level to be positive.

Remark: The reason behind increasing the level in this example is to increase the power of the test from 0. In general, one of the first requirements of a test is to have a small-enough level so that the probability of concluding a false positive, (i.e. rejecting the null while the null is true) is controlled.

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Generating Speech Output

Testing the Support of a Uniform Variable: Determine the Threshold

1/1 point (graded)

As above, let $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Unif}[0, \theta]$ for an unknown parameter θ and consider tests of the form

$$\psi_{n,C} = \mathbf{1}(\max_{1 \leq i \leq n} X_i > C)$$

to decide between the null and alternative hypotheses

$$H_0 : \theta \leq 1/2$$

$$H_1 : \theta > 1/2.$$

Let $\alpha_{\psi_{n,C}}(\theta)$ and $\beta_{\psi_{n,C}}(\theta)$ be the type 1 and type 2 errors respectively.

Determine the smallest threshold C such that the test $\psi_{n,C}$ has level α .

(Enter the roots of x as a power of x , e.g. enter $\mathbf{x^{(1/3)}}$ for $\sqrt[3]{x} = x^{1/3}$.)

$C =$

$(1-\alpha)^{(1/n)/2}$

✓ Answer: $1/2 * (1-\alpha)^{(1/n)}$

$$\frac{(1-\alpha)^{\frac{1}{n}}}{2}$$

STANDARD NOTATION

Solution:

Following similar computation as in a previous problem where $C = 1/2$, we have $\mathbf{P}_{\theta}(\psi_{n,C} = 1) = 1 - \left(\frac{C}{\theta}\right)^n$. Since the smallest level is

$$\begin{aligned} \alpha &= \max_{\theta \in \Theta_0} p_{\theta}(\psi_{n,C} = 1) \\ &= p_{1/2}(\psi_{n,C} = 1) = 1 - \left(\frac{C}{1/2}\right)^n, \end{aligned}$$

Generating Speech Output

a test with threshold $C = \frac{1}{2}\sqrt{1-\alpha}$ or smaller will have level α .

Remark: Notice the threshold C depends on n , α , as well as the value of θ at the boundary of Θ_0 and Θ_1 .

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 2 Foundation of Inference:Lecture 6: Introduction to Hypothesis Testing, and Type 1 and Type 2 Errors / 16. Level of a Statistical test

Add a Post

◀ All Posts

Confused about the shape of the curve

question posted 3 days ago by [XuYang2015](#)

I am a little confused here. According to the curve, it seems that level alpha is equal to power pi. It is not true, there must be something wrong with the curve. Actually, I don't understand why the curve can be continuous without interruption, a "S" shape. The x-axis of the curve is true mu? If it is, in the interval of theta 0, the curve should be a constant, because $\sqrt{n}*(\bar{X}_n - \mu)/\sigma$ is asymptotically normal, meanwhile \bar{X}_n is a r.v and C is a constant here, $P(\sqrt{n}*(\bar{X}_n - \mu)/\sigma > C)$ is a constant. Am I right?

This post is visible to everyone.

Add a Response


2 responses

Erocha (Community TA)

3 days ago

Do you mean this graph:

Generating Speech Output



note this is the value of $P_\mu[\psi = 1]$ (a.k.a. power function), and it is continuous indeed. If you have access to the book "All of Statistics" see example 10.2 in Chapter 10.

See this:

Example 11.2 Let $X_1, \dots, X_n \sim N(\mu, \sigma)$ where σ is known. We want to test $H_0 : \mu \leq 0$ versus $H_1 : \mu > 0$. Hence, $\Theta_0 = (-\infty, 0]$ and $\Theta_1 = (0, \infty)$. Consider the test:

reject H_0 if $T > c$

where $T = \bar{X}$. The rejection region is $R = \{x^n : T(x^n) > c\}$. Let Z denote a standard normal random variable. The power function is

$$\begin{aligned} \beta(\mu) &= P_\mu(\bar{X} > c) \\ &= P_\mu\left(\frac{\sqrt{n}(\bar{X} - \mu)}{\sigma} > \frac{\sqrt{n}(c - \mu)}{\sigma}\right) \\ &= P\left(Z > \frac{\sqrt{n}(c - \mu)}{\sigma}\right) \\ &= 1 - \Phi\left(\frac{\sqrt{n}(c - \mu)}{\sigma}\right). \end{aligned}$$

Generating Speech Output

This function is increasing in μ . Hence

$$\text{size} = \sup_{\mu < 0} \beta(\mu) = \beta(0) = 1 - \Phi\left(\frac{\sqrt{n}c}{\sigma}\right).$$

To get a size α test, set this equal to α and solve for c to get

$$c = \frac{\sigma\Phi^{-1}(1 - \alpha)}{\sqrt{n}}.$$

So we reject when $\bar{X} > \sigma\Phi^{-1}(1 - \alpha)/\sqrt{n}$. Equivalently, we reject when

$$\frac{\sqrt{n}(\bar{X} - 0)}{\sigma} > z_{\alpha}. \quad \blacksquare$$

posted 2 days ago by [CoolZ](#)

Add a comment

XuYang2015

2 days ago



Thank you for your detailed explanations. Just a small question to check if I correctly understand the example you gave (example 11.2). In the example, because the power function is continuously increasing, thus the sup of the power function in the interval $\Theta_0 (-\infty, 0]$ is equal to the inf of the power function in the interval $\Theta_1 (0, \infty)$. It means that if we control the level of the test statistics at α , the power of the test statistics is also α ? Am I right?

Generating Speech Output

Yes, with my understanding in this example.

$\alpha = \Phi\left(\frac{\sqrt{n}(c-0)}{\sigma}\right)$, $\text{power} = \lim_{\mu \rightarrow 0^+} \Phi\left(\frac{\sqrt{n}(c-\mu)}{\sigma}\right)$, giving this function is increasing in μ .

and we know $\Phi\left(\frac{\sqrt{n}(c-\mu)}{\sigma}\right)$ is continuous at 0 when $\sigma \neq 0$, thus $\alpha = \text{power}$

posted 2 days ago by [CoolZ](#)

Add a comment

Showing all responses

Add a response:

Preview

Submit

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

Generating Speech Output

Learn About Verified Certificates

© All Rights Reserved

Generating Speech Output