

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Lecture 3: Parametric Statistical](#)

[Course](#) > [Unit 2 Foundation of Inference](#) > [Models](#)

> 11. Identifiability exercises

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

11. Identifiability exercises

Identifiability of Statistical Models 2

1/1 point (graded)

Let $X_i = Y_i^2$ where $Y_1, \dots, Y_n \stackrel{iid}{\sim} \mathcal{U}([0, a])$ for some unknown parameter a . We observe the i.i.d. samples X_1, \dots, X_n , but not the Y_i 's themselves.

Hint: Compute the cdf of X_i .

Is the parameter a identifiable from the common distribution the X_i 's?

☒ Yes

☐ No



Solution:

Write $X_i \sim X$ and note that X is supported on the interval $[0, a^2]$. Let us compute the CDF of X in terms of a .

$$\mathbf{P}(X \leq t) = \mathbf{P}(Y \leq \sqrt{t}) = \min \left(\int_0^{\sqrt{t}} \frac{1}{a} dy, 1 \right) = \min \left(\frac{\sqrt{t}}{a}, 1 \right).$$

For different values of a , the CDF of X are different; hence a is identifiable.

Submit

You have used 1 of 1 attempt

i Answers are displayed within the problem

Identifiability of Statistical Models 3

1/1 point (graded)

Let $X_i = \mathcal{I}(Y_i \geq a/2)$ where $Y_1, \dots, Y_n \stackrel{iid}{\sim} \mathcal{U}([0, a])$ for some unknown parameter a . We observe the independent samples X_1, \dots, X_n but not the Y_i 's themselves.

Is the parameter a identifiable from the common distribution of the X_i 's?

☐ Yes

☒ No



Solution:

Note that X is a Bernoulli random variable with parameter $p := P(\mathcal{I}(Y_i \geq \frac{a}{2}) = 1) = P(Y_i \geq \frac{a}{2})$.

For any choice of a , we have by the distribution of Y_i that $p = P(Y_i \geq a/2) = 1/2$. Hence, for any choice of a , the random variable X is distributed as $\text{Ber}(1/2)$. The parameter a is not identifiable.

[Submit](#)

You have used 1 of 1 attempt

i Answers are displayed within the problem

Review of terminology

1/1 point (graded)

You have access to samples $X_1, \dots, X_n \stackrel{iid}{\sim} P_{\theta^*}$ where $\theta^* \in \mathbb{R}$ is a true, unknown parameter specifying the distribution. You construct a statistical model $((-\infty, \infty), \{P_{\theta}\}_{\theta \in \mathbb{R}})$ for this statistical experiment. Your goal is to uncover the true parameter θ^* .

Imagine that somehow you are able to figure out the true distribution P_{θ^*} . Which assumptions (individually, each on its own) below are sufficient to recover the true parameter θ^* from the distribution?

(Choose all that apply.)

☐ There is another value $\theta' \in \mathbb{R}$ such that $\theta' \neq \theta^*$ but P_{θ^*} and $P_{\theta'}$ are the same distribution.

☐ The given statistical model $((-\infty, \infty), \{P_{\theta}\}_{\theta \in \mathbb{R}})$ is well-specified.

☒ The parameter θ is identifiable for the given statistical model.

[Submit](#)

You have used 1 of 1 attempt

Discussion

Topic: Unit 2 Foundation of Inference:Lecture 3: Parametric Statistical Models / 11. Identifiability exercises

[Hide Discussion](#)[Add a Post](#)

[◀ All Posts](#)

2nd question

question posted 6 days ago by [stephenjersey](#).

What is the meaning of the 'I' in ' $X_i = I(Y_i \geq a/2)$ ' in the 2nd question 'Identifiability of Statistical Models 3'? Thanks.

This post is visible to everyone.



[markweitzman](#)

6 days ago - marked as answer 6 days ago by [ya_mukhin](#) (Staff)

Indicator function.



@stephenjersey, does that help or would a more detailed explanation be useful?

posted 6 days ago by [ya_mukhin](#) (Staff)



That's clear! Thanks!

posted 6 days ago by [stephenjersey](#).



Can it be assumed that $a > 0$ since the interval $[0, a]$ and $Y \sim U([0, a])$ only makes sense iff $a > 0$, so we can rule out the possibility of a non-positive zero?

posted 4 days ago by [sandipan dey](#)



Yes, that seems a reasonable assumption.

posted 3 days ago by [mrBB](#)



Thank you @mrBB, so I guess we can assume the same for the 1st question too, right?



posted about 12 hours ago by [sandipan dey](#).

Add a comment

Preview

Submit

[Learn About Verified Certificates](#)

© All Rights Reserved