



<u>Lecture 14: Wald's Test, Likelihood</u> <u>Ratio Test, and Implicit Hypothesis</u>

<u>Course</u> > <u>Unit 4 Hypothesis testing</u> > <u>Test</u>


> 11. Likelihood Ratio Test

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. **Upgrade now**

11. Likelihood Ratio Test Likelihood Ratio Test



https://courses.edx.org/courses/course-v1:MITx+18.6501x+3T2019/courseware/unit4/u04s04_hypotesting/

In the general form of the likelihood ratio test, we have an unknown parameter $\theta^* \in \mathbb{R}^d$, and we are deciding between two hypotheses of the form

$$egin{aligned} H_0: (heta_{r+1}^*, \dots, heta_d^*) &= (heta_{r+1}^{(0)}, \dots, heta_d^{(0)}) \ H_1: (heta_{r+1}^*, \dots, heta_d^*) &
eq (heta_{r+1}^{(0)}, \dots, heta_d^{(0)}) \,. \end{aligned}$$

for some $r \geq 0$.

Thus Θ_0 , the region defined by the null hypothesis, is

$$\Theta_0 := \{ \mathbf{v} \in \mathbb{R}^d : (v_{r+1}, \ldots, v_d) = (heta_{r+1}^{(0)}, \ldots, heta_d^{(0)}) \}$$

where $(heta_{r+1}^{(0)},\ldots, heta_d^{(0)})$ consists of *known* values.

The likelihood ratio test involves the test-statistic

$$T_{n}=2\left(\ell_{n}\left(\widehat{ heta_{n}}^{MLE}
ight)-\ell_{n}\left(\widehat{ heta_{n}}^{c}
ight)
ight)$$

where ℓ_n is the log-likelihood.

The estimator $\widehat{\theta_n^c}$ is the **constrained MLE** , and it is defined to be

$$\widehat{ heta_{n}^{c}} = \operatorname{argmax}_{ heta \in \Theta_{0}} \ell_{n}\left(X_{1}, \ldots, X_{n}; heta
ight).$$

Which of the following are possible? (Choose all that apply.)

$$oxedsymbol{igsquare} \ell_n\left(\widehat{ heta_n}^{MLE}
ight) < \ell_n\left(\widehat{ heta_n}^c
ight)$$

$$lacksquare \ell_n \, (\widehat{ heta_n}^{MLE}) = \ell_n \, (\widehat{ heta_n}^c)$$

$$lacksquare \ell_n \, (\widehat{ heta_n}^{MLE}) > \ell_n \, (\widehat{ heta_n}^c)$$



Solution:

Recall that the MLE is defined by the optimization problem

$$\widehat{ heta_{n}^{MLE}} = \operatorname{argmax}_{ heta \in \Theta} \ell_{n}\left(X_{1}, \ldots, X_{n}; heta
ight)$$

In particular, we find the maximizer over the \emph{entire} parameter space Θ . The constrained MLE

$$\widehat{ heta_{n}^{c}} = \operatorname{argmax}_{ heta \in \Theta_{0}} \ell_{n}\left(X_{1}, \ldots, X_{n}; heta
ight)$$

finds the maximum over a subset of Θ , so it is not possible that $\ell_n\left(\widehat{\theta_n}^{MLE}\right) < \ell_n\left(\widehat{\theta_n}^c\right)$. However, it may be the case that $\ell_n\left(\widehat{\theta_n}^{MLE}\right) = \ell_n\left(\widehat{\theta_n}^c\right)$ or $\ell_n\left(\widehat{\theta_n}^{MLE}\right) > \ell_n\left(\widehat{\theta_n}^c\right)$. In general, we will have that $\ell_n\left(\widehat{\theta_n}^{MLE}\right) \geq \ell_n\left(\widehat{\theta_n}^c\right)$.

Remark: The likelihood ratio test is a natural test in a situation where we only care about *some* (e.g., the last d-r coordinates) of the unknowns involved in the parameter $\theta^* \in \mathbb{R}^d$.

Submit

You have used 1 of 3 attempts

1 Answers are displayed within the problem

Concept Check: Test-Statistic for the Likelihood Ratio Test

1/1 point (graded)

Suppose we are hypothesis testing between a null and alternative of the form

$$H_0:(heta^*_{r+1},\ldots, heta^*_d)\ =(heta^{(0)}_{r+1},\ldots, heta^{(0)}_d)$$

$$H_1: (heta^*_{r+1}, \ldots, heta^*_d) \
eq (heta^{(0)}_{r+1}, \ldots, heta^{(0)}_d) \,.$$

Above, $\theta^* \in \mathbb{R}^d$ is an unknown parameter while the values $\theta_{r+1}^{(0)}, \dots, \theta_d^{(0)}$ are known. To perform the likelihood ratio test, we define the test statistic

$$T_{n}=2\left(\ell_{n}\left(\widehat{ heta_{n}^{MLE}}
ight)-\ell_{n}\left(\widehat{ heta_{n}^{c}}
ight)
ight)$$

Assume that the technical conditions needed for the MLE to be a consistent estimator are satisfied, and assume that the null-hypothesis is true.

Which of the following are true about the above test statistic T_n ? (Choose all that apply. Refer to the slides.)

 $ightharpoonup T_n$ is a pivotal statistic; *i.e.*, it converges to a pivotal distribution.

 $luellimits_n T_n$ is asymptotically normal.

$$lacksquare T_n \xrightarrow[n o \infty]{(d)} \chi^2_{d-r}$$

$$lineq T_n extstyle rac{(d)}{n o\infty} \chi^2_r$$



Solution:

We examine the choices in order.

• The first answer choice is correct. Under the null hypothesis,

$$T_n \xrightarrow[n \to \infty]{(d)} \chi^2_{d-r}.$$

The distribution χ^2_r is pivotal because it does not depend on the specific value of the true parameter θ^* . Hence T_n is also a pivotal statistic.

- The second answer choice is incorrect. T_n is not asymptotically normal; rather it is asymptotically a χ^2 random variable, as stated in the previous bullet. Note that the normal distribution and χ distribution are very different from each other (e.g., χ^2 has significantly heavier tails).
- The third answer choice is correct. As stated in the first bullet, $T_n \xrightarrow[n \to \infty]{(d)} \chi^2_{d-r}$ assuming the null hypothesis and the technical conditions mentioned in the problem statement.
- The fourth answer choice is incorrect. It is true that T_n converges to a χ^2 random variable, but this choice gives the wrong number of degrees of freedom.

Remark: Be careful not to be confused about the following point. While the parameter space corresponding to H_0 is $\Theta_0 = \mathbb{R}^r$ which, intuitively, has r free variables, the test statistic T_n converges to a χ^2 distribution with d-r degrees of freedom. This convergence fact follows from a technical result of Wilks, and we do not discuss aspects of its proof here.

Submit

You have used 1 of 3 attempts

1 Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 4 Hypothesis testing:Lecture 14: Wald's Test, Likelihood Ratio Test, and Implicit Hypothesis Test / 11. Likelihood Ratio Test

Add a Post

Show all posts

by recent activity ▼

There are no posts in this topic yet.

| × | | | |
|---|-----------------------------------|------------|--------------|
| | | | |
| | Learn About Verified Certificates | | |
| | 2001117112000 10111100 | © All Righ | nts Reserved |