

<u>Help</u> 🗘

sandipan_dey 🗸

<u>Course</u> <u>Progress</u> <u>Dates</u> <u>Discussion</u> <u>Syllabus</u> <u>Outline</u> <u>laff routines</u> <u>Community</u>

☆ Course / Week 9: Vector Spaces / 9.5 Span, Linear Independence, and Bases

(3)

Next >

9.5.2 Linear Independence

☐ Bookmark this page


< Previous

■ Calculator

Show all posts

Week 9 due Dec 9, 2023 18:12 IST Completed

9.5.2 Linear Independence



Add a Post

by recent activity >

? The column space of A and linearly independency of A's columns

Hi, In the last of this unit Dr. Robert van de Geijn said these two are equivalent: 1- The column space of A is R^n 2- A has linearly independent co...

Question on 9.5.2 Linear Independence
Greetings! I do not quite understand that a matrix, which has more columns than rows, indicates we have vectors in the null space. It is mentic

Homework 9.5.2.1

1/1 point (graded)

$$\operatorname{Span}\left(\left(egin{array}{c}1\0\1\end{array}
ight),\left(egin{array}{c}0\0\1\end{array}
ight)=\operatorname{Span}\left(\left(egin{array}{c}1\0\1\end{array}
ight),\left(egin{array}{c}0\0\1\end{array}
ight),\left(egin{array}{c}1\0\3\end{array}
ight)
ight)$$

TRUE ✓ ✓ Answer: TRUE

•
$$S\subset T$$
: Let $x\in \mathrm{Span}egin{pmatrix}1\\0\\1\end{pmatrix}$, $\begin{pmatrix}0\\0\\1\end{pmatrix}$ Then there exist $lpha_0$ and $lpha_1$ such that $x=lpha_0egin{pmatrix}1\\0\\1\end{pmatrix}+lpha_1egin{pmatrix}0\\0\\1\end{pmatrix}$. This in turn means that

$$x=lpha_0egin{pmatrix}1\0\1\end{pmatrix}+lpha_1egin{pmatrix}0\0\1\end{pmatrix}+(0)egin{pmatrix}1\0\3\end{pmatrix}.$$

Hence
$$x \in \operatorname{Span}\left(\begin{pmatrix}1 \ 0 \ 1\end{pmatrix}, \begin{pmatrix}0 \ 0 \ 1\end{pmatrix}, \begin{pmatrix}1 \ 0 \ 3\end{pmatrix}\right)$$

•
$$T\subset S$$
: Let $x\in \mathrm{Span}\left(egin{pmatrix}1\\0\\1\end{pmatrix},egin{pmatrix}0\\0\\1\end{pmatrix}+eta_1egin{pmatrix}0\\0\\1\end{pmatrix}+lpha_2egin{pmatrix}1\\0\\3\end{pmatrix}$. But $egin{pmatrix}1\\0\\3\end{pmatrix}=egin{pmatrix}1\\0\\0\\1\end{pmatrix}+2egin{pmatrix}0\\0\\0\\1\end{pmatrix}$. Hence

$$x=lpha_0egin{pmatrix}1\0\1\end{pmatrix}+lpha_1egin{pmatrix}0\0\1\end{pmatrix}+lpha_2egin{pmatrix}1\0\1\end{pmatrix}+2egin{pmatrix}0\0\1\end{pmatrix}\end{pmatrix}=(lpha_0+lpha_2)egin{pmatrix}1\0\0\1\end{pmatrix}+(lpha_1+2lpha_2)egin{pmatrix}0\0\0\1\end{pmatrix}.$$

Therefore
$$x \in \operatorname{Span}\left(\begin{pmatrix}1\\0\\1\end{pmatrix}, \begin{pmatrix}0\\0\\1\end{pmatrix}\right)$$
.

Submit

1 Answers are displayed within the problem

Homework 9.5.2.2

1/1 point (graded)

Let the set of vectors $\{v_0, v_1, \dots, v_{n-1}\} \subset \mathbb{R}^m$ be linearly dependent. Then at least one of these vectors can be written as a linear combination of the others.

TRUE ✓ ✓

✓ Answer: TRUE

 $\chi_0u_0+\chi_1u_1+\cdots+\chi_{n-1}u_{n-1}=0$ and for alleast one $j, 0 \leq j \leq n, \chi_j \neq 0$. Dut then

$$\chi_j a_j = -\chi_0 a_0 + -\chi_1 a_1 - \dots - \chi_{j-1} a_{j-1} - \chi_{j+1} a_{j+1} - \dots - \chi_{n-1} a_{n-1}$$

and therefore

$$a_j = -rac{\chi_0}{\chi_j} a_0 + -rac{\chi_1}{\chi_j} a_1 - \dots - rac{\chi_{j-1}}{\chi_j} a_{j-1} - rac{\chi_{j+1}}{\chi_j} a_{j+1} - \dots - rac{\chi_{n-1}}{\chi_j} a_{n-1}.$$

In other words, a_i can be written as a linear combination of the other n-1 vectors.

Submit

Answers are displayed within the problem

Homework 9.5.2.3

1/1 point (graded)

Let $U \in \mathbb{R}^{n \times n}$ be an upper triangular matrix with nonzeroes on its diagonal. Then its columns are linearly independent.

TRUE ✓ ✓ Answer: TRUE

We saw in a previous week that Ux=b has a unique solution if U is upper triangular with nonzeroes on its diagonal. Hence Ux=0 has the unique solution x=0 (the zero vector). This implies that U has linearly independent columns.

Submit

Answers are displayed within the problem

Homework 9.5.2.4

1/1 point (graded)

Let $L \in \mathbb{R}^{n \times n}$ be a lower triangular matrix with nonzeroes on its diagonal. Then its rows are linearly independent. (Hint: How do the rows of L relate to the columns of L^T ?)

ALWAYS

✓ Answer: ALWAYS

We saw in a previous week that Lx = b has a unique solution if L is lower triangular with nonzeroes on its diagonal. Hence Lx = 0 has the unique solution x = 0 (the zero vector). This implies that L has linearly independent columns.

Submit

Answers are displayed within the problem

Previous

Next >

© All Rights Reserved



edX

About

Affiliates

edX for Business

<u>Open edX</u>

Careers

News

Legal

Terms of Service & Honor Code

Privacy Policy

Accessibility Policy

<u>Trademark Policy</u>

<u>Sitemap</u>

Cookie Policy

Your Privacy Choices

Connect

<u>Idea Hub</u>

Contact Us

Help Center

<u>Security</u>

Media Kit















© 2023 edX LLC. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>