


L2 PROBLEM 4 (3/3 points)

1. Are the following two distributions equivalent?

```
import random
def dist1():
    return random.random() * 2 - 1

def dist2():
    if random.random() > 0.5:
        return random.random()
    else:
        return random.random() - 1
```

- ☒ Yes 
- ☐ No


EXPLANATION:

The random.random() distribution is uniform, so both `dist1` and `dist2` are a uniform distribution over [-1.0, 1.0).

2. Are the following two distributions equivalent?

```
import random
def dist3():
    return int(random.random() * 10)

def dist4():
    return random.randrange(0, 10)
```

- ☒ Yes 
- ☐ No

EXPLANATION:

The random.random() distribution is uniform, and so is the random.randrange() distribution, so both `dist3` and `dist4` are a discrete uniform distribution over [0, 1, 2, 3, 4, 5, 6, 7, 8, 9].

3. Are the following two distributions equivalent?

```
import random
def dist5():
    return int(random.random() * 10)

def dist6():
    return random.randint(0, 10)
```

- ☐ Yes

EXPLANATION:

The `random.random()` distribution is uniform, and so is the `random.randint()` distribution. However unlike `random.randrange(start, end)`, `random.randint(start, end)` returns a distribution that is inclusive of both the given start and end points.

Thus `dist5` is a discrete uniform distribution over `[0, 1, 2, 3, 4, 5, 6, 7, 8, 9]`, but `dist6` is a discrete uniform distribution over `[0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10]`.

You can code a simple simulation to see what a distribution looks like using dictionaries:

```
d1 = {}
for i in range(10000):
    x = random.randrange(10)
    d1[x] = d1.get(x, 0) + 1
d2 = {}
for i in range(10000):
    x = int(random.random()*10)
    d2[x] = d2.get(x, 0) + 1
d3 = {}
for i in range(10000):
    x = random.randint(0, 10)
    d3[x] = d3.get(x, 0) + 1
```

Examine the values of the three dictionaries to see what sort of distribution results!

Question to ponder: Should all the values of the dictionaries be equal? That is, should `d1[x] == d1[y]` for all values of `x` and `y`, where `x != y` and both `x` and `y` are values in `[0, 1, 2, 3, 4, 5, 6, 7, 8, 9]`?

Check

Hide Answer

Show Discussion

 New Post



EdX offers interactive online classes and MOOCs from the world's best universities. Online courses from MITx, HarvardX, BerkeleyX, UTx and many other universities. Topics include biology, business, chemistry, computer science, economics, finance, electronics, engineering, food and nutrition, history, humanities, law, literature, math, medicine, music, philosophy, physics, science, statistics and more. EdX is a non-profit online initiative created by founding partners Harvard and MIT.

© 2014 edX, some rights reserved.

Terms of Service and Honor Code (<https://www.edx.org/edx-terms-service>)

About & Company Info

About

(<https://www.edx.org/about-us>)

News

(<https://www.edx.org/news>)

Contact

(<https://www.edx.org/contact>)


FAQ

(<https://www.edx.org/student-faq>)


Follow Us

 Twitter

(<https://twitter.com/edXOnline>)

 Facebook


(<http://www.facebook.com/EdxOnline>)

 Meetup

(<http://www.meetup.com/edX-Global-Community>)

 LinkedIn

(<http://www.linkedin.com/company-edx>)

 Google+

Privacy Policy (Revised 4/16/2014) (<https://www.edx.org/edx-privacy-policy>)

edX Blog
(<https://www.edx.org/edx-blog>)

(<https://plus.google.com/+edXOnline>)

Donate to edX
(<https://www.edx.org/donate>)

Jobs at edX
(<https://www.edx.org/jobs>)