



Bookmarks

- ▶ [Module 1: The Basics of R and Introduction to the Course](#)
- ▶ [Entrance Survey](#)
- ▶ [Module 2: Fundamentals of Probability, Random Variables, Distributions, and Joint Distributions](#)
- ▶ [Module 3: Gathering and Collecting Data, Ethics, and Kernel Density Estimates](#)
- ▶ [Module 4: Joint, Marginal, and Conditional Distributions & Functions of Random Variable](#)

Module 8: Causality, Analyzing Randomized Experiments, & Nonparametric Regression > Analyzing Randomized Experiments > More on the Fisher Exact Test - Quiz

## More on the Fisher Exact Test - Quiz

🔖 Bookmark this page

### Question 1

1/1 point (graded)

In this segment, Professor Duflo describes the process for conducting the Fisher exact test. In this test, all possible \_\_\_\_\_ are tested while keeping \_\_\_\_\_ fixed. This is called \_\_\_\_\_.


- ☐ a. Observations ; treatment assignments ; bootstrapping
- ☐ b. Treatment assignments ; outcomes ; bootstrapping
- ☐ c. Observations ; treatment assignments ; a permutation test
- ☒ d. Treatment assignments ; the observations ; a permutation test ✓

### Explanation


In the Fisher exact test, (simulated) treatment effects are calculated under all possible treatment assignments, keeping the number of treated and control units the same as in the original assignment and holding the observations themselves constant. For example, we first assume that the first half of

- ▶ [Module 5: Moments of a Random Variable, Applications to Auctions, & Intro to Regression](#)
- ▶ [Module 6: Special Distributions, the Sample Mean, the Central Limit Theorem, and Estimation](#)
- ▶ [Module 7: Assessing and Deriving Estimators - Confidence Intervals, and Hypothesis Testing](#)
- ▼ [Module 8: Causality, Analyzing Randomized Experiments, & Nonparametric Regression](#)

### Causality

Finger Exercises due Nov 21, 2016  
at 05:00 IST 

### Analyzing Randomized Experiments

Finger Exercises due Nov 21, 2016  
at 05:00 IST 

the units are treated and the second are control, and we compute the treatment effect under this assignment.

Submit

You have used 1 of 2 attempts

✓ Correct (1/1 point)


### Question 2

1/1 point (graded)


After calculating the treatment effect under all possible treatment assignments, the next step is to:

- ☒ a. Count the number of cases in which the absolute value of the simulated treatment effect is greater than the treatment effect actually observed, divide by the total number of cases, this is the relevant p-value ✓
- ☐ b. Count the number of cases in which the absolute value of the simulated treatment effect is less than the treatment effect actually observed, divide by the total number of cases, this is the relevant p-value
- ☐ c. Average the p-values from each of the possible treatment effects
- ☐ d. Count the number of cases when the treatment effect is at least as large as the observed case

**Use of Randomization and****Nonparametric Regression**

Finger Exercises due Nov 21, 2016  
at 05:00 IST 

**Module 8: Homework**

Homework due Nov 14, 2016 at  
05:00 IST 

► Exit Survey

**Explanation**

The next step in the Fisher exact test would be to count up the number of cases in which the absolute value of the calculated treatment effect is greater than the treatment effect actually observed, and divide by the total number of cases to get the relevant p-value: under the sharp null hypothesis (the treatment effect is zero for everyone), the observed treatment effect should be somewhere in the middle of the distribution of simulated differences. The null will be rejected if the observed treatment effect is very much to the right (or the left) of the distribution.

Submit

You have used 1 of 2 attempts

✓ Correct (1/1 point)

**Question 3**

1/1 point (graded)

In the honey example discussed in class, there are six observations and 20 possible treatment assignments (only one of which is actually observed). For 16 of the possible treatment assignments, the absolute value of the calculated treatment effect is larger than the one actually observed (which was 1), such that the associated (exact) p-value is 0.8.

True or false: This p-value tells us that we can reject the null of no effect.

☐ a. True

☒ b. False ✓

**Explanation**

This is false. The p-value of 0.8 is large, which indicates that the observed treatment effect of 1 is likely due to chance, and we can conclude that the observed treatment effect of 1 is insignificant.

[Submit](#)

You have used 1 of 2 attempts

✓ Correct (1/1 point)

**Discussion****Topic:** Module 8 / More on the Fisher Exact Test - Quiz[Show Discussion](#)

© All Rights Reserved



© 2016 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.



