

[Course](#)

[Progress](#)

[Dates](#)


[Discussion](#)


[Syllabus](#)

[Outline](#)

[laff routines](#)

[Community](#)

 [Course](#) / [Week 3: Matrix-Vector Operations](#) / [3.2 Special Matrices](#)



< Previous	 	 	 	 			Next >
------------	---	---	--	---	---	---	--------

3.2.5 Transpose Matrix

 Bookmark this page

Week 3 due Oct 18, 2023 06:12 IST

3.2.5 Transpose Matrix

Summary

$$A = \begin{pmatrix} \alpha_{0,0} & \alpha_{0,1} & \cdots & \alpha_{0,n-1} \\ \alpha_{1,0} & \alpha_{1,1} & \cdots & \alpha_{1,n-1} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{m-1,0} & \alpha_{m-1,1} & \cdots & \alpha_{m-1,n-1} \end{pmatrix}$$
$$A^T = \begin{pmatrix} \alpha_{0,0} & \alpha_{0,1} & \cdots & \alpha_{m-1,0} \\ \alpha_{0,1} & \alpha_{1,1} & \cdots & \alpha_{m-1,1} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{0,n-1} & \alpha_{1,n-1} & \alpha_{2,n-1} & \alpha_{m-1,n-1} \end{pmatrix}$$

23 / 23

▶ 5:46 / 5:46

▶ 2.0x 🔊 🗑️ 📄 🗣️

Video

📄 [Download video file](#)

Transcripts

📄 [Download SubRip \(.srt\) file](#)

📄 [Download Text \(.txt\) file](#)

Reading Assignment

0 points possible (ungraded)
Read Unit 3.2.5 of the notes. [\[LINK\]](#)

☒ Done

✓

Submit

✓ Correct

Discussion

Topic: Week 3 / 3.2.5

Hide Discussion

Add a Post

Show all posts ▼

by recent

Calculator

https://learning.edx.org/course/course-v1:UTAustinX+UT.5.05x+1T2022/block-v1:UTAustinX+UT.5.05x+1T2022+type@sequential+block@8ae8cc0cab746e3903a9537de97d6ef/block-v1:UTAustinX+UT.5.05x+1T20... 2/7

<div><div>?</div><div>Picture Flame likes it, but Matlab does not. Getting a "input matrices must have same number of columns"</div><div>function [B_out] = Transpose_unb(A, B) [AL, AR] = FLA_Part_1x2(A, ... 0, 'FLA_LEFT'); [BT, ... BB] = FLA_Part_2x1(B, ... 0, 'FLA_TOP'); while...</div></div>	3
<div><div>?</div><div>'T' on b1t</div><div>Hi, I just wanted ask about the transpose label on b1t. At 4:50, professor van de Geijn explains that 'it has to do with the fact that it's a row of B...</div></div>	3
<div><div>?</div><div>Transpose_unb_var2(A, B) not working correctly in MATLAB</div><div>Anyone else getting an error transposing using the alternative code? The first row is transposed fine but the following rows stay the same. The f...</div></div>	6
<div><div></div><div>Homework 3.2.5.8 - is it clearer if the question is changed to an always/sometimes/never question?</div><div>While an identity matrix remains unchanged after transposition. Other matrices (i.e. symmetric matrices) have the same property.</div></div>	4

Homework 3.2.5.1

15/15 points (graded)

Let $A = \begin{pmatrix} -1 & 0 & 2 & 1 \\ 2 & -1 & 1 & 2 \\ 3 & 1 & -1 & 3 \end{pmatrix}$ and $x = \begin{pmatrix} -1 \\ 2 \\ 4 \end{pmatrix}$.

What is $A^T =$

<input type="text" value="-1"/>	✓ Answer: -1	<input type="text" value="2"/>	✓ Answer: 2	<input type="text" value="3"/>	✓ Answer: 3
<input type="text" value="0"/>	✓ Answer: 0	<input type="text" value="-1"/>	✓ Answer: -1	<input type="text" value="1"/>	✓ Answer: 1
<input type="text" value="2"/>	✓ Answer: 2	<input type="text" value="1"/>	✓ Answer: 1	<input type="text" value="-1"/>	✓ Answer: -1
<input type="text" value="1"/>	✓ Answer: 1	<input type="text" value="2"/>	✓ Answer: 2	<input type="text" value="3"/>	✓ Answer: 3

What is $x^T =$

<input type="text" value="-1"/>	<input type="text" value="2"/>	<input type="text" value="4"/>
✓	✓	✓
Answer: -1	Answer: 2	Answer: 4

Explanation

$$A^T = \begin{pmatrix} -1 & 0 & 2 & 1 \\ 2 & -1 & 1 & 2 \\ 3 & 1 & -1 & 3 \end{pmatrix}^T = \begin{pmatrix} -1 & 2 & 3 \\ 0 & -1 & 1 \\ 2 & 1 & -1 \\ 1 & 2 & 3 \end{pmatrix}$$
 and

$$x^T = \begin{pmatrix} -1 \\ 2 \\ 4 \end{pmatrix} = (-1 \quad 2 \quad 4)$$

Submit

i Answers are displayed within the problem

Homework 3.2.5.2

1/1 point (graded)

Consider the following algorithm.

Algorithm: $[B] := \text{TRANSPOSE_ALTERNATIVE}(A, B)$

Partition $A \rightarrow \left(\begin{array}{c} A_T \\ \cdot \end{array} \right)$, $B \rightarrow \left(\begin{array}{c|c} B_L & B_R \end{array} \right)$

Calculator

where A_T has 0 rows, B_L has 0 columns

while $m(A_T) < m(A)$ do

 Repartition

$$\left(\frac{A_T}{A_B} \right) \rightarrow \left(\frac{A_0}{a_1^T} \right), \left(B_L \mid B_R \right) \rightarrow \left(B_0 \mid b_1 \mid B_2 \right)$$

 where a_1 has 1 row, b_1 has 1 column

Continue with

$$\left(\frac{A_T}{A_B} \right) \leftarrow \left(\frac{A_0}{a_1^T} \right), \left(B_L \mid B_R \right) \leftarrow \left(B_0 \mid b_1 \mid B_2 \right)$$

endwhile

What belongs in the blank?

(Check all that apply)

✓ $b_1 := (a_1^T)^T$

$$\square \quad a_1 := b_1^T$$

□ $a_1 := 0$

□ $b_1 := 0$



Submit

i Answers are displayed within the problem

Homework 3.2.5.3

1/1 point (graded)

Implement the functions

- `Transpose_unb(A, B)`
- `Transpose_alternative_unb(A, B)`

(As before, implement as many as you enjoy implementing and/or until you "get the point". Then move on. We suggest you implement at least one of these.)

Some links that will come in handy:

- Spark (alternatively, open the file LAFF-2.0xM/Spark/index.html)
- PictureFLAME (alternatively, open the file LAFF-2.0xM/PictureFLAME/PictureFLAME.html)

 Calculator

You will need these in many future exercises. Bookmark them!

☒ Done/Skip



Answer:

- View a document that we put together that has most algorithms and MATLAB implementations that are homework problems in this week:

Week 3 algorithms and implementations.

This document is best viewed two pages, side by side, so that you can see the algorithm on the left and its implementation on the right.

Submit

Answers are displayed within the problem

Homework 3.2.5.4

1/1 point (graded)

The transpose of a lower triangular matrix is an upper triangular matrix.

Always

Answer: Always

Submit

Answers are displayed within the problem

Homework 3.2.5.5

1/1 point (graded)

The transpose of a strictly upper triangular matrix is a strictly lower triangular matrix.

Always

Answer: Always

Submit

Answers are displayed within the problem

Homework 3.2.5.6

1/1 point (graded)

The transpose of the identity matrix is the identity matrix.

Always



Submit

Correct (1/1 point)

Homework 3.2.5.7

8/8 points (graded)

$\begin{pmatrix} 0 & 1 \end{pmatrix}^T =$

Calculator

$\begin{pmatrix} 1 & 0 \end{pmatrix}$

<input type="text" value="0"/>	✓ Answer: 0	<input type="text" value="1"/>	✓ Answer: 1
<input type="text" value="1"/>	✓ Answer: 1	<input type="text" value="0"/>	✓ Answer: 0

$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}^T =$

<input type="text" value="0"/>	✓ Answer: 0	<input type="text" value="1"/>	✓ Answer: 1
<input type="text" value="-1"/>	✓ Answer: -1	<input type="text" value="0"/>	✓ Answer: 0

Submit

Answers are displayed within the problem

Homework 3.2.5.8

1/1 point (graded)
If $\mathbf{A} = \mathbf{A}^T$ then $\mathbf{A} = \mathbf{I}$ (The identity matrix).

FALSE

▼

✓ Answer: FALSE

Explanation
See Homework 3.2.5.7

Submit

Answers are displayed within the problem



[News](#)

Legal

- [Terms of Service & Honor Code](#)
 - [Privacy Policy](#)
 - [Accessibility Policy](#)
 - [Trademark Policy](#)
 - [Sitemap](#)
 - [Cookie Policy](#)
 - [Your Privacy Choices](#)
-

Connect

- [Idea Hub](#)
 - [Contact Us](#)
 - [Help Center](#)
 - [Security](#)
 - [Media Kit](#)
-



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)