



< Previous

✓

✓

Next >

11.2.3 Newton-Raphson method

 Bookmark this page

MO2.10

The Newton-Raphson method (often referred to just as Newton's method) fits a tangent line to the point $(x^k, r(x^k))$ on the graph of r , and defines x^{k+1} at the intersection of this tangent line with the x axis. The slope of this graph at the point x^k is $r'(x^k)$, and so the tangent line at $(x^k, r(x^k))$ satisfies the following equation,

$$r_{\text{tan}}(x) = r(x^k) + (x - x^k) r'(x^k),$$

Then, finding where this tangent intersects the x axis, defines the Newton iterate: $r_{\text{tan}}(x^{k+1}) = 0$. Specifically,

$$x^{k+1} = x^k - \frac{r(x^k)}{r'(x^k)}. \tag{11.7}$$

An alternative (but equivalent) derivation is to perform a first-order Taylor series approximation of $r(x)$ about the current iterate x^k :

$$r(x^k + \Delta x) \approx r(x^k) + r'(x^k) \Delta x$$

and then set this Taylor series approximation to zero to find Δx .

$$r(x^k) + r'(x^k) \Delta x = 0 \Rightarrow \Delta x = -\frac{r(x^k)}{r'(x^k)}$$

And then update x^k by Δx :

$$x^{k+1} = x^k + \Delta x = x^k - \frac{r(x^k)}{r'(x^k)}$$

which is the same result as Equation (11.7).

Convergence for Newton's method is very fast, when it occurs. Under some (reasonable) conditions, the error $\epsilon^k \equiv |x^* - x^k|$ obeys the following inequality:



$$|\epsilon^{k+1}| \leq C(\epsilon^k)^2,$$

Discussions

All posts sorted by recent activity

edX

< Previous

Next >

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)

Video on Newton's method and its application



[Start of transcript. Skip to the end.](#)

