

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Course](#) > [Unit 2 Foundation of Inference](#) > [Lecture 5: Delta Method and Confidence Intervals](#)

12. Interpretation of the frequentist  
> Confidence Interval

**Audit Access Expires Dec 24, 2019**

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

## 12. Interpretation of the frequentist Confidence Interval

### Frequentist Interpretation of a Confidence Interval

we're not really going to talk about them except for their connections to hypothesis testing.

So next week, we'll start hypothesis testing in a slightly mild way, mostly coming from--

so you will have some exercises that like are soft and connect them to confidence intervals.


There's going to be a lot of terminology.

We'll talk about type I, type II error, one-sided, two-sided tests.

We're going to talk about power.

We're going to talk about p values.

We're going to see a bunch of examples.



### Video

[Download video file](#)

### Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

## Frequentist Interpretation of a Confidence Interval

1/1 point (graded)

In a particular experiment, you gather data in the form of a sample  $X_1, \dots, X_n \stackrel{iid}{\sim} P_\theta$ , and construct a confidence interval  $\mathcal{I}$  with level 90% for the true (unknown) parameter  $\theta$ .

After conducting the experiment, there are two possibilities:

- $\mathcal{I}$  contains  $\theta$  (We refer to this as a **success**.)

we're going to see a bunch of new words.

So make sure that you're here, or make sure that you take a look at the notes.

Again, office hours this afternoon are actually right upstairs in 2-290.

- $\mathcal{I}$  does not contain  $\theta$  (We refer to this as a **failure**.)

Suppose you repeat the experiment above  $T$  total times, and assume that the experiments are jointly independent. Moreover, the value of the unknown parameter  $\theta$ , is always assumed to be the same. After conducting these  $T$  experiments, you will have constructed  $T$  confidence intervals  $\mathcal{I}_1, \mathcal{I}_2, \dots, \mathcal{I}_T$ .

As  $T$  grows very large, what percentage of experiments do you expect to be successes?

✓ Answer: 90 %

**Solution:**

By the definition of confidence interval, we know that for the  $j$ -th experiment ( $1 \leq j \leq T$ ) that

$$P(\mathcal{I}_j \ni \theta) = 90\%.$$

Consider the indicator random variables  $\mathbf{1}(\theta \in \mathcal{I}_1), \mathbf{1}(\theta \in \mathcal{I}_2), \dots, \mathbf{1}(\theta \in \mathcal{I}_T)$ . Since the experiments are jointly independent, this means that  $\mathbf{1}(\theta \in \mathcal{I}_1), \mathbf{1}(\theta \in \mathcal{I}_2), \dots, \mathbf{1}(\theta \in \mathcal{I}_T)$  are independent. Moreover, for all  $j$ , the random variable  $\mathbf{1}(\theta \in \mathcal{I}_j)$  is Bernoulli because it can only take value 0 or 1. It follows that  $\mathbf{1}(\theta \in \mathcal{I}_1), \mathbf{1}(\theta \in \mathcal{I}_2), \dots, \mathbf{1}(\theta \in \mathcal{I}_T)$  are identically distributed, because for all  $j$ ,

$$P(\mathbf{1}(\theta \in \mathcal{I}_j) = 1) = P(\mathcal{I}_j \ni \theta) = 90\%.$$

In summary,  $\mathbf{1}(\theta \in \mathcal{I}_1), \mathbf{1}(\theta \in \mathcal{I}_2), \dots, \mathbf{1}(\theta \in \mathcal{I}_T) \stackrel{iid}{\sim} \text{Ber}(0.9)$ . By the strong law of large numbers,

$$\lim_{T \rightarrow \infty} \frac{\sum_{j=1}^T \mathbf{1}(\theta \in \mathcal{I}_j)}{T} = \mathbb{E}[\mathbf{1}(\theta \in \mathcal{I}_j)] = 0.9$$

almost surely. Since

$$\frac{\sum_{j=1}^T \mathbf{1}(\theta \in \mathcal{I}_j)}{T} = \frac{\text{Number of successes}}{\text{Total number of experiments}},$$

the correct response is 90%.

Submit

You have used 1 of 1 attempt

**i** Answers are displayed within the problem

## Discussion

Hide Discussion

**Topic:** Unit 2 Foundation of Inference:Lecture 5: Delta Method and Confidence Intervals / 12. Interpretation of the frequentist Confidence Interval

Add a Post

Show all posts ▼

by recent activity ▼

💬 [Such a detailed solution...](#)  
[...for something so simple!](#)

2

Learn About Verified Certificates

© All Rights Reserved