



[Lecture 14: Wald's Test, Likelihood
Ratio Test, and Implicit Hypothesis](#)

[Course](#) > [Unit 4 Hypothesis testing](#) > [Test](#)

> 5. Introduction to Wald's Test

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

5. Introduction to Wald's Test

Introduction to Wald's Test



▶ 7:41 / 7:41

▶ 1.50x 🔊 🗑️ 📄 🗨️

Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Review: Manipulating Multivariate Gaussians

1/1 point (graded)

Recall that a **multivariate Gaussian** $\mathcal{N}(\vec{\mu}, \Sigma)$ is a random vector $\mathbf{Z} = [Z^{(1)}, \dots, Z^{(n)}]^T$ where $Z^{(1)}, \dots, Z^{(n)}$ are **jointly Gaussian**, meaning that the density of \mathbf{Z} is given by the joint pdf

$$f: \mathbb{R}^n \rightarrow \mathbb{R}$$

$$\mathbf{Z} \mapsto \frac{1}{(2\pi)^{n/2} \sqrt{\det(\Sigma)}} \exp\left(-\frac{1}{2}(\mathbf{Z} - \vec{\mu})^T \Sigma^{-1} (\mathbf{Z} - \vec{\mu})\right)$$

where

$$\vec{\mu}_i = \mathbb{E}[Z^{(i)}], \quad (\text{vector mean}).$$

$$\Sigma_{ij} = \text{Cov}(Z^{(i)}, Z^{(j)}) \quad (\text{positive definite covariance matrix}).$$

Suppose that $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \Sigma)$. Let \mathbf{M} denote an $n \times n$ matrix.

What is the distribution of \mathbf{MZ} ?

☐ $\mathcal{N}(\mathbf{0}, \Sigma)$

☐ $\mathcal{N}(\mathbf{0}, \mathbf{M}\Sigma)$

☐ $\mathcal{N}(\mathbf{0}, \Sigma\mathbf{M})$

☒ $\mathcal{N}(\mathbf{0}, \mathbf{M}\Sigma\mathbf{M}^T)$



Solution:

Linear transformations, e.g. \mathbf{MZ} , of Gaussian vectors are still Gaussian vectors. Hence, we only need to figure out the mean and covariance matrix of \mathbf{MZ} . By linearity of expectation:

$$\mathbb{E}[(\mathbf{MZ})_i] = \mathbb{E}\left[\sum_{j=1}^n \mathbf{M}_{ij} Z^{(j)}\right] = 0$$

for all i , so $\mathbb{E}[\mathbf{MZ}] = \mathbf{0}$. Or equivalently, in vector notation, (which is still correct, by linearity of expectation):

$$\mathbb{E}[\mathbf{MZ}] = \mathbf{M}\mathbb{E}[\mathbf{Z}] = \mathbf{M}\mathbf{0} = \mathbf{0}.$$

Next we compute the covariance. We will use the vector notation. Observe that

$$\Sigma = \mathbb{E}[\mathbf{ZZ}^T].$$

The covariance matrix of \mathbf{MZ} is given by

$$\mathbb{E}[(\mathbf{MZ})(\mathbf{MZ})^T] = \mathbb{E}[\mathbf{MZZ}^T\mathbf{M}^T] = \mathbf{M} \cdot \mathbb{E}[\mathbf{ZZ}^T] \mathbf{M}^T = \mathbf{M}\Sigma\mathbf{M}^T,$$

where we applied linearity of expectation for the third equality and the definition of Σ in the final equality.

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Aside: Rotation of Standard Gaussian

4/4 points (graded)

Suppose

$$\mathbf{M} = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix}.$$

Is it true that $\mathbf{M}^T \mathbf{M} = \mathbf{1}_{2 \times 2}$?

(Here $\mathbf{1}_{2 \times 2} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ is the identity matrix in 2 dimensions.)

☒ True

☐ False



Now, let $\mathbf{Z} \sim \mathcal{N}_2(\mathbf{0}, \mathbf{1}_{2 \times 2})$, i.e. \mathbf{Z} is a standard Gaussian in 2 dimensions. Is it true that $\mathbf{MZ} \sim \mathcal{N}_2(\vec{\mu}, \Sigma_{\mathbf{MZ}})$, for some $\vec{\mu}, \Sigma_{\mathbf{MZ}}$?

☒ True

☐ False



Find the mean $\vec{\mu} = \mathbb{E}[\mathbf{MZ}]$ and covariance matrix $\Sigma_{\mathbf{MZ}}$ of \mathbf{MZ} .

(Enter your answer as a vector or matrix. For example, type **[1,3]** for the vector $\begin{pmatrix} 1 \\ 3 \end{pmatrix}$; type **[[1,2],[5,1]]** for the matrix $\begin{pmatrix} 1 & 2 \\ 5 & 1 \end{pmatrix}$. Note the square brackets, and the commas as separators.)

$\vec{\mu} = \mathbb{E}[\mathbf{MZ}] =$

[0,0]

✓ Answer: [0,0]

$\Sigma_{\mathbf{MZ}} =$

[[1,0],[0,1]]

✓ Answer: [[1,0],[0,1]]

STANDARD NOTATION

Solution:

$$\begin{aligned} \mathbf{M}\mathbf{M}^T &= \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} \cos \phi & \sin \phi \\ -\sin \phi & \cos \phi \end{pmatrix} \\ &= \begin{pmatrix} \cos^2 \phi + \sin^2 \phi & \cos \phi \sin \phi - \cos \phi \sin \phi \\ \cos \phi \sin \phi - \cos \phi \sin \phi & \sin^2 \phi + \cos^2 \phi \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \end{aligned}$$

Hence $\mathbf{M}^T\mathbf{M} = \mathbf{1}_{2 \times 2}$ or equivalently $\mathbf{M}^T = \mathbf{M}^{-1}$

Remark: Geometrically, \mathbf{M} rotates a vector \mathbf{z} by an angle ϕ counterclockwise. Hence $\|\mathbf{M}\mathbf{z}\| = \|\mathbf{z}\|$ for any nonzero \mathbf{z} .

- Recall a main property of (multivariate) Gaussian variables is that any linear transformation of them remain (multivariate) Gaussian.
- Compute the mean and covariance of $\mathbf{M}\mathbf{Z}$:

$$\begin{aligned} \mathbb{E}[\mathbf{M}\mathbf{Z}] &= \mathbf{M}\mathbf{0} = \mathbf{0} \\ \Sigma_{\mathbf{M}\mathbf{Z}} &= \mathbf{M}\Sigma_{\mathbf{Z}}\mathbf{M}^T = \mathbf{M}\mathbf{1}_{2 \times 2}\mathbf{M}^T = \mathbf{M}\mathbf{M}^{-1} = \mathbf{1}_{2 \times 2}. \end{aligned}$$

Hence, $\mathbf{M}\mathbf{Z} \sim \mathcal{N}_d(\mathbf{0}, \mathbf{1}_{2 \times 2})$, i.e. a **standard** Gaussian vector.

Remark: Real matrices satisfying $\mathbf{M}^T = \mathbf{M}^{-1}$ (or equivalently $\mathbf{M}\mathbf{M}^T = \mathbf{M}^T\mathbf{M} = \mathbf{1}_{d \times d}$,) are called **orthogonal** matrices. In general, in d dimensions and for any orthogonal matrix \mathbf{M} , $\mathbf{M}\mathbf{Z}$ is also a **standard** multivariate Gaussian vector if \mathbf{Z} is a standard multivariate Gaussian.

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Review: Asymptotic Normality of the MLE

1/1 point (graded)

Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathbf{P}_{\theta^*}$ for some true parameter $\theta^* \in \mathbb{R}^d$. We construct the associated statistical model $(\mathbb{R}, \{\mathbf{P}_{\theta}\}_{\theta \in \mathbb{R}^d})$ and the maximum likelihood estimator $\hat{\theta}_n^{MLE}$ for θ^* .

Recall that, under some technical conditions,

$$\sqrt{n}(\hat{\theta}_n^{MLE} - \theta^*) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(0, \mathcal{I}(\theta^*)^{-1})$$

where $\mathcal{I}(\theta^*)$ denotes the Fisher information. That is, the MLE $\hat{\theta}_n^{MLE}$ is asymptotically normal with asymptotic covariance matrix $\mathcal{I}(\theta^*)^{-1}$.

Standardize the statement of asymptotic normality above. Answer by finding the power a of the Fisher information $\mathcal{I}(\theta^*)$ such that the following is true:

$$\sqrt{n\mathcal{I}(\theta^*)^a}(\hat{\theta}_n^{MLE} - \theta^*) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(0, I_{d \times d})$$

where $I_{d \times d}$ denotes the $d \times d$ identity matrix.

Hint: Use the result of the previous problem.

$a =$ ✓ Answer: 1/2

STANDARD NOTATION

Solution:

By the result of the previous problem, if $\mathbf{X} \sim \mathcal{N}(\mathbf{0}, \mathcal{I}(\theta^*)^{-1})$, then $\mathcal{I}(\theta^*)^{1/2}\mathbf{X}$ is mean 0 and has covariance matrix

$$\mathcal{I}(\theta^*)^{1/2}\mathcal{I}(\theta^*)^{-1}\left(\mathcal{I}(\theta^*)^{1/2}\right)^T = \mathcal{I}(\theta^*)^{1/2}\mathcal{I}(\theta^*)^{-1}\mathcal{I}(\theta^*)^{1/2} = I_{d \times d}.$$

Indeed, $\mathcal{I}(\theta^*)^{1/2}\mathbf{X} \sim \mathcal{N}(\mathbf{0}, I_{d \times d})$.

By the asymptotic normality of the MLE,

$$\sqrt{n}(\hat{\theta}_n^{MLE} - \theta^*) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(\mathbf{0}, \mathcal{I}(\theta^*)^{-1})$$

so that, by continuity,

$$\sqrt{n} \mathcal{I}(\theta^*)^{1/2} (\hat{\theta}_n^{MLE} - \theta^*) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{I}(\theta^*)^{1/2} \mathbf{N}(\mathbf{0}, \mathcal{I}(\theta^*)^{-1}) = \mathbf{N}(\mathbf{0}, I_{d \times d}).$$

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 4 Hypothesis testing: Lecture 14: Wald's Test, Likelihood Ratio Test, and Implicit Hypothesis Test / 5.
Introduction to Wald's Test

Add a Post

Show all posts ▼

by recent activity ▼

✓ [Fisher information matrix](#)

4

? [Fisher information](#)

1

💬 [\[Staff\] Inconsistence in denoting Identity matrices](#)

2

Learn About Verified Certificates

© All Rights Reserved