

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Course](#) > [Unit 3:...](#) > [6 Deco...](#) > 15. Pro...

15. Properties of the matrix exponential

Some properties and computation

MIT180312016-V027200

No comments



▶ 0:00 / 9:28

▶ 2.0x



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)[Download Text \(.txt\) file](#)

Properties of the matrix exponential

1. $e^{\mathbf{0}} = \mathbf{I}$ (here $\mathbf{0}$ is the zero matrix)

Proof: $e^{\mathbf{0}} = \mathbf{I} + \mathbf{0} + \frac{\mathbf{0}^2}{2!} + \cdots = \mathbf{I}$

2. $\frac{d}{dt}e^{\mathbf{A}t} = \mathbf{A}e^{\mathbf{A}t}$

Proof: Take the derivative of $e^{\mathbf{A}t}$ term by term.

3. If $\mathbf{AB} = \mathbf{BA}$, then $e^{\mathbf{A}+\mathbf{B}} = e^{\mathbf{A}}e^{\mathbf{B}}$. Here are a few (but not all) special cases when $\mathbf{AB} = \mathbf{BA}$:

- $\mathbf{A} = c\mathbf{I}$
- $\mathbf{B} = -\mathbf{A}$
- $\mathbf{B} = \mathbf{A}^{-1}$

Warning: In general, this fails, that is, if $\mathbf{AB} \neq \mathbf{BA}$, then usually $e^{\mathbf{A}+\mathbf{B}} \neq e^{\mathbf{A}}e^{\mathbf{B}}$, since there are counterexamples even for real 2x2 matrices.

4. If $\mathbf{A} = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$, then $e^{\mathbf{A}} = \begin{pmatrix} e^{\lambda_1} & 0 \\ 0 & e^{\lambda_2} \end{pmatrix}$.

Proof: $\mathbf{A}^2 = \begin{pmatrix} \lambda_1^2 & 0 \\ 0 & \lambda_2^2 \end{pmatrix}$, $\mathbf{A}^3 = \begin{pmatrix} \lambda_1^3 & 0 \\ 0 & \lambda_2^3 \end{pmatrix}$, and so on. Thus

$$e^{\mathbf{A}} = \mathbf{I} + \mathbf{A} + \frac{\mathbf{A}^2}{2!} + \dots = \begin{pmatrix} 1 + \lambda_1 + \frac{\lambda_1^2}{2!} + \dots & 0 \\ 0 & 1 + \lambda_2 + \frac{\lambda_2^2}{2!} + \dots \end{pmatrix} = \begin{pmatrix} e^{\lambda_1} & 0 \\ 0 & e^{\lambda_2} \end{pmatrix}.$$

(A similar statement holds for diagonal matrices of any size.)

5. Given \mathbf{A} is **diagonalizable**, that is,

$$\mathbf{A} = \mathbf{SDS}^{-1},$$

where \mathbf{D} is the diagonal matrix of eigenvalues and \mathbf{S} is the invertible matrix whose columns are eigenvectors:

$$\mathbf{D} = \begin{pmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_n \end{pmatrix}$$

$$\mathbf{S} = \begin{pmatrix} | & | & \cdots & | \\ \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \\ | & | & \cdots & | \end{pmatrix} \quad \text{where } \mathbf{v}_i \text{ corresponds to } \lambda_i.$$

Then

$$e^{\mathbf{A}} = \mathbf{S}e^{\mathbf{D}}\mathbf{S}^{-1}.$$

Proof: Expand $e^{\mathbf{A}}$ as a power series and use the following cancellations for each term:

$$\mathbf{A}^n = \mathbf{S}\underbrace{\mathbf{D}\mathbf{S}^{-1}\mathbf{S}}_{\text{cancels}}\cdots\underbrace{\mathbf{S}^{-1}\mathbf{S}\mathbf{D}\mathbf{S}^{-1}\mathbf{S}}_{\text{cancels}}\cdots\underbrace{\mathbf{S}^{-1}\mathbf{S}\mathbf{D}\mathbf{S}^{-1}\mathbf{S}}_{\text{cancels}}\cdots\underbrace{\mathbf{S}^{-1}\mathbf{S}\mathbf{D}\mathbf{S}^{-1}}_{\text{cancels}} = \mathbf{S}\mathbf{D}^n\mathbf{S}^{-1}.$$

6. $e^{\mathbf{A}t} = \mathbf{X}(t)\mathbf{X}(0)^{-1}$ for any fundamental matrix \mathbf{X} .

Proof: Both $e^{\mathbf{A}t}$ and $\mathbf{X}(t)\mathbf{X}(0)^{-1}$ satisfy the matrix differential equation $\dot{\mathbf{Y}} = \mathbf{A}\mathbf{Y}$ and the same initial conditions $\mathbf{Y}(0) = \mathbf{I}$. The uniqueness and existence theorem then guarantees these to be the same. (You can also think of the columns of $e^{\mathbf{A}t}$ separately and apply the existence and uniqueness theorem.)

Problem 15.1 Use the matrix exponential to find the solution to the system

$$\dot{x} = 2x + y$$

$$\dot{y} = 2y$$

satisfying $x(0) = 5$ and $y(0) = 7$.

Solution: This is $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x}$ with $\mathbf{A} = \begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}$. Write $\mathbf{A} = \mathbf{D} + \mathbf{N}$ with $\mathbf{D} = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$ (diagonal) and $\mathbf{N} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. Then $\mathbf{N}^2 = 0$, so

$$e^{\mathbf{N}t} = \mathbf{I} + \mathbf{N}t = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}.$$

Then $\mathbf{D}t$ and $\mathbf{N}t$ commute (a scalar times \mathbf{I} commutes with any matrix of the same size), so

$$\begin{aligned} e^{\mathbf{A}t} &= e^{\mathbf{D}t + \mathbf{N}t} \\ &= e^{\mathbf{D}t} e^{\mathbf{N}t} \\ &= \begin{pmatrix} e^{2t} & 0 \\ 0 & e^{2t} \end{pmatrix} \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} e^{2t} & te^{2t} \\ 0 & e^{2t} \end{pmatrix} \\ \begin{pmatrix} x \\ y \end{pmatrix} &= e^{\mathbf{A}t} \begin{pmatrix} 5 \\ 7 \end{pmatrix} \\ &= \begin{pmatrix} e^{2t} & te^{2t} \\ 0 & e^{2t} \end{pmatrix} \begin{pmatrix} 5 \\ 7 \end{pmatrix} \\ &= \begin{pmatrix} 5e^{2t} + 7te^{2t} \\ 7e^{2t} \end{pmatrix}. \end{aligned}$$

Remark: The matrix exponential is a fundamental matrix even when \mathbf{A} is deficient, as in the example above.

Compute the exponential matrix

0.5/1 point (graded)

Compute the exponential matrix $e^{\mathbf{A}}$ for $\mathbf{A} = \begin{pmatrix} -2 & 1 \\ 1 & -2 \end{pmatrix}$.

(Enter **[a,b;c,d]** for the matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$. That is, use commas to separate entries within the same row and **semicolon** to separate rows, and **square bracket** the entire matrix. Similarly for larger matrices.)

$$e^{\mathbf{A}} = [0.2088333, 2.7182818; 2.7182818, 0.2088333] *$$

Answer: $[\cosh(1)/e^2, \sinh(1)/e^2; \sinh(1)/e^2, \cosh(1)/e^2]$

Solution:

Since \mathbf{A} is symmetric, it is diagonalizable. Observe that

$$\mathbf{A} = -2\mathbf{I} + \mathbf{B} \quad \text{where } \mathbf{B} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

and

$$(-2\mathbf{I})\mathbf{B} = \mathbf{B}(-2\mathbf{I}).$$

This implies that $e^{\mathbf{A}t} = e^{-2t}\mathbf{I}e^{\mathbf{B}}$. Let us compute $e^{\mathbf{B}}$:


$$\begin{aligned} e^{\mathbf{B}} &= \mathbf{I} + \mathbf{B} + \frac{\mathbf{B}^2}{2} + \frac{\mathbf{B}^3}{3!} + \frac{\mathbf{B}^4}{4!} + \cdots & \left(\mathbf{B} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right) \\ &= \mathbf{I} + \mathbf{B} + \frac{\mathbf{I}}{2} + \frac{\mathbf{B}}{3!} + \frac{\mathbf{I}}{4!} + \cdots & \text{since } \mathbf{B}^2 = \mathbf{I} \\ &= \mathbf{I} \left(1 + \frac{1}{2} + \frac{1}{4!} + \cdots \right) + \mathbf{B} \left(1 + \frac{1}{3!} + \frac{1}{5!} + \cdots \right) \\ &= \mathbf{I} \cosh(1) + \mathbf{B} \sinh(1) = \begin{pmatrix} \cosh(1) & \sinh(1) \\ \sinh(1) & \cosh(1) \end{pmatrix}. \end{aligned}$$

Therefore,

$$e^{\mathbf{A}t} = e^{-2t} \begin{pmatrix} \cosh(1) & \sinh(1) \\ \sinh(1) & \cosh(1) \end{pmatrix} = e^{-2} \begin{pmatrix} \cosh(1) & \sinh(1) \\ \sinh(1) & \cosh(1) \end{pmatrix}.$$

Submit

You have used 3 of 3 attempts

 Answers are displayed within the problem

15. Properties of the matrix exponential

Hide Discussion

Topic: Unit 3: Solving systems of first order ODEs using matrix methods
/ 15. Properties of the matrix exponential

Add a Post

Show all posts ▼		by recent activity ▼	
?	<u>There is no t in the question</u> There is no t in the question.	2	▼
?	<u>Different solution.</u> Please note that in solving this problem, I used a different approach, namely exploiting this property $e^{A\cdot}$...	2	▼
💬	<u>Typo at answer box</u>	2	▼
?	<u>Grader Problem</u> I gave the numerical value of the correct solution in my three attempts and was marked wrong. Help! Th...	5	▼
💬	<u>BCH formula</u>	1	▼
💬	<u>Verify the grader</u> I think something is going wrong with the grader. If you try the matlab command <code>expm(A)</code> or calculating...	2	▼

Learn About Verified Certificates

© All Rights Reserved