

edX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Course](#) > [Unit 2 Foundation of Inference](#) > [Lecture 4: Parametric Estimation and Confidence Intervals](#) > 3. Bias of Estimators; Jensen's Inequality

### Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

## 3. Bias of Estimators; Jensen's Inequality

### Bias Estimators and an application of Jensen's Inequality

itself, which is precisely the definition of a convex

function.

The way you want to remember Jensen, and the way it goes,

is by saying the chord represents the expectation,



and of course, the function represents  
the function  
of the expectation.

End of transcript. Skip to the start.



## Video

[Download video file](#)

## Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

## The Expectation of the Average

1/1 point (graded)

Let  $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{U}([a, a+1])$  where  $a$  is an unknown parameter. Let  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$  denote the sample mean. In terms of  $a$ , what is  $\mathbb{E}[\bar{X}_n]$ ?

$\mathbb{E}[\bar{X}_n] =$   ✓ Answer: a+1/2

**Solution:**

Note that since the  $X_i$ 's are identically distributed, by linearity of expectation,

$$\mathbb{E}[\bar{X}_n] = \frac{1}{n} \sum_{i=1}^n \mathbb{E}[X_i] = \mathbb{E}[X_1] = a + \frac{1}{2}.$$

You have used 1 of 3 attempts

**i** Answers are displayed within the problem

## Computing Bias

1/1 point (graded)

**Recall:** Let  $\hat{\theta}_n$  denote an estimator for a true parameter  $\theta$ . Here  $n$  specifies the sample size. The **bias** of  $\hat{\theta}_n$  is defined to be

$$\mathbb{E}[\hat{\theta}_n] - \theta.$$

Let  $X_1, \dots, X_n$  be defined as in the previous question. Compute the bias of the estimator  $\bar{X}_n$  with respect to the parameter  $a$ .

✓ Answer: .5

**Solution:**

The bias is given by  $\mathbb{E}[\bar{X}_n] - a = 1/2$ , where we applied the previous part. Note that this implies that  $\bar{X}_n - \frac{1}{2}$  is an unbiased estimator.

Submit

You have used 1 of 3 attempts

**i** Answers are displayed within the problem

(Optional) Jensen's Inequality

Show

## (Optional) Expectation of nonlinear functions and Jensen's Inequality

0 points possible (ungraded)

Let  $X$  be a positive random variable with expectation  $\lambda$ . How does  $\mu = \mathbb{E} \left[ \frac{1}{X} \right]$  compare to  $\frac{1}{\lambda}$ ?

☐ In general,  $\mu$  and  $\lambda$  are not comparable

☒  $\mu \geq \frac{1}{\lambda}$

☐  $\mu \leq \frac{1}{\lambda}$



### Solution:

Note that the function  $x \mapsto \frac{1}{x}$  is a convex function on  $(0, \infty)$ , hence we can use Jensen's inequality that implies

$$\mathbb{E}[f(X)] \geq f(\mathbb{E}[X])$$

for all convex functions  $f$  to conclude

$$\mu = \mathbb{E} \left[ \frac{1}{X} \right] \geq \frac{1}{\mathbb{E}[X]} = \frac{1}{\lambda}.$$

[Submit](#)

You have used 1 of 1 attempt

**i** Answers are displayed within the problem

## Discussion

[Hide Discussion](#)

**Topic:** Unit 2 Foundation of Inference:Lecture 4: Parametric Estimation and Confidence Intervals  
/ 3. Bias of Estimators; Jensen's Inequality

[Add a Post](#)

Show all posts ▼

by recent activity ▼

☒ [computing bias](#)

I'm assuming  $\theta$  is  $\mu$  of a Uniform Random Variable?. If yes, then that estimator is unbiased and the answer wouldn't include "a" but since i...

6

[Learn About Verified Certificates](#)

© All Rights Reserved