EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the Privacy Policy.





Lecture 16: Goodness of Fit Tests Continued: Kolmogorov-Smirnov test, Kolmogorov-Lilliefors test,

Course > Unit 4 Hypothesis testing > Quantile-Quantile Plots

- 2. Review: Cumulative Distribution
- > Functions

2. Review: Cumulative Distribution Functions

Warmup: Integration limits of CDF

3/3 points (graded)

The cumulative distribution function $\Phi:\mathbb{R}\to\mathbb{R}$ of the standard normal $\mathcal{N}\left(0,1\right)$ can be written as

$$\Phi \left(z
ight) =\int_{A}^{B\left(z
ight) }rac{1}{\sqrt{2\pi }}e^{C\left(x
ight) }dx$$

where B(z) is a function of z and C(x) is a function of x. Write down the integration limits A, B(z), as well as the function C(x) in the integrand.

Enter inf for ∞ .

$$A = egin{bmatrix} -inf \ \hline -inf \end{bmatrix}$$
 $ullet$ Answer: -inf

2. Review: Cumulative Distribution Functions | Lecture 16: Goodness of Fit Tests Continued: Kolmogorov-Smirnov test, Kolmogorov-Lilliefors test, Quantile-Quantile Plots | 18.6501x Courseware | edX

STANDARD NOTATION

Solution:

Recall the cdf of a distribution ${\bf P}$ is a function F such that

$$F\left(x
ight) =P\left(X\leq x
ight) ,\quad X\sim \mathbf{P}.$$

The density of a standard Gaussian is given by $\phi\left(x
ight)=rac{1}{\sqrt{2\pi}}e^{-x^{2}/2}$. Therefore, the CDF of $\mathcal{N}\left(0,1
ight)$ is given by

$$\Phi \left(z
ight) =\int_{-\infty }^{z}rac{1}{\sqrt{2\pi }}e^{-x^{2}/2}\,dx.$$

Hence
$$A=-\infty$$
 , $B=z$, and $C=-x^2/2$.

Submit

You have used 2 of 3 attempts

1 Answers are displayed within the problem

Review



Video Download video file **Transcripts** Download SubRip (.srt) file Download Text (.txt) file

Review: Cumulative Distribution Function

1/1 point (graded)

2. Review: Cumulative Distribution Functions | Lecture 16: Goodness of Fit Tests Continued: Kolmogorov-Smirnov test, Kolmogorov-Lilliefors test, Quantile-Quantile Plots | 18.6501x Courseware | edX Let X and Y be real-valued random variables, both distributed according to a distribution \mathbf{P} . (We make no assumption about their joint distribution). Let F denote the **cdf** of \mathbf{P} .

Which of the following are true about the cdf F? (Choose all that apply.)

 $P(X \leq t)$ and $P(Y \leq t)$ are random variables.

ightharpoons For all $t\in\mathbb{R}$, $F\left(t
ight)=P\left(X\leq t
ight)$ and $F\left(t
ight)=P\left(Y\leq t
ight)$

 $\bigcap F(t) = P(X \le t) = P(Y \le t)$ only if X and Y are independent.

 $arprojlim_{t o\infty}F(t)=1$

 $\lim_{t\to-\infty}F(t)=0.$

 $\int F(t) dt = 1$



Solution:

We examine the choices in order.

- The first choice is incorrect. The probability of an event is a real number, not a random variables, so both ${f P}(X < t)$ and ${f P}(Y < t)$ are deterministic numbers.
- The second choice is correct. The joint distribution of X and Y is irrelevant– so long as X and Y have the same distribution, it will be true that $\mathbf{P}(X \le t) = \mathbf{P}(Y \le t)$. By definition, both of these quantities are equal to F(t).
- The third choice is incorrect. As stated in the previous bullet, regardless of the joint distribution of X and Y, as long as they are identically distributed, it is true that $\mathbf{P}(X \leq t) = \mathbf{P}(Y \leq t)$.

- 2. Review: Cumulative Distribution Functions | Lecture 16: Goodness of Fit Tests Continued: Kolmogorov-Smirnov test, Kolmogorov-Lilliefors test, Quantile-Quantile Plots | 18.6501x Courseware | edX
 - The fourth choice is correct. Observe that $\mathbf{P}(X \leq \infty) = 1$ because X is real-valued. Moreover, F is an increasing function of t. Therefore, $\lim_{t \to \infty} F(t) = \mathbf{P}(X \leq \infty) = 1$.
 - ullet The fifth choice is correct, since $\lim_{t o -\infty}F\left(t
 ight) =\mathbf{P}\left(X\leq -\infty
 ight) =0.$
 - The final choice is incorrect. The statement is true for the **pdf** not the cdf:

$$\int_{-\infty}^{\infty}f\left(t
ight) dt=1\qquad ext{if }f\left(t
ight) ext{ is pdf of }X.$$

Since the cdf F(t) has limit $\lim_{t \to \infty} F(t) = 1$, the integral over $\mathbb R$ of F(t) diverges.

Submit

You have used 1 of 2 attempts

• Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 4 Hypothesis testing:Lecture 16: Goodness of Fit Tests Continued: Kolmogorov-Smirnov test, Kolmogorov-Lilliefors test, Quantile-Quantile Plots / 2. Review: Cumulative Distribution Functions

Add a Post

Show all posts ▼

Show all posts ▼

Goodness of fit test goes against the grain of some of the principles of Hypothesis testing learnt thus far?

8 new_ 13

© All Rights Reserved