EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the Privacy Policy.





<u>Unit 0. Course Overview, Syllabus,</u> <u>Guidelines, and Homework on</u>

Homework 0: Probability and Linear

> <u>algebra Review</u>

> 3. Gaussian random variables

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. **Upgrade now**

3. Gaussian random variables

Moments of Gaussian random variables

5/5 points (graded)

Course > Prerequisites

Let X be a Gaussian random variable with mean μ and variance σ^2 . Compute the following moments:

Remember that we use the terms **Gaussian random variable** and **normal random variable** interchangeably.

(Enter your answers in terms of μ and σ .)

$$\mathbb{E}\left[X^3\right] = \boxed{\begin{array}{c} \text{mu}^3 + 3 \text{*mu*sigma}^2 \\ \hline \mu^3 + 3 \cdot \mu \cdot \sigma^2 \end{array}} \qquad \qquad \textbf{Answer: 3 * sigma}^2 \text{*mu + mu}^3$$

$$\mathbb{E}\left[X^{4}\right] = \boxed{ \begin{array}{c} \text{mu}^{4} + 6 * \mu^{2} \cdot \sigma^{2} + 3 \cdot \sigma^{4} \end{array}} \qquad \qquad \textbf{Answer: 3 * sigma}^{4} + 6 * sigma}^{2} \qquad \qquad \textbf{Answer: 3 * sigma}^{4} + 6 * sigma}^{2} = \boxed{ \begin{array}{c} \mu^{4} + 6 \cdot \mu^{2} \cdot \sigma^{2} + 3 \cdot \sigma^{4} \end{array} }$$

Write ${f P}(X>0)$ in terms of the **cumulative distribution function (cdf)** Φ of the standard Gaussian distribution, that is,

$$\Phi\left(x
ight) =\mathbf{P}\left(Z\leq x
ight) ,\quad x\in\mathbb{R},$$

where $Z \sim \mathcal{N}\left(0,1
ight)$ is a standard normal variable. (Enter Phi for Φ .)

STANDARD NOTATION

Solution:

We can write a general Gaussian variable $X \sim \mathcal{N}\left(\mu,\sigma^2\right)$ as $X = \sigma Z + \mu$, where $Z \sim \mathcal{N}\left(0,1\right)$ is a standard normal variable. Hence, the calculation can be made by factoring out the corresponding polynomials and calculating (or looking up) the moments of Z:

$$egin{array}{ll} \mathbb{E}\left[Z
ight] = & 0 \ \mathbb{E}\left[Z^2
ight] = & 1 \ \mathbb{E}\left[Z^3
ight] = & 0 \ \mathbb{E}\left[Z^4
ight] = & 3. \end{array}$$

As an example, let us compute $\mathbb{E}\left[X^3
ight]$. Denote the density of a standard normal distribution by $arphi\left(z
ight)$, i.e.,

$$\phi\left(z
ight)=rac{1}{\sqrt{2\pi}}e^{-rac{z^{2}}{2}}.$$

With this, we calculate

$$egin{array}{lll} \mathbb{E}\left[X^3
ight] &=& \int_{-\infty}^{\infty} \left(\sigma z + \mu
ight)^3 \phi\left(z
ight) dz \ \\ &=& \sigma^3 \mathbb{E}\left[Z^3
ight] + 3\sigma^2 \mu \mathbb{E}\left[Z^2
ight] + 3\sigma \mu^2 \mathbb{E}\left[Z
ight] + \mu^3 \ \\ &=& 3\sigma^2 \mu + \mu^3. \end{array}$$

For $\mathsf{Var}(X^2)$, we can use the formula $\,\mathsf{Var}(X^2) = \mathbb{E}\,[X^4] - \left(\mathbb{E}\,[X^2]\right)^2$.

Similarly, we can express the probability $\mathbf{P}\left(X>0
ight)$ as

$$\mathbf{P}\left(X>0
ight) = \mathbf{P}\left(\sigma Z + \mu > 0
ight) = \mathbf{P}\left(\sigma Z > -\mu
ight) \ = \mathbf{P}\left(Z>-rac{\mu}{\sigma}
ight) = 1 - \Phi\left(-rac{\mu}{\sigma}
ight).$$

Submit

You have used 1 of 4 attempts

1 Answers are displayed within the problem

Covariance of Gaussians

4/4 points (graded)

Recall that **i.i.d.** stands for **independent and identically distributed**. A collection of random variables X_1, \ldots, X_n are **i.i.d.** if all of them follow the same distribution, and each X_i does not contain information about the other realizations.

Let X,Y be i.i.d. **standard** normal random variables, that is, $X,Y \sim \mathcal{N}\left(0,1
ight)$.

Recall that the **covariance** of two random variables X and Y, denoted by $\mathsf{Cov}(X,Y)$, is defined as

$$\mathsf{Cov}\left(X,Y\right) \; = \; \mathbb{E}\left[\left(X-\mathbb{E}\left[X\right]\right)\left(Y-\mathbb{E}\left[Y\right]\right)\right]. \tag{1.3}$$

Compute the following variances and covariances.

$$Var(X+Y)=$$
 2 Answer: 2

STANDARD NOTATION

Solution:

Note that by the definition of a standard Gaussian random variable,

$$\mathbb{E}\left[X
ight] = \mathbb{E}\left[Y
ight] = 0 \quad \mathbb{E}\left[X^2
ight] = \mathbb{E}\left[Y^2
ight] = 1.$$

With this, compute

$$\mathsf{Var}(X+Y) = \mathsf{Var}(X) + \mathsf{Var}(Y)$$
 (independence)
= 1+1=2,

$$egin{array}{ll} (XY) &=& \mathbb{E}\left[(XY)^2
ight] - (\mathbb{E}\left[XY
ight])^2 \ &=& \mathbb{E}\left[X^2
ight]\mathbb{E}\left[Y^2
ight] - \mathbb{E}[X]^2\mathbb{E}[Y]^2 \end{array} \qquad \qquad ext{(independence)}$$

$$= 1 \times 1 - 0 = 1,$$

$$(X, X + Y) = \mathbb{E}[X(X + Y)] - \mathbb{E}[X]\mathbb{E}[X + Y]$$

$$= \mathbb{E}[X^2] + \mathbb{E}[XY] - \mathbb{E}[X](\mathbb{E}[X] + \mathbb{E}[Y]) \qquad \text{(linearity of expectation)}$$

$$= \mathbb{E}[X^2] + \mathbb{E}[X]\mathbb{E}[Y] - \mathbb{E}[X]^2 - \mathbb{E}[X]\mathbb{E}[Y] \qquad \text{(independence)}$$

$$= 1,$$

$$(X, XY) = \mathbb{E}[X(XY)] - \mathbb{E}[X]\mathbb{E}[XY]$$

$$= \mathbb{E}[X^2]\mathbb{E}[Y] - \mathbb{E}[X]^2\mathbb{E}[Y] \qquad \text{(independence)}$$

$$= 1 \cdot 0 - 0 \cdot 0 = 0.$$

: 8. Covariance, 9. Covariance properties, and 10. the variance of a sum in Lecture 12, *Sums of independent random variables; covariance, and correlation*.

Submit

You have used 1 of 3 attempts

• Answers are displayed within the problem

True or False: Variance, covariance and independence

2/2 points (graded)

For each of the statements below, determine whether it is true (meaning, always true) or false (meaning, not always true).

ullet For any two random variables, $\operatorname{\sf Var}(X+Y)=\operatorname{\sf Var}(X)+\operatorname{\sf Var}(Y).$

True

● False

• If the covariance, Cov(X,Y) between two random variables X,Y is 0, then X and Y are independent.

True

False

STANDARD NOTATION

Solution:

• The first item is False. For any two random variables, it is known that,

$$Var(X + Y) = Var(X) + Var(Y) + 2Cov(X, Y).$$

In particular, if $\mathrm{Cov}\left(X,Y\right) \neq 0$, this does not hold.

ullet The second item is also false. As a simple example, let $X\sim \mathrm{Unif}\,[-1,1]$ and let $Y=X^2$. Then,

$$\mathrm{Cov}\left(X,Y
ight)=\mathbb{E}\left[XY
ight]-\mathbb{E}\left[X
ight]\mathbb{E}\left[Y
ight]=\mathbb{E}\left[X^3
ight]-\mathbb{E}\left[X
ight]\mathbb{E}\left[X^2
ight]=0,$$

using the fact that X is centered and symmetric around 0, and its odd moments vanish. Even though they are uncorrelated, they are (highly) dependent, Y is obtained from X, intuitively!

Submit

You have used 1 of 3 attempts

1 Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 0. Course Overview, Syllabus, Guidelines, and Homework on Prerequisites:Homework 0: Probability and Linear algebra Review / 3. Gaussian random variables

Add a Post

Show all posts ▼

Book on statistics
I have not taken the recommended earlier course mentioned as prerequisite for this course. Can someone recommend any book followed in that ea... 7

I Pinned

Covariance of Gaussians

For those who are overwhelmed with the first problem (like myself)

30

? Struggling with this section questions Covariance of Gaussians, a hint will be really helpful	2
? Two dependent random variables covariance=0 example Are there any examples when Cov(X,Y) between two random variables X,Y is 0 and they are dependent? I can't come up with the one. Any ideas?	4
Last part of first question Lam stuck on the last part of the first question: "Write P(X>0) in terms of the cumulative distribution function (cdf) Φ of the standard Gaussian	trib
✓ Notation Bug? on last part of Moments of Gaussian random variables question Is there something wrong with the standard notation Phi(0) to denote the normal cumulative distribution function at zero?	6
Where did you learn this "moments" thing? I managed to get the correct answers after some googling and tedious calculations but I have never heard of the "moments" until now. It wasn't tage.	14 aug
 Exercise 1 Hi guys I am unsure as to why on the last part of the question when it asks you to write P(x>0) as a Cumulative distribution function the answer is 	3 no
? [Staff] First part N/A in the mobile app I almost missed the 5 point worth first part of this question. It doesn't appear in the mobile app. Could you check please? Thank you in advance	1
• [Staff] [Typo]	3
Format for question 4 Liget: Invalid Input: X, Y not permitted in answer as a variable. Without revealing answers, is there an issue with putting answers as their function for	<u>2</u>

Learn About Verified Certificates

© All Rights Reserved