



[Lecture 14: Wald's Test, Likelihood
Ratio Test, and Implicit Hypothesis](#)

[Course](#) > [Unit 4 Hypothesis testing](#) > [Test](#)

> 6. Wald's Test Continued

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

6. Wald's Test Continued

Review: Chi-Squared Distribution

2/2 points (graded)

Which of the following random variables follow a χ_d^2 distribution?

(Choose all that apply. In the choices, "I" denotes the $d \times d$ identity matrix.)

☐ $Z_1 + Z_2 \dots + Z_d$ where $Z_i \sim \mathcal{N}(\mu, \sigma^2)$

☐ $Z_1^2 + Z_2^2 \dots + Z_d^2$ where $Z_i \sim \mathcal{N}(\mu, \sigma^2)$ for some $\mu, \sigma \in \mathbb{R}$

☐ $Z_1^2 + Z_2^2 \dots + Z_d^2$ where $Z_i \sim \mathcal{N}(\mu, \sigma^2)$ for some $\mu, \sigma \in \mathbb{R}$ and are independent

☐ $Z_1 + Z_2 \dots + Z_d$ where $Z_i \sim \mathcal{N}(0, 1)$

☐ $Z_1^2 + Z_2^2 \dots + Z_d^2$ where $Z_i \sim \mathcal{N}(0, 1)$

☒ $Z_1^2 + Z_2^2 \dots + Z_d^2$ where $Z_i \sim \mathcal{N}(0, 1)$ and are independent



☐ $\|\mathbf{Z}\|$ where \mathbf{Z} is multivariate Gaussian $\mathbf{Z} \sim \mathcal{N}_d(\vec{\mu}, \Sigma_{\mathbf{Z}})$ for some $\vec{\mu} \in \mathbb{R}^d$ and $d \times d$ matrix $\Sigma_{\mathbf{Z}}$

☐ $\|\mathbf{Z}\|$ where \mathbf{Z} is multivariate Gaussian $\mathbf{Z} \sim \mathcal{N}_d(\vec{\mu}, I)$ for some $\vec{\mu} \in \mathbb{R}^d$

☐ $\|\mathbf{Z}\|$ where \mathbf{Z} is multivariate Gaussian $\mathbf{Z} \sim \mathcal{N}_d(\mathbf{0}, I)$

☐ $\|\mathbf{Z}\|^2$ where \mathbf{Z} is multivariate Gaussian $\mathbf{Z} \sim \mathcal{N}_d(\vec{\mu}, \Sigma_{\mathbf{Z}})$ for some $\vec{\mu} \in \mathbb{R}^d$ and $d \times d$ matrix $\Sigma_{\mathbf{Z}}$

☐ $\|\mathbf{Z}\|^2$ where \mathbf{Z} is multivariate Gaussian $\mathbf{Z} \sim \mathcal{N}_d(\vec{\mu}, I)$ for some $\vec{\mu} \in \mathbb{R}^d$

☒ $\|\mathbf{Z}\|^2$ where \mathbf{Z} is multivariate Gaussian $\mathbf{Z} \sim \mathcal{N}_d(\mathbf{0}, I)$



Solution:

The χ^2 distribution with d degrees of freedom is by definition the distribution of

$$Z_1^2 + Z_2^2 \dots + Z_d^2 \quad \text{where } Z_i \stackrel{iid}{\sim} \mathcal{N}(0, 1)$$

or equivalently the distribution of

$$\|\mathbf{Z}\|^2 \quad \text{where } \mathbf{Z} \sim \mathcal{N}_d(\mathbf{0}, \mathbf{1}),$$

whose components are independent because the off-diagonal elements of the covariance matrix $\mathbf{1}$ are all 0.

Remark: Recall from a problem on the previous page that the vector $\mathbf{M}\mathbf{Z}$, where $\mathbf{M}^T = \mathbf{M}^{-1}$ (or equivalently $\mathbf{M}\mathbf{M}^T = \mathbf{M}^T\mathbf{M} = \mathbf{1}_{d \times d}$), is also a **standard** multivariate Gaussian vector. Hence $\|\mathbf{M}\mathbf{Z}\|^2$ also follows a χ_d^2 distribution.

Submit

You have used 2 of 3 attempts

i Answers are displayed within the problem

Review: Writing the Norm Squared

1/1 point (graded)

Which of the following equals the squared norm $\|\mathbf{Ax}\|^2$ of the vector \mathbf{Ax} , where \mathbf{A} is a **symmetric** $d \times d$ matrix and \mathbf{x} is a vector in \mathbb{R}^d ?

(Choose all that apply.)

☒ $(\mathbf{Ax})^T (\mathbf{Ax})$

☐ $(\mathbf{Ax}) (\mathbf{Ax})^T$

☒ $\mathbf{x}^T \mathbf{A}^T \mathbf{Ax}$

☒ $\mathbf{x}^T \mathbf{A}^2 \mathbf{x}$



Solution:


$$\begin{aligned}\|\mathbf{Ax}\|^2 &= (\mathbf{Ax})^T (\mathbf{Ax}) = \mathbf{x}^T \mathbf{A}^T \mathbf{Ax} \\ &= \mathbf{x}^T \mathbf{A}^T \mathbf{Ax} \quad (\text{since } \mathbf{A}^T = \mathbf{A}) = \mathbf{x}^T \mathbf{A}^2 \mathbf{x}\end{aligned}$$

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Wald's Test Continued



Video

[Download video file](#)

Transcripts

[Download SubRip\(.srt\) file](#)

[Download Text \(.txt\) file](#)

Deriving Wald's Test

1/1 point (graded)

Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathbf{P}_{\theta^*}$ for some true parameter $\theta^* \in \mathbb{R}^d$. We construct the associated statistical model $(\mathbb{R}, \{\mathbf{P}_{\theta}\}_{\theta \in \mathbb{R}^d})$ and the maximum likelihood estimator $\hat{\theta}_n^{MLE}$ for θ^* .

Your goal is to use hypothesis testing to decide between two hypotheses:

$$H_0 : \theta^* = \mathbf{0}$$

$$H_1 : \theta^* \neq \mathbf{0}.$$

Assuming that the null hypothesis is true, the asymptotic normality of the MLE $\hat{\theta}_n^{MLE}$ implies that the following random variable

$$\left\| \sqrt{n} \mathcal{I}(\mathbf{0})^{1/2} (\hat{\theta}_n^{MLE} - \mathbf{0}) \right\|^2$$

converges to a χ_k^2 distribution. What is the degree of freedom k of this χ_k^2 distribution?

$$\left\| \sqrt{n} \mathcal{I}(\mathbf{0})^{1/2} (\hat{\theta}_n^{MLE} - \mathbf{0}) \right\|^2 \xrightarrow[n \rightarrow \infty]{(d)} \chi_k^2 \text{ for } k =$$

d

✓ Answer: d

STANDARD NOTATION

Solution:

From the previous problem, we know that under the assumption $X_1, \dots, X_n \stackrel{iid}{\sim} P_{\mathbf{0}}$,

$$\sqrt{n} \mathcal{I}(\mathbf{0})^{1/2} (\hat{\theta}_n^{MLE} - \mathbf{0}) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(\mathbf{0}, I_{d \times d}).$$

Next, if $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, I_{d \times d})$, then $Z_1, \dots, Z_d \stackrel{iid}{\sim} \mathcal{N}(0, 1)$. Hence,

$$\|\mathbf{Z}\|_2^2 = Z_1^2 + Z_2^2 + \cdots + Z_d^2 \sim \chi_d^2$$

by definition of the χ^2 distribution with d degrees of freedom. Hence by continuity, we have

$$\left\| \sqrt{n} \mathcal{I}(\mathbf{0})^{1/2} (\hat{\theta}_n^{MLE} - \mathbf{0}) \right\|_2^2 \xrightarrow[n \rightarrow \infty]{(d)} \chi_d^2.$$

Remark: The above allows us to derive **Wald's test**. For the given null and alternative hypotheses:

$$H_0 : \theta^* = \mathbf{0}$$

$$H_1 : \theta^* \neq \mathbf{0},$$

we define the test statistic

$$W_n := \left\| \sqrt{n} \mathcal{I}(\mathbf{0})^{1/2} (\hat{\theta}_n^{MLE} - \mathbf{0}) \right\|^2 = n (\hat{\theta}_n^{MLE} - \mathbf{0})^T \mathcal{I}(\mathbf{0}) (\hat{\theta}_n^{MLE} - \mathbf{0}).$$

Then, then Wald's test of level α is the test

$$\psi_\alpha = \mathbf{1}(W_n > q_\alpha(\chi_d^2)),$$

where $q_\alpha(\chi_d^2)$ is the $1 - \alpha$ -quantile of the (pivotal) distribution χ_d^2 .

Submit

You have used 1 of 2 attempts

i Answers are displayed within the problem

Discussion

[Hide Discussion](#)

Topic: Unit 4 Hypothesis testing:Lecture 14: Wald's Test, Likelihood Ratio Test, and Implicit Hypothesis Test / 6.
Wald's Test Continued

[Add a Post](#)

Show all posts ▼

by recent activity ▼

? [Are we going back to asymptotic distributions?](#)

1 ▼

We needed chi-squared and T test so that we can say something when the sample size is small. Since, we are going back to asymptotic distributions, does that mean there is ...

? [Review: Chi-Squared Distribution](#)

3 ▼

Quick question, as I am confused about the wording: ' $Z_1 + Z_2 + \dots + Z_d$ where $Z_i \sim N(0,1)$ and are independent' is a ~~Deleted by MW-CTA~~, but ' $Z_1 + Z_2 + \dots + Z_d$ where $Z_i \sim N(\mu, \dots$

[Learn About Verified Certificates](#)

© All Rights Reserved