



You are taking "[Exam \(Timed, No Correctness Feedback\)](#)," as a timed exam. [Show more](#)

End My Exam

44:21:15



< Previous



5. Scalar multiplication

Bookmark this page

Calculator



Hide Notes


Lecture due Aug 18, 2021 20:30 IST Completed



Practice

Scalar multiplication with a vector

[Start of transcript. Skip to the end.](#)



PROFESSOR: Another thing that we can do with vectors is we can scale them. We can stretch them or contract them. So if we write something like 2 times v, what it means is it's a vector in the same direction as v,

▶

0:00 / 0:00

▶

2.0x

🔊

🔍

📺

🗣️


Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)
[Download Text \(.txt\) file](#)

We can scale vectors. For example, $2\vec{v}$ is in the same direction as \vec{v} but is twice as long.



Definition 5.1

A **scalar** is a (real or complex) number. In this class, we will almost always consider real scalars.

To multiply a vector $\vec{v} = \langle v_1, v_2 \rangle$ by a scalar c , we multiply each component by c as follows:

$$c\vec{v} = c\langle v_1, v_2 \rangle = \langle cv_1, cv_2 \rangle.$$

🧮

Calculator

📝

Hide Notes

- If \vec{v} is in the **same direction** as \vec{w} , then $\vec{v} = \lambda \vec{w}$ for some **positive number** $\lambda > 0$.
- If \vec{v} is in the **opposite direction** as \vec{w} , then $\vec{v} = \lambda \vec{w}$ for some **negative number** $\lambda < 0$.

Question: Is $(2, 3)$ in the same direction as the vector $(4, 7)$?

This is the same as asking is $(4, 7) = \lambda (2, 3)$ for some positive λ ? For this to be true, we would need


$$(4, 7) = \lambda (2, 3) = (2\lambda, 3\lambda) \implies 4 = 2\lambda \text{ and } 7 = 3\lambda.$$

But this would mean $\lambda = 2$ and $\lambda = 7/3$, which is not possible. Since there is no λ that works, the vectors are not in the same direction.

On the other hand,

$$(4, 6) = 2 (2, 3)$$

and so $(4, 6)$ is in the same direction as $(2, 3)$.



▼ Spoiler: Scalar multiplication in higher dimension.

Consider a scalar c and a vector with n components given by $\vec{v} = \langle v_1, v_2, \dots, v_n \rangle$. Then $c\vec{v}$ is given by

$$c\vec{v} = c\langle v_1, v_2, \dots, v_n \rangle = \langle cv_1, cv_2, \dots, cv_n \rangle. \tag{3.7}$$

[Hide](#)

Scalar multiplication and graders

0 points possible (ungraded)
This problem is to help familiarize yourself with entering vectors or their scalar multiples into an automatically graded problem. It is ungraded.

Given a vector \vec{v} and a scalar c , enter $c\vec{v}$ for the following values.

(Enter your answer as a vector with two components inside square brackets, e.g., [1,1]. Note you can type 2*[1,1] for the vector $\langle 2, 2 \rangle$.)

$\vec{v} = \langle 3, 4 \rangle, c = 3,$

$c\vec{v} =$

✓ Answer: [9,12]

Calculator

Hide Notes

$\vec{w} = \langle 2, -8 \rangle, c = -7,$

✓ Answer: [-14,56]

$\vec{u} = \langle -32, 24 \rangle, c = -1/4, c\vec{u} =$

✓ Answer: [8,-6]

Solution:

We have:

$3\langle 3, 4 \rangle = \langle (3)(3), (3)(4) \rangle = \langle 9, 12 \rangle$

(3.8)

$-7\langle 2, -8 \rangle = \langle (-7)(2), (-7)(-8) \rangle = \langle -14, 56 \rangle$

(3.9)

$-\frac{1}{4}\langle -32, 24 \rangle = \langle (-1/4)(-32), (-1/4)(24) \rangle = \langle 8, -6 \rangle$

(3.10)

Submit

You have used 1 of 3 attempts

Answers are displayed within the problem

Why are they called scalars?

2.0/2 points (graded)

When we do math with vectors, we call real numbers scalars because when we multiply vectors by scalars, it has the effect of scaling the length of the vector.

Find the length of the vector $\langle 3, 4 \rangle$.

Find the length of the vector $c\langle 3, 4 \rangle = \langle 3c, 4c \rangle$, where c is a scalar.

(Hint: c could be positive, negative, or zero. For help in entering formulas, use the input help button.)

$|\langle 3, 4 \rangle| =$

✓ Answer: 5

$|\langle 3c, 4c \rangle| =$

✓ Answer: 5*abs(c)

? INPUT HELP

Solution:

$|\langle 3, 4 \rangle| = \sqrt{3^2 + 4^2} = 5$

$|\langle 3c, 4c \rangle| = \sqrt{3^2c^2 + 4^2c^2} = |c|\sqrt{3^2 + 4^2} = 5|c|$

Submit

You have used 2 of 3 attempts

Answers are displayed within the problem

Sketch the vectors

1.0/1 point (graded)


Let $\vec{v} = \langle -1, 2 \rangle$ and $c = -1/2$. Sketch the original vector \vec{v} in blue, and sketch the vector after scalar multiplication $c\vec{v}$ in orange.

You can move a drawn arrow around the page using the select button an clicking on the arrow to drag

Calculator

Hide Notes

drawn arrow, click select and then click and drag either the start or end point of the vector.



Answer: See solution.


Great job!

Solution:

The vector $\vec{v} = \langle -1, 2 \rangle$ begins at the origin and extends to the coordinate $(-1, 2)$. The vector

$$c\vec{v} = \frac{-1}{2}\langle -1, 2 \rangle = \langle 1/2, -1 \rangle \tag{3.11}$$

starts at the origin and extends to the coordinate $(1/2, -1)$.



Submit You have used 1 of 25 attempts

i Answers are displayed within the problem

5. Scalar multiplication

Hide Discussion

Topic: Unit 2: Geometry of Derivatives / 5. Scalar multiplication

Add a Post

Show all posts	by recent activity
[Staff] HTML Problem	7
Sketching the Vectors	4
assumption for scaling vectors We are told in the lecture that if vector v is in the same direction as vector w , that means we can multiply vector w with...	3
[Edit] Relabeling vector from (3,7) to (4,7) Hello, in the second diagram from the top of this page, the orange vector is labelled (3,7), but the vector discussed in text before it i...	2

< Previous Next >



edX

- About
- Affiliates
- edX for Business
- Open edX
- Careers
- News

Legal

- Terms of Service & Honor Code
- Privacy Policy
- Accessibility Policy
- Trademark Policy
- Sitemap

Connect

- Blog
- Contact Us
- Help Center
- Media Kit

Calculator

Hide Notes

[Donate](#)



© 2021 edX Inc. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)